5th INTERNATIONAL TOKAT SCIENTIFIC RESEARCH CONGRESS

SCIENCE • SOCIETY • SUSTAINABLE FUTURE

PROCEEDINGS BOOK

Editör

Dr. Ulduz Farhad Qahramanova

Tokat

TARİHİN DOĞANIN VE BİLİMİN BULUŞTUĞU ŞEHİR

Tokat

A City Where History, Nature and Science Meet

ISBN: 978-625-90699-1-3

5th INTERNATIONAL TOKAT SCIENTIFIC RESEARCH CONGRESS

August 7-9, 2026 / Tokat, TÜRKİYE

INTERNATIONAL SCIENTIFIC RESEARCH AND STRATEGY DEVELOPMENT ASSOCIATION

UBS

www.ubsder.org

PROCEEDINGS BOOK

EDITOR IN CHIEF

Dr. Ulduz Fərhad Qəhrəmanova

ISBN: 978-625-90699-1-3

by

ACADEMIC SHARING PLATFORM PUBLISHING HOUSE®

All Rights Reserved.
The authors are solely responsible for the ethical and legal content of their contributions.
APP Publications – 2026 ©

Publication Date: 20.09.2026

CONGRESS ID

CONGRESS TITLE

5th INTERNATIONAL
TOKAT SCIENTIFIC RESEARCH CONGRESS

DATE and PLACE

August 7-9, 2026 / Tokat, TÜRKİYE

ORGANIZATION

UBS-Institute of International Scientific Research and Strategy Development Association, Akademik Paylaşım Platformu Eğitim Ltd.Şti.,Ankara,Türkiye

CONGRESS CHAIRS

Dr. Ulduz Fərhad Qəhrəmanova

EDITOR IN CHIEF

Dr. Ulduz Fərhad Qəhrəmanova

CONGRESS ORGANIZERS

Uluslararası Bilimsel Araştırma ve Strateji Geliştirme Derneği, Türkiye

PARTICIPANT COUNTRIES (6):

Azerbaijan, Nigeria, Thailand, Türkiye, United Kingdom, Uzbekistan

NUMBER OF ACCEPTED PAPERS (70):

(Türkiye-25, Other Countries-45)

PRESENTATION

Oral presentation

EVALUATION PROCESS

All applications have undergone a double-blind peer review process

ISBN: 978-625-90699-1-3

ORGANIZING COMMITTEE

Prof. Dr. Hakkı ÇİFTÇİ – Çukurova University

Prof. Dr. Mehmet Salih SAYAR – Dicle University

Prof. Dr. Hasan TUTAR – Bolu Abant İzzet Baysal University

Prof. Dr. Ahmet DEVECİ – Gaziantep University

Prof. Dr. Sibel AKOVA – Yalova University

Prof. Dr. Mesut BULUT – Atatürk University

Prof. Dr. Abdulkadir KIRBAŞ – Atatürk University

Assoc. Prof. Dr. Münevver BAYAR – Bilim University

Assoc. Prof. Dr. Muhammad Zia-ur-Rehman – University of Malaya & NDU

Assoc. Prof. Dr. Fatma KÖPRÜLÜ – Near East University

Assoc. Prof. Dr. Melda Medine SUNAY – Bursa Technical University

Assoc. Prof. Dr. Günel HUMMETOVA – Azerbaijan State Pedagogical University

Assist. Prof. Dr. Gönül GÜL – Çankırı Karatekin University

Dr. Zeynep YILMAZ ŞİMŞEK – Bingöl University

Dr. Ulduz Farhad Qahramanova – Azerbaijan State Pedagogical University

Assist. Prof. Dr. Ananda MAJUMDAR – USA

Dr. Keshab Chandra MANDAL – India

Lect. Ayla DEVECİ – Kilis 7 Aralık University

INTERNATIONAL SCIENTIFIC RESEARCH AND STRATEGY DEVELOPMENT ASSOCIATION
www.ubsder.org

SCIENTIFIC BOARD MEMBERS

Prof. Dr. Yaşar Hamza OCAK – Kütahya Dumlupınar University – Türkiye

Prof. Dr. Sinem SOMUNOĞLU İKİNCİ – Bursa Uludağ University – Türkiye

Prof. Dr. Anderj JEAN – Gas Petroleum University of Romania – Romania

Prof. Dr. Müjdat AVCI – Osmaniye Korkut Ata University – Türkiye

Prof. Dr. Basim ALHADİDİ – Albalqa Applied University – Jordan

Prof. Dr. Ashraf M. A. AHMAD – Princess Sumaya University for Technology, Amman – Jordan

Prof. Dr. Raymond KOSALA – The Joseph Wibowo Center for Advanced Learning, Jakarta – Indonesia

Prof. Dr. Wen-Lin YANG – National University of Tainan – Taiwan

Prof. Dr. Gülzar İBRAHİMOVA – Baku Eurasian University – Azerbaijan

Assoc. Prof. Dr. M. Can YARDIMCI – Afyon Kocatepe University, Afyon – Türkiye

Assoc. Prof. Dr. Muhammad Zia-ur-Rehman – Universiti Malaya (Malaysia) & National Defence University (Pakistan) – Malaysia / Pakistan

Assoc. Prof. Dr. Aytekin ZEYNALOVA – Baku State University – Azerbaijan

Assoc. Prof. Dr. Sibel AKOVA – Yalova University – Türkiye

Assoc. Prof. Dr. Fahri ÖZSUNGUR – Mersin University – Türkiye

Assoc. Prof. Dr. Gönül SAMEDOVA – Azerbaijan State Pedagogy University – Azerbaijan

Assist. Prof. Dr. Abdullah Yüksel BARUT – İstanbul Gelişim University, İstanbul – Türkiye

Assist. Prof. Dr. Hayri YILDIRIM – Dicle University, Diyarbakır – Türkiye

Assist. Prof. Dr. Michal Wojtaszek – Warsaw University of Life Sciences – Poland

Assist. Prof. Dr. Maral Jamalova – Szent Istvan University – Hungary

Assist. Prof. Dr. Nicholas Imbeah – Takoradi Technical University – Ghana

Assist. Prof. Dr. Subhashish DEY – Gudlavalleru Engineering College, Civil Engineering Department, Andhra Pradesh – India

Assist. Prof. Dr. Ananda MAJUMDAR – North Central Theological Seminary, Minneapolis – USA

Assist. Prof. Dr. Subhashish DEY – Civil Engineering Department, Gudlavalleru Engineering College, Andhra Pradesh – India

UBSDER

Çankaya - Ankara

06-139-180

Konu: Kongre Düzenleme Kurulu 25.05.2026

Sayı:2026/210

İLGİLİ KURUMA

İçişleri Bakanlığı tarafından tahsis edilen 06-139-180 tescil kodu ile Tüzel kişiliğe sahip olan Uluslararası Bilimsel Araştırma ve Strateji Geliştirme Derneği, 5253 sayılı kanuna uygun olarak "Bilimsel Araştırmalar ve Akademik Çalışmalar" alanında ulusal ve uluslararası düzeyde faaliyetlerini yürütmektedir.

Kurumumuzun Yönetim Kurulu tarafından, 25.05.2026 tarihinde saat 14:00'da "Bilimsel Çalışmalarda Strateji Geliştirme" görüşmeleri ile "Bilimsel Kongreler Düzenlenmesi" gündemleri ile toplanmış ve alınan 9 nolu karara istinaden aşağıda detayları yazılı olan bilimsel etkinliğin düzenlenmesine ve etkinliğe ilişkin akademisyenlerin resmi olarak görevlendirilmesine oy birliği ile karar verilmiştir.

Bilgi ve gereğini arz ederim.

Dr. Nadire KANTARCIOĞLU

UBSDER Başkanı

N. Kantarcıoğlu

Etkinlik Adı : 5. ULUSLARARASI TOKAT BİLİMSEL ARAŞTIRMALAR KONGRESİ

Etkinlik Tarihi: 7-9 Ağustos 2026, Tokat/TÜRKİYE

www.ubsder.org

KONGRE BAŞKANI

Dr. Ulduz Farhad Qahramanova – Azerbaijan State Pedagogical University

Düzenleme Kurulu Üyeleri

Prof. Dr. Hakkı ÇİFTÇİ – Çukurova University

Prof. Dr. Mehmet Salih SAYAR – Dicle University

Prof. Dr. Hasan TUTAR – Bolu Abant İzzet Baysal University

Prof. Dr. Ahmet DEVECİ – Gaziantep University

Prof. Dr. Sibel AKOVA – Yalova University

Prof. Dr. Mesut BULUT – Atatürk University

Prof. Dr. Abdulkadir KIRBAŞ – Atatürk University

Assoc. Prof. Dr. Münevver BAYAR – Bilim University

Assoc. Prof. Dr. Muhammad Zia-ur-Rehman – University of Malaya & NDU

Assoc. Prof. Dr. Fatma KÖPRÜLÜ – Near East University

Assoc. Prof. Dr. Melda Medine SUNAY – Bursa Technical University

Assoc. Prof. Dr. Günel HUMMETOVA – Azerbaijan State Pedagogical University

Assist. Prof. Dr. Gönül GÜL – Çankırı Karatekin University

Dr. Zeynep YILMAZ ŞİMŞEK – Bingöl University

Dr. Ulduz Farhad Qahramanova – Azerbaijan State Pedagogical University

Assist. Prof. Dr. Ananda MAJUMDAR – USA

Dr. Keshab Chandra MANDAL – India

Lect. Ayla DEVECİ – Kilis 7 Aralık University

INTERNATIONAL SCIENTIFIC RESEARCH AND STRATEGY DEVELOPMENT ASSOCIATION
UBS
www.ubsdcr.org

5. ULUSLARARASI TOKAT BİLİMSEL ARAŞTIRMALAR KONGRESİ

7-9 Ağustos 2026, Tokat/TÜRKİYE

Sayı:2026/230

İlgili Makama:

Uluslararası Bilimsel Araştırma ve Strateji Geliştirme Derneği (UBSDER) tarafından organize edilen "**5. ULUSLARARASI TOKAT BİLİMSEL ARAŞTIRMALAR KONGRESİ**" Ağustos 7-9, 2026 tarihinde 6 farklı ülkeden (Türkiye'den 25, yurt dışından 45 bildiri sunuldu) olmak üzere toplam 70 adet sunum yapılarak, çevrimiçi olarak düzenlenmiştir.

Kongremiz, 16 Ocak 2020'de Akademik Teşvik Ödeneği Yönetmeliğine getirilen "Tebliğlerin sunulduğu yurt içinde veya yurt dışındaki etkinliğin uluslararası olarak nitelendirilebilmesi için Türkiye dışından en az beş farklı ülkeden sözlü tebliğ sunan konuşmacının katılım sağlaması ve tebliğlerin yarıdan fazlasının Türkiye dışından katılımcılar tarafından sunulması esastır." Değişikliğine uygun olarak düzenlenmiştir.

Bilgilerinize Arz Ederim

20.09.2026

Dr. Nadire KANTARCIOĞLU

UBSDER Başkanı

Kongremize katılan ülkeler:

Azerbaijan, Nigeria, Thailand, Türkiye, United Kingdom, Uzbekistan

INTERNATIONAL SCIENTIFIC RESEARCH AND STRATEGY DEVELOPMENT ASSOCIATION
www.ubsder.org

INSTITUTE CONTACT INFORMATION:
TURGUT ÖZAL MAH. AKKENT 6 TWINS TRADE CENTER ANKARA
TEL: 0 312 566 00 20 FAKS: 0 312 566 0021
Web : www.ubsder.org ubsder@ubsder.org

5th INTERNATIONAL TOKAT SCIENTIFIC RESEARCH CONGRESS

August 7-9, 2026 / Tokat, TÜRKİYE

www.ubsder.org

CONGRESS PROGRAM

Meeting Link:
https://us06web.zoom.us/j/4603636838?pwd=Gx9rIRf5cUTPsZZ4pilDYkYWaaEAc6.1&omn=84556738761

Meeting ID: 460 363 6838
Password: 121314

www.ubsder.org

PARTICIPANT COUNTRIES (6):
Azerbaijan, Nigeria, Thailand, Türkiye, United Kingdom, Uzbekistan

Türkiye: 25 **Other Countries: 45**

Opening Ceremony

Online

Date: **August 8, 2026**

Time: 14:00

Meeting Link:
https://us06web.zoom.us/j/4603636838?pwd=Gx9rIRf5cUTPsZZ4pilDYkYWaaEAc6.1&omn=84556738761

Meeting ID: 460 363 6838
Password: 121314

Speech:

Dr. Nadire KANTARCIOĞLU
UBSDER President, Türkiye

Speech:

Süreyya KAYA
Folk Poet of the Ministry of Culture and Tourism of the Republic of Türkiye

INTERNATIONAL SCIENTIFIC RESEARCH AND STRATEGY DEVELOPMENT ASSOCIATION
UBS
www.ubsder.org

ÖNEMLİ, DİKKATLE OKUYUNUZ LÜTFEN / IMPORTANT, PLEASE READ CAREFULLY

Önemli, Dikkatle Okuyunuz Lütfen

- ✓ Kongremizde Yazım Kurallarına uygun gönderilmiş ve bilim kurulundan geçen bildiriler için online (video konferans sistemi üzerinden) sunum imkanı sağlanmıştır.
- ✓ Online sunum yapabilmek için **https://zoom.us/join** sitesi üzerinden giriş yaparak "Meeting ID or Personal Link Name" yerine ID numarasını girerek oturuma katılabilirsiniz.
- ✓ Zoom uygulaması ücretsizdir ve hesap oluşturmaya gerek yoktur.
- ✓ Zoom uygulaması kaydolmadan kullanılabilir.
- ✓ Uygulama tablet, telefon ve PC'lerde çalışıyor.
- ✓ Her oturumdaki sunucular, sunum saatinden 15 dk öncesinde oturuma bağlanmış olmaları gerekmektedir.
- ✓ Tüm kongre katılımcıları canlı bağlanarak tüm oturumları dinleyebilir.
- ✓ Moderatör – oturumdaki sunum ve bilimsel tartışma (soru-cevap) kısmından sorumludur.

Dikkat Edilmesi Gerekenler- TEKNİK BİLGİLER

- ✓ Bilgisayarınızda mikrofon olduğuna ve çalıştığına emin olun.
- ✓ Zoom'da ekran paylaşma özelliğine kullanabilmelisiniz.
- ✓ Kabul edilen bildiri sahiplerinin mail adreslerine Zoom uygulamasında oluşturduğumuz oturuma ait ID numarası gönderilecektir.
- ✓ **Katılım belgeleri kongre sonunda tarafınıza pdf olarak gönderilecektir**
- ✓ Kongre programında yer ve saat değişikliği gibi talepler dikkate alınmayacaktır

IMPORTANT, PLEASE READ CAREFULLY

- ✓ To be able to attend a meeting online, login via **https://zoom.us/join** site, enter ID "Meeting ID or Personal Link Name" and solidify the session.
- ✓ The Zoom application is free and no need to create an account.
- ✓ The Zoom application can be used without registration.
- ✓ The application works on tablets, phones and PCs.
- ✓ The participant must be connected to the session 15 minutes before the presentation time.
- ✓ All congress participants can connect live and listen to all sessions.
- ✓ Moderator is responsible for the presentation and scientific discussion (question-answer) section of the session.

Points to Take into Consideration - TECHNICAL INFORMATION

- ✓ Make sure your computer has a microphone and is working.
- ✓ You should be able to use screen sharing feature in Zoom.
- ✓ **Attendance certificates will be sent to you as pdf at the end of the congress.**
- ✓ Requests such as change of place and time will not be taken into consideration in the congress program.

Before you login to Zoom please indicate your name_surname and HALL number,

exp. Hall-1, Hakan ÖZTÜRK

Meeting ID: 460 363 6838

Passcode: 121314

IMPORTANT, PLEASE READ CAREFULLY

- *To be able to make a meeting online, login via https://zoom.us/join site, enter ID instead of "Meeting ID or Personal Link Name" and solidify the session.*
- *The Zoom application is free and no need to create an account.*
- *The Zoom application can be used without registration.*
- *The application works on tablets, phones and PCs.*
- *Speakers must be connected to the session 10 minutes before the presentation time.*
- *Online presentations; Presentation time is 10 minutes. There is 5 minutes of question and answer time.*
- *All congress participants can connect live and listen to all sessions.*
- *During the session, your camera should be turned on at least %70 of session period*
- *Moderator is responsible for the presentation and scientific discussion (question-answer) section of the session.*

TECHNICAL INFORMATION

- *Make sure your computer has a microphone and is working.*
- *You should be able to use screen sharing feature in Zoom.*
- *Attendance certificates will be sent to you as pdf at the end of the congress.*
- *Moderator is responsible for the presentation and scientific discussion (question-answer) section of the session.*

INTERNATIONAL SCIENTIFIC RESEARCH AND STRATEGY DEVELOPMENT ASSOCIATION
UBS
www.ubsder.org

08.08.2026	SESSION-1 / HALL-1
TOKAT	TÜRKİYE TIME: 14:30

CHAIR: ***Assist. Prof. Dr. Fikret BUDAK***

Author(s)	Affiliation	Title
Rukiye YATCI Assist. Prof. Dr. Fikret BUDAK	Düzce University	THE EFFECTS OF DIFFERENT FERTILISER TYPES ON THE YIELD AND QUALITY OF HUNGARIAN VETCH (VICIA PANNONICA CRANTZ) UNDER ANTALYA CONDITIONS
Edanur SÖNMEZ Assist. Prof. Dr. Fikret BUDAK	Düzce University	DETERMINATION OF AGRICULTURAL CHARACTERISTICS OF SOME FORAGE PEA [Pisum sativum ssp. arvense (L.) Poir.] VARIETIES UNDER ZONGULDAK ECOLOGICAL CONDITIONS
Hilal Özge DEMİR Assist. Prof. Dr. Fikret BUDAK	Düzce University	INVESTIGATION OF THE YIELD AND QUALITY CHARACTERISTICS OF SOME ITALIAN RYEGRASS (LOLIUM MULTIFLORUM LAM.) CULTIVARS UNDER BALIKESİR ECOLOGICAL CONDITIONS

www.ubsder.org

08.08.2026	SESSION-1 / HALL-2
TOKAT	TÜRKİYE TIME: 14:30

CHAIR: ***Lecturer Dr. İsmail KILIÇ***

Author(s)	Affiliation	Title
Gökay ÜSTYOL Prof. Dr. Mehmet Sait İZGİ Assoc. Prof. Dr. Erhan ONAT	Bitlis Eren University Siirt University	STUDY OF THE CATALYTIC HYDROLYSIS KINETICS OF SODIUM BOROHYDRIDE IN THE PRESENCE OF A SUPPORTED BIMETALLIC CATALYST
Assoc. Prof. Dr. Erhan ONAT Gökay ÜSTYOL Prof. Dr. Mehmet Sait İZGİ	Bitlis Eren University Siirt University	STUDY OF THE HYDROLYSIS OF SODIUM BOROHYDRIDE IN THE PRESENCE OF A GRAPHITIC CARBON NITRIDE-SUPPORTED BIMETALLIC CATALYST
Lecturer Dr. İsmail KILIÇ	Yozgat Bozok University	NUMERICAL MODELLING OF GLUED LAMINATED TIMBER (GLULAM) BEAMS: CURRENT APPROACHES, EXPERIMENTAL VALIDATION, RESEARCH GAPS, AND FUTURE TRENDS
Assoc. Prof. Dr. Omer CIHAN Assoc. Prof. Dr. Mustafa Guven GOK	Sakarya University Gaziantep University	THE ROLE OF MATERIAL SELECTION IN LIGHTWEIGHTING ELECTRIC VEHICLES

www.ubsder.org

5th INTERNATIONAL TOKAT SCIENTIFIC RESEARCH CONGRESS August 7-9, 2026 / Tokat, TÜRKİYE ONLINE	
08.08.2026	SESSION-1 / HALL-3
TOKAT	TÜRKİYE TIME: 14:30

CHAIR: *Prof. Dr. Metin AKSOY*

Author(s)	Affiliation	Title
Dr. Erdal Taha AYDOĞDU Dr. Dursun YILMAZ	Tokat Gaziosmanpaşa University	THE CONSTRUCTION OF A LEADERSHIP PROFILE THROUGH COMMUNICATION IN THE CONTEXT OF THE NATO ANKARA SUMMIT: THE CASE OF PRESIDENT RECEP TAYYIP ERDOĞAN
Dr. Dursun YILMAZ Dr. Erdal Taha AYDOĞDU	Tokat Gaziosmanpaşa University	SYMBOLIC COMMUNICATION IN SPORTS DIPLOMACY AND THE INTERNATIONAL VISIBILITY OF THE PALESTINE ISSUE: THE CASE OF LAMINE YAMAL
Lecturer Yücel NAKTİYOK Lecturer Fatih UZUNBOY	Atatürk University	THE IMPACT OF PRODUCT- AND COUNTRY-SPECIFIC RULES TO BE COMPLIED WITH IN EXPORTS ON TURKEY'S FOREIGN TRADE
Lecturer Yücel NAKTİYOK	Atatürk University	THE GENERAL COURSE OF FOREIGN TRADE IN TURKEY BEFORE AND AFTER THE PANDEMIC
Prof. Dr. Metin AKSOY Lect. Muhlise CANLI	Gümüşhane University	THE THEORETICAL VALUE OF SMALL SETTLEMENTS IN MIGRATION RESEARCH: A CRITICAL ASSESSMENT OF METROPOLIS-CENTRED APPROACHES
Prof. Dr. Metin AKSOY Lect. Muhlise CANLI	Gümüşhane University	THE INVISIBLE GOVERNANCE OF MIGRATION IN SMALL TOWNS: LOCAL INSTITUTIONS, EVERYDAY PRACTICES, AND INSTITUTIONAL LIMITS
Muhammad FAISAL	Joint Director (AICU) at Ministry of IT & Telecommunication, Pakistan	MULTI-AGENT ARTIFICIAL INTELLIGENCE SYSTEMS FOR COLLABORATIVE BUSINESS DECISION INTELLIGENCE

www.ubsder.org

08.08.2026	SESSION-1 / HALL-4
TOKAT	TÜRKİYE TIME: 14:30

CHAIR: ***Ulduz Fərhad Qəhrəmanova***
CHAIR OF VICE: ***Emil Raul oğlu AĞAYEV***

Author(s)	Affiliation	Title
GÜNEL HÜMMƏTOVA (Pəniyeva) Saleh	Ph.D. in Philology, Senior Lecturer; Azerbaijan State Pedagogical University, Department of Literature Teaching Technology	MÜHARİBƏ DÖVRÜ UŞAQLARININ MƏNƏVİ DÜNYASI VƏ İNSANLIQ DƏRSLƏRİ MAQSUD İBRAHİMBƏYOVUN "PÜSTƏ AĞACI" ƏSƏRİNDƏ SİMVOLİZM VƏ SOSİAL REALLIQ
GÜNEL HÜMMƏTOVA (Pəniyeva) Saleh	Ph.D. in Philology, Senior Lecturer; Azerbaijan State Pedagogical University, Department of Literature Teaching Technology	EASTERN RESONANCE AND THE SOCIO-CULTURAL IMPACT SPECTRUM OF ALISHER NAVOI'S HERITAGE IN THE HISTORY OF AZERBAIJANI LITERARY THOUGHT
Ulduz Fərhad Qəhrəmanova	Azerbaijan State Pedagogical University, Associate Professor at the Department of "Literature"	NIZAMI GANJAVI'S POEM "THE TREASURY OF SECRETS" III CONVERSATION: THE WORLD'S TROUBLES AND THE SUFFERING OF MANKIND
Emil Raul oğlu AĞAYEV	Azerbaijan State Pedagogical University	THE ROLE OF FINE ARTS EDUCATION IN A COLLABORATIVE LEARNING ENVIRONMENT
Asgarova Ganira Ali gizi	Azerbaijan State Pedagogical College under Azerbaijan State Pedagogical University, Azerbaijan	ABOUT SOME VERBS USED IN AZERBAIJAN DIALECTS AND ACCENTS

www.ubsder.org

08.08.2026	SESSION-1 / HALL-5
TOKAT	TÜRKİYE TIME: 14:30

CHAIR *Asst. Prof. Dr. Hakan AVAN*

Author(s)	Affiliation	Title
Asst. Prof. Dr. Hakan AVAN	Kahramanmaraş Sütçü İmam University	THE LONELINESS EPIDEMIC IN CHILDREN AND ADOLESCENTS: HEALTH OUTCOMES AND NURSING APPROACHES
Asst. Prof. Dr. Hakan AVAN	Kahramanmaraş Sütçü İmam University	ARTIFICIAL INTELLIGENCE-ASSISTED NURSING CARE FOR CHILDREN AND ADOLESCENTS: CURRENT PRACTICES, OPPORTUNITIES, AND ETHICAL ISSUES
Büşra ERDEN	Hacı Bektaş Veli University,	LAYERS OF MEANING OF THE CONCEPT OF SIRAT IN THE QUR'AN
Lecturer Havana KAYNAK	Gümüşhane University	TEXT ADAPTATION FOR THE B1 LEVEL IN TEACHING TURKISH TO FOREIGNERS: A CASE STUDY OF SAIT FAIK ABASIYANIK'S STORY "İPEKLI MENDIL"
Lecturer Havana KAYNAK Yağmur ÜLKMEN	Independent Researcher Gümüşhane University	LITERARY TEXT ADAPTATION IN TEACHING TURKISH TO FOREIGNERS: ADAPTING AHMET HAMDI TANPINAR'S SHORT STORY "FAL" TO THE B1 LEVEL
Yasemin ALBAYRAK Assoc. Prof. Dr. Sıddık BAKIR	Atatürk University	AN EVALUATION OF CURRENT GENERATIVE AI TOOLS USED IN LANGUAGE TEACHING FROM THE PERSPECTIVE OF TURKISH LANGUAGE EDUCATION: OPPORTUNITIES AND RISKS
Yasemin ALBAYRAK Assoc. Prof. Dr. Sıddık BAKIR	Atatürk University	AI-SUPPORTED APPROACHES TO INCREASING STUDENT PARTICIPATION IN TEACHING TURKISH TO FOREIGNERS

www.ubsder.org

5th INTERNATIONAL TOKAT SCIENTIFIC RESEARCH CONGRESS
August 7-9, 2026 / Tokat, TÜRKİYE
ONLINE

08.08.2026	SESSION-1 / HALL-6
TOKAT	TÜRKİYE TIME: 14:30

CHAIR ***Asst. Prof. Dr. Fesih KESKİN***

Author(s)	Affiliation	Title
Moses Adeolu AGOI Olayemi Grace ABIMBOLA Yewande Adeyosola ODUNAYO Emmanuel Taiwo AGOI	Lagos State University of Education, Lagos, Nigeria Loughborough University, United Kingdom	NEXT-GENERATION AUTONOMOUS DESIGN SYSTEMS: LEVERAGING GENERATIVE AI, COMPUTER VISION, AND DIGITAL TWINS FOR SMART ARCHITECTURAL PRACTICE
Moses Adeolu AGOI Bamidele Olumuyiwa ALABA Gbenga Oyewole OGUNSANWO Hendri Hermawan ADINUGRAHA	Lagos State University of Education, Lagos Nigeria. Tai Solarin University of Education, Ogun Nigeria. Universitas Islam Negeri K.H Abdurrahman Wahid Pekalongan.	ADVANCING VETERINARY DISEASE MANAGEMENT THROUGH MACHINE LEARNING-DRIVEN ANTIMICROBIAL RESISTANCE PREDICTION
Prof. Dr. Murat Subaşı Assoc. Prof. Dr. Serpil Yorgancı	Atatürk University	ARTIFICIAL INTELLIGENCE INTEGRATION IN MATHEMATICS EDUCATION: AN INTEGRATIVE FRAMEWORK GROUNDED IN SELF-DETERMINATION THEORY
Prof. Dr. Murat Subaşı Assoc. Prof. Dr. Serpil Yorgancı	Atatürk University	MANAGING COGNITIVE LOAD IN AI-SUPPORTED MATHEMATICS LEARNING
M.Sc., Esra YILMAZ ÖNDER	Tokat Gaziosmanpaşa University	MAPPING RECENT TRENDS IN ARTIFICIAL INTELLIGENCE RESEARCH FOR SUSTAINABLE AGRICULTURE: A BIBLIOMETRIC AND SYSTEMATIC REVIEW
Assistant Prof. Dr. Serap SARP	Abdullah Gül University	TRANSFORMATION FACTORS FOR RESPONSIBLE AI-DRIVEN CONSUMER ENGAGEMENT: A QUALITATIVE EXPLORATION
Ramazan AKYOL	Adıyaman University	A STRUCTURAL COMPARISON OF THE LEARNING OBJECTIVES IN THE 2018 SOCIAL STUDIES CURRICULUM AND THE LEARNING OUTCOMES IN THE 2024 CURRICULUM

www.ubsder.org

09.08.2026	SESSION-2 / HALL-1
TOKAT	TÜRKİYE TIME: 13:00

CHAIR: ***Mamadaliyev Boburjon Odiljon o'g'li***

Author(s)	Affiliation	Title
Anorboev D.G Irismetov M.E. Razhabov K.N	Republican Scientific and Practical Center for the Study of Traumatology and Otropedia, Tashkent	OUR EXPERIENCE WITH REINSERTATION OF THE ANTERIOR CRUCIATE LIGAMENT OF THE KNEE JOINT
Mamadaliyev Boburjon Odiljon o'g'li; Normatova Shakhnoza Anvarovna	4th- student of Pediatrics, Central Asian Medical University; Vice-Rector for Research and Innovation, PhD, Associate Professor Central Asian Medical University	NEURODEVELOPMENTAL AND NUTRITIONAL DETERMINANTS OF ATTENTION AND CHILD DEVELOPMENT IN THE FERGANA REGION: AN INTEGRATED CLINICAL AND EPIDEMIOLOGICAL STUDY
Nishonova Dilafruz Tajimatovna	Central Asian Medical University medical university Assistant of the Department of "Anatomy and mikroanatomy"	IMPROVING THE METHODOLOGY OF TEACHING ANATOMY TO FUTURE DOCTORS IN A DIGITAL LEARNING ENVIRONMENT.
Kholikova Oyistakhon Yuldashevna	Fergana Medical Institute of Public Health, Doctor of Philosophy, Head of the Department of Folk Medicine and Pharmacology	TELEMEDICINE AND ARTIFICIAL INTELLIGENCE INTEGRATION: NEW OPPORTUNITIES IN CLINICAL PHARMACOLOGY
Ph.D. Musaev Farxod Taxirovich	Central Aisan Medical University, Head of the Department of Phthisiology, Pulmonology and Metabolic Diseases, Fergana, Uzbekistan	ANALYSIS OF THE MAIN CLINICAL AND FUNCTIONAL PARAMETERS AFTER TREATMENT IN PATIENTS WITH LOW AND HIGH RISK OF CHRONIC OBSTRUCTIVE PULMONARY DISEASE
Dilrabo Salizhanovna Tadjibaeva	Assistant, Central Asian Medical University Uzbekistan, Fergana	THE EFFECT OF SPINAL PROTRUSION ON THE FUNCTIONING OF INTERNAL ORGANS
Nilufar Akhmedova Ozoda Atabaeva	Doctor of Medical Sciences, Associate Professor Department of Hospital Pediatrics and Traditional Medicine Tashkent State Medical University; Department of Anatomy, Pathological Anatomy and Histology Tashkent State Medical University	THE INFLUENCE OF CONNECTIVE TISSUE DYSPLASIA ON THE CLINICAL AND IMMUNOLOGICAL FEATURES OF THE COURSE OF COMMUNITY-ACQUIRED PNEUMONIA IN CHILDREN

09.08.2026	SESSION-2 / HALL-2
TOKAT	TÜRKİYE TIME: 13:00

CHAIR: ***Ruzmatova Hiloloy Kabilovna***

Author(s)	Affiliation	Title
Ruzmatova Hiloloy Kabilovna PhD, Associate Professor Sh.A. Normatova	Central Asian Medical University, Fergana, Uzbekistan	PREVENTIVE GERIATRICS: MODERN APPROACHES AND EVALUATION OF EFFECTIVENESS
Ismailov Akhroridin Zakhridinovich Kakhkharova Safia Sarvarjon kizi	Central Asian Medical University; Central Asian Medical University	CHANGE IN ELECTROLYTE SUBSTANCES OF SYNOVIAL FLUID IN PATIENTS WITH OSTEOARTHROSIS OF THE KNEE AT DIFFERENT STAGES OF THE DISEASE
S.Z.Sayfuddinov; Proff. DSc. A.L.Kasimov	Central Asian Medical University, Assistent of the Department of Surgical Sciences, Fergana, Uzbekistan; Andijan State Medical Institute, Professor of the Department of General Surgery and Transplantology, Andijan, Uzbekistan	CHOLEDOCHOLITHIASIS COMPLICATED BY OBSTRUCTIVE JAUNDICE: MODERN STRATEGIES FOR DIAGNOSIS AND TREATMENT
Khabibullayev Ne‘matillo Muminjonova Shohzoda	Student of Central Asian Medical University; Student of Central Asian Medical University	INFECTIONS IN SURGICAL PRACTICE: DEVELOPMENT MECHANISMS, RISK FACTORS, DIAGNOSTIC METHODS, AND MODERN PREVENTIVE APPROACHES
Muminjonova Shohzoda Khabibullayev Ne‘matillo	Student of Central Asian Medical University; Student of Central Asian Medical University	BACTERIAL AND VIRAL INFECTIONS IN SURGICAL PRACTICE: DIAGNOSTIC, THERAPEUTIC, AND PREVENTIVE APPROACHES
Khabibullayev Ne‘matillo Muminjonova Shohzoda	Student of Central Asian Medical University; Student of Central Asian Medical University	POSTOPERATIVE SURGICAL INFECTIONS: DEVELOPMENT MECHANISMS, DIAGNOSTIC METHODS, AND PREVENTIVE STRATEGIES

www.ubsder.org

09.08.2026	SESSION-2 / HALL-3
TOKAT	TÜRKİYE TIME: 13:00

CHAIR: ***Foziljon Rasulov***

Author(s)	Affiliation	Title
Mutabarova Mohinora Khabibullayev Ne'matillo	Students Central Asian Medical University Uzbekistan, Fergana City; Students Central Asian Medical University Uzbekistan, Fergana City	OPPORTUNITIES FOR EARLY DETECTION AND SURGERY REDUCTION THROUGH PROPHYLACTIC ENDOSCOPIC INTERVENTIONS
Foziljon Rasulov	Fergana Medical Institute of Public Health Department of Microbiology, Virology, and Immunology The Republic of Uzbekistan. Fergana city.	STUDY OF THE EFFECT OF HERBAL PREPARATIONS ON IMMUNOGENETICS AND HEMATOPOIESIS IN THE EXPERIMENT
Ph.D.GM Mamatkhanova	Department of Fundamentals of Preventive Medicine, Public Health, Physical Education, and Sports Fergana Medical Institute of Public Health. Uzbekistan, Fergana.	BENEFITS OF IMPLEMENTING AN ELECTRONIC HEALTHCARE SYSTEM
Siddikov Bobirjon Tursinali ugli	Fergana Regional Health Department	ACUTE CARDIOVASCULAR DISEASES IN THE PRIMARY HEALTH CARE SYSTEM
Yulduz Djabbarova Kasimovna Maqsudova Barno Jaxongirovna	Doctor of Medical Sciences, Academician of the Academy of Medical and Technical Sciences of the Russian Federation, , Professor of the Department of Obstetrics and Gynecology, Reproductology of the Tashkent State Medical University, Professor of the Department of Obstetrics and Gynecology,Neonatology of the Central Asian Medical Master's degree student of the Department of Obstetrics and Gynecology, International Medical University, Central Asian Medical University, Student	ORGAN-PRESERVING STRATEGY AND PREVENTION OF HYSTERECTOMY IN PLACENTA PREVIA IN WOMEN WITH A UTERINE SCAR
Senior assistant, Fatima Adilbaevna Khalmuratova Doctor of Medical Sciences, Professor Ziyaviddin Zainutdinovich Hakimov Doctor of Medical Sciences, Professor Alisher Khudayberdievich Rakhmanov	Karakalpakstan Medical Institute, Nukus, Karakalpakstan, Uzbekistan Department of Pharmacology, Tashkent State Medical University , Tashkent, Uzbekistan Institute of Medical, Scientific and Innovative Research at the Tashkent State Medical University, Tashkent, Uzbekistan	ASSESSMENT OF THE ANTIOXIDANT ACTIVITY OF THE DRUG "RUTAN" IN ACUTE TOXIC HEPATITIS AT THE PREPUBERTAL AGE
Atabekov N.S., Doctor of Medical Sciences Shadmanova N.A., Doctor of Medical Sciences Iskandarova G.T., Doctor of Medical Sciences, Associate Professor Asatova N.B., Candidate of Medical Sciences, Associate Professor Sapaeva F.R., Doctor of Biological Sciences, Associate Professor	Committee for Sanitary-Epidemiological Welfare and Public Health of the Republic of Uzbekistan Center for Professional Development of Medical Workers under the Ministry of Health of the Republic of Uzbekistan	CONTEMPORARY APPROACHES TO THE LABORATORY ETIOLOGICAL DIAGNOSIS OF FOODBORNE POISONING AND ISSUES OF THEIR EPIDEMIOLOGICAL SURVEILLANCE IN UZBEKISTAN

5th INTERNATIONAL TOKAT SCIENTIFIC RESEARCH CONGRESS August 7-9, 2026 / Tokat, TÜRKİYE ONLINE	
09.08.2026	SESSION-2 / HALL-4
TOKAT	TÜRKİYE TIME: 13:00

CHAIR: ***Sulaymanov Dilshod Abdurakhimovich***

Author(s)	Affiliation	Title
Sulaymanov Dilshod Abdurakhimovich	PhD, MD Head of department of oncology and radiology, Central Asian Medical University, Fergana, Uzbekistan.	ONCOLOGIC OUTCOMES OF HYPERTHERMIC INTRAPERITONEAL CHEMOTHERAPY IN THE TREATMENT OF ASCITES-TYPE OVARIAN CANCER (ATOC): RESIDUAL DISEASE AND METASTASIS
Daripov Shahobiddin Israiljonovich	Central Asian Medical University, Senior Lecturer, Department of Social Sciences and Physical Culture. Uzbekistan, Fergana	HEALTHCARE REFORMS AS A STRATEGIC FACTOR IN POVERTY REDUCTION
Shakhnoza Kholboboyeva Asadullayeva	Central Asian Medical University Fergana, Uzbekistan	AGE- SPECIFIC PREVENTIVE STRATEGIES FOR ATHEROSCLEROSIS AND THEIR EFFECTIVENESS
Shodmonov Usmonbek Bakhodir ugli	Central Asian Medical University, Fergana, Uzbekistan	MICROORGANISMS CAUSING HIGHLY DANGEROUS INFECTIOUS DISEASES IN SOIL AND THEIR ECOLOGY
Abdukarimova Marhabo Ruzamatovna Ergashboev Ozodbek Sherzod ugli	Students Department of dentistry International Medical University, Central Asian Medical University Uzbekistan, Fergana City; Students Department of dentistry International Medical University, Central Asian Medical University Uzbekistan, Fergana City	OPTIMIZATION OF MAXILLOFACIAL PLASTIC SURGERY METHODS FOR DENTITIAL DEFECTS

www.ubsder.org

09.08.2026	SESSION-2 / HALL-5
TOKAT	TÜRKİYE TIME: 13:00

CHAIR: ***Madrakhimova Zukhra Mamasodikovna***

Author(s)	Affiliation	Title
Madrakhimova Zukhra Mamasodikovna	Lecturer, Central Asian Medical University, Fergana Region, Uzbekistan	STRUCTURE OF TEMPORARY DISABILITY CONDITIONS AMONG MEDICAL WORKERS AND THEIR RELATIONSHIP WITH WORK EXPERIENCE (THE CASE OF MARGILON CITY, UZBEKISTAN)
Sarimsakov Mahamadjalol Isakjonovich	Senior Lecturer, Fergana Medical Institute of Public Health Uzbekistan, Fergana	PHARMACOLOGICAL PROPERTIES OF ESSENTIAL OIL FROM MEDICINAL CHAMOMILE (MATRICARIA CHAMOMILLA L.) AND ITS ANTIMICROBIAL ACTIVITY
Assistant Tohir Mukaramov	Central Asian Medical University, Department of Hygienic and Medical Process Modeling, Uzbekistan, Fergana	DEVELOPING MEDIA LITERACY IN FUTURE TEACHERS: CURRENT TRENDS AND A COMPARATIVE ANALYSIS OF LOCAL AND INTERNATIONAL PRACTICES
Akhunjonova Khakimakhon Abdumannabovna Mirzayev Muhammadjon Ubaydullo oglu	Central Asian Medical University, Fergana; Central Asian Medical University, Fergana	PREVENTION OF CORRUPTION IN THE HEALTHCARE SYSTEM
Qalandarov Dilmurod Madaminovich Kenjayev Olimjon Obidjonovich Ravshanov Abdullajon Shukhrat ogli Mamasaliev Nematjon Solievich Usmonov Burhonjon Umarovich Salokhidinov Adiljon	Andijan State Medical Institute Central Asian Medical University Andijan State Medical Institute	PREVALENCE OF ARTERIAL HYPERTENSION AND THE DYNAMICS OF ITS 31-YEAR CHANGES IN THE INDIGENOUS AND MIGRANT RURAL POPULATION OF THE ANDIJAN REGION
DSC. Prof, Xakimhon Xanbabayev Assoc. Prof., Akhrorjon ADBUAMANONOV	Department of “Digital technology and artificial intelligence”, Kokand State University, Fergana, Uzbekistan; Department of “Modeling of Medical Hygiene Processes”, Central Asian Medical University, Fergana, Uzbekistan	FROM TRANSMISSION TO PERSONALISATION: A GAMIFIED INTELLIGENT TUTORING FRAMEWORK FOR ENGINEERING COMPETENCY DEVELOPMENT

www.ubsder.org

09.08.2026	SESSION-2 / HALL-6
TOKAT	TÜRKİYE TIME: 13:00

CHAIR: ***Sayidmirzayeva Nazokat Gayratjonovna***

Author(s)	Affiliation	Title
Sayidmirzayeva Nazokat Gayratjonovna Shadmanova Hargiza Abidovna	Assistant of Central Asian Medical University, Fergana, Uzbekistan; Center for Retraining and Advanced Training of Personnel in the Field of Sanitary and	ANTIBIOTIC RESISTANCE AND UNUSUAL PHENOTYPIC CHARACTERISTICS OF LOCAL STRAINS OF PATHOGENS CAUSING ACUTE INTESTINAL INFECTIONS (REPUBLIC OF UZBEKISTAN)
M. Mamajonov	Central Asian Medical University, Department of Hygienic and Medical Process Modeling, Uzbekistan, Fergana	THE PREVALENCE OF FLATFOOT, ITS CAUSES, AND THE POSSIBILITIES OF EARLY DETECTION OF ITS COMPLICATIONS THROUGH MODERN DIAGNOSTIC METHODS
M.M. Mamajonov M.B. To'xtasheva	Associate Professor, Central Asian Medical University, Department of Hygienic and Medical Process Modeling, Uzbekistan, Fergana; Student, General Medicine Program, Central Asian Medical University	NEUROBIOLOGICAL MECHANISMS OF COGNITIVE IMPAIRMENT IN ALZHEIMER'S DISEASE
G.N. Rayimov N.A. Salokhitdinov	Central Asian Medical University, Department of Surgery and surgical subjects, Fergana, Uzbekistan; Central Asian Medical University, Department of Surgery and surgical subjects, Fergana, Uzbekistan	RESULTS OF SURGICAL TREATMENT OF POST-TRAUMATIC PANCREATITIS IN COMBINED ABDOMINAL TRAUMA
Sh.Sh. Dekhkonov G.N. Rayimov O. Juraboev	Central Asian Medical University, Department of Surgery and surgical subjects, Fergana, Uzbekistan; Central Asian Medical University, Department of Surgery and surgical subjects, Fergana, Uzbekistan; Central Asian Medical University, Department of Surgery and surgical subjects, Fergana, Uzbekistan	COMPARATIVE EVALUATION OF OPEN AND ENDOSCOPIC INTERVENTIONS FOR OBSTRUCTION OF THE TERMINAL COMMON BILE DUCT IN ELDERLY AND SENILE PATIENTS
Dr. Nattapong Wattanakul Kanya Srisuk	Chiang Mai University, Thailand	ONE HEALTH APPROACH IN THE SURVEILLANCE OF EMERGING ZOONOTIC DISEASES IN SOUTHEAST ASIA

www.ubsder.org

CONTENT

CONFERENCE ID	I
ORGANIZING COMMITTEE	II
SCIENTIFIC BOARD MEMBERS	III
PROGRAM	V
CONTENT	VI

	AUTHOR(S)	TITLE	PAGE NUMBER
1)	Rukiye YATCI Assist. Prof. Dr. Fikret BUDAK	THE EFFECTS OF DIFFERENT FERTILISER TYPES ON THE YIELD AND QUALITY OF HUNGARIAN VETCH (VICIA PANNONICA CRANTZ) UNDER ANTALYA CONDITIONS	1-26
2)	Edanur SÖNMEZ Assist. Prof. Dr. Fikret BUDAK	DETERMINATION OF AGRICULTURAL CHARACTERISTICS OF SOME FORAGE PEA [Pisum sativum ssp. arvense (L.) Poir.] VARIETIES UNDER ZONGULDAK ECOLOGICAL CONDITIONS	27-43
3)	Hilal Özge DEMİR Assist. Prof. Dr. Fikret BUDAK	INVESTIGATION OF THE YIELD AND QUALITY CHARACTERISTICS OF SOME ITALIAN RYEGRASS (LOLIUM MULTIFLORUM LAM.) CULTIVARS UNDER BALIKESİR ECOLOGICAL CONDITIONS	44
4)	Assoc. Prof. Dr. Erhan ONAT Gökay ÜSTYOL Prof. Dr. Mehmet Sait İZGİ	STUDY OF THE HYDROLYSIS OF SODIUM BOROHYDRIDE IN THE PRESENCE OF A GRAPHITIC CARBON NITRIDE-SUPPORTED BIMETALLIC CATALYST	45-53
5)	Gökay ÜSTYOL Prof. Dr. Mehmet Sait İZGİ Assoc. Prof. Dr. Erhan ONAT	STUDY OF THE CATALYTIC HYDROLYSIS KINETICS OF SODIUM BOROHYDRIDE IN THE PRESENCE OF A SUPPORTED BIMETALLIC CATALYST	54-60
6)	Lecturer Dr. İsmail KILIÇ	NUMERICAL MODELLING OF GLUED LAMINATED TIMBER (GLULAM) BEAMS: CURRENT APPROACHES, EXPERIMENTAL VALIDATION, RESEARCH GAPS, AND FUTURE TRENDS	61-78

ISBN: 978-625-90699-1-3

7)	Assoc. Prof. Dr. Omer CIHAN Assoc. Prof. Dr. Mustafa Guven GOK	THE ROLE OF MATERIAL SELECTION IN LIGHTWEIGHTING ELECTRIC VEHICLES	79-93
8)	Dr. Dursun YILMAZ Dr. Erdal Taha AYDOĞDU	SYMBOLIC COMMUNICATION IN SPORTS DIPLOMACY AND THE INTERNATIONAL VISIBILITY OF THE PALESTINE ISSUE: THE CASE OF LAMINE YAMAL	94-104
9)	Dr. Erdal Taha AYDOĞDU Dr. Dursun YILMAZ	THE CONSTRUCTION OF A LEADERSHIP PROFILE THROUGH COMMUNICATION IN THE CONTEXT OF THE NATO ANKARA SUMMIT: THE CASE OF PRESIDENT RECEP TAYYIP ERDOĞAN	105-106
10)	Lecturer Yücel NAKTİYOK Lecturer Fatih UZUNBOY	THE IMPACT OF PRODUCT- AND COUNTRY-SPECIFIC RULES TO BE COMPLIED WITH IN EXPORTS ON TURKEY'S FOREIGN TRADE	107-117
11)	Lecturer Yücel NAKTİYOK	THE GENERAL COURSE OF FOREIGN TRADE IN TURKEY BEFORE AND AFTER THE PANDEMIC	118-128
12)	Prof. Dr. Metin AKSOY Lect. Muhlise CANLI	THE THEORETICAL VALUE OF SMALL SETTLEMENTS IN MIGRATION RESEARCH: A CRITICAL ASSESSMENT OF METROPOLIS-CENTRED APPROACHES	129-136
13)	GÜNEL HÜMMƏTOVA (Pəniyeva) Saleh	MÜHARİBƏ DÖVRÜ UŞAQLARININ MƏNƏVİ DÜNYASI VƏ İNSANLIQ DƏRSLƏRİ MAQSUD İBRAHİMBƏYOVUN "PÜSTƏ AĞACI" ƏSƏRİNDƏ SİMVOLİZM VƏ SOSİAL REALLIQ	137-139
14)	Prof. Dr. Metin AKSOY Lect. Muhlise CANLI	THE INVISIBLE GOVERNANCE OF MIGRATION IN SMALL TOWNS: LOCAL INSTITUTIONS, EVERYDAY PRACTICES, AND INSTITUTIONAL LIMITS	140-148
15)	GÜNEL HÜMMƏTOVA (Pəniyeva) Saleh	EASTERN RESONANCE AND THE SOCIO-CULTURAL IMPACT SPECTRUM OF ALISHER NAVOI'S HERITAGE IN THE HISTORY OF AZERBAIJANI LITERARY THOUGHT	149-153
16)	Ulduz Fərhad Qəhrəmanova	NIZAMI GANJAVI'S POEM "THE TREASURY OF SECRETS" III CONVERSATION: THE WORLD'S TROUBLES AND THE SUFFERING OF MANKIND	154-155

ISBN: 978-625-90699-1-3

17)	Emil Raul oğlu AĞAYEV	THE ROLE OF FINE ARTS EDUCATION IN A COLLABORATIVE LEARNING ENVIRONMENT	156-158
18)	Asgarova Ganira Ali gizi	ABOUT SOME VERBS USED IN AZERBAIJAN DIALECTS AND ACCENTS	159-166
19)	Asst. Prof. Dr. Hakan AVAN	THE LONELINESS EPIDEMIC IN CHILDREN AND ADOLESCENTS: HEALTH OUTCOMES AND NURSING APPROACHES	167-175
20)	Asst. Prof. Dr. Hakan AVAN	ARTIFICIAL INTELLIGENCE-ASSISTED NURSING CARE FOR CHILDREN AND ADOLESCENTS: CURRENT PRACTICES, OPPORTUNITIES, AND ETHICAL ISSUES	176-188
21)	Büşra ERDEN	LAYERS OF MEANING OF THE CONCEPT OF SIRAT IN THE QUR'AN	189-195
22)	Lecturer Havana KAYNAK Yağmur ÜLKMEN	LITERARY TEXT ADAPTATION IN TEACHING TURKISH TO FOREIGNERS: ADAPTING AHMET HAMDI TANPINAR'S SHORT STORY "FAL" TO THE B1 LEVEL	196-218
23)	Lecturer Havana KAYNAK	TEXT ADAPTATION FOR THE B1 LEVEL IN TEACHING TURKISH TO FOREIGNERS: A CASE STUDY OF SAIT FAIK ABASIYANIK'S STORY "İPEKLI MENDIL"	219-235
24)	Yasemin ALBAYRAK Assoc. Prof. Dr. Sıddık BAKIR	AN EVALUATION OF CURRENT GENERATIVE AI TOOLS USED IN LANGUAGE TEACHING FROM THE PERSPECTIVE OF TURKISH LANGUAGE EDUCATION: OPPORTUNITIES AND RISKS	236
25)	Yasemin ALBAYRAK Assoc. Prof. Dr. Sıddık BAKIR	AI-SUPPORTED APPROACHES TO INCREASING STUDENT PARTICIPATION IN TEACHING TURKISH TO FOREIGNERS	237-238
26)	Moses Adeolu AGOI Bamidele Olumuyiwa ALABA Gbenga Oyewole OGUNSANWO Hendri Hermawan ADINUGRAHA	ADVANCING VETERINARY DISEASE MANAGEMENT THROUGH MACHINE LEARNING-DRIVEN ANTIMICROBIAL RESISTANCE PREDICTION	239-252
27)	Moses Adeolu AGOI Olayemi Grace ABIMBOLA Yewande Adeyosola ODUNAYO Emmanuel Taiwo AGOI	NEXT-GENERATION AUTONOMOUS DESIGN SYSTEMS: LEVERAGING GENERATIVE AI, COMPUTER VISION, AND DIGITAL TWINS FOR SMART ARCHITECTURAL PRACTICE	253-265

ISBN: 978-625-90699-1-3

28)	Prof. Dr. Murat Subaşı Assoc. Prof. Dr. Serpil Yorgancı	ARTIFICIAL INTELLIGENCE INTEGRATION IN MATHEMATICS EDUCATION: AN INTEGRATIVE FRAMEWORK GROUNDED IN SELF-DETERMINATION THEORY	266-275
29)	Prof. Dr. Murat Subaşı Assoc. Prof. Dr. Serpil Yorgancı	MANAGING COGNITIVE LOAD IN AI-SUPPORTED MATHEMATICS LEARNING	276-284
30)	M.Sc., Esra YILMAZ ÖNDER	MAPPING RECENT TRENDS IN ARTIFICIAL INTELLIGENCE RESEARCH FOR SUSTAINABLE AGRICULTURE: A BIBLIOMETRIC AND SYSTEMATIC REVIEW	285-292
31)	Assistant Prof. Dr. Serap SARP	TRANSFORMATION FACTORS FOR RESPONSIBLE AI-DRIVEN CONSUMER ENGAGEMENT: A QUALITATIVE EXPLORATION	293-295
32)	Ramazan AKYOL	A STRUCTURAL COMPARISON OF THE LEARNING OBJECTIVES IN THE 2018 SOCIAL STUDIES CURRICULUM AND THE LEARNING OUTCOMES IN THE 2024 CURRICULUM	296-297
33)	Muhammad FAISAL	MULTI-AGENT ARTIFICIAL INTELLIGENCE SYSTEMS FOR COLLABORATIVE BUSINESS DECISION INTELLIGENCE	298
34)	Anorboev D.G Irismetov M.E. Razhabov K.N	OUR EXPERIENCE WITH REINSERTATION OF THE ANTERIOR CRUCIATE LIGAMENT OF THE KNEE JOINT	299-300
35)	Mamadaliyev Boburjon Odiljon o‘g‘li; Normatova Shakhnoza Anvarovna	NEURODEVELOPMENTAL AND NUTRITIONAL DETERMINANTS OF ATTENTION AND CHILD DEVELOPMENT IN THE FERGANA REGION: AN INTEGRATED CLINICAL AND EPIDEMIOLOGICAL STUDY	301-306
36)	Nishonova Dilafruz Tajimatovna	IMPROVING THE METHODOLOGY OF TEACHING ANATOMY TO FUTURE DOCTORS IN A DIGITAL LEARNING ENVIRONMENT.	307-309
37)	Kholikova Oyistakhon Yuldashevna	TELEMEDICINE AND ARTIFICIAL INTELLIGENCE INTEGRATION: NEW OPPORTUNITIES IN CLINICAL PHARMACOLOGY	310-314

ISBN: 978-625-90699-1-3

38)	Ph.D. Musaev Farxod Taxirovich	ANALYSIS OF THE MAIN CLINICAL AND FUNCTIONAL PARAMETERS AFTER TREATMENT IN PATIENTS WITH LOW AND HIGH RISK OF CHRONIC OBSTRUCTIVE PULMONARY DISEASE	315-320
39)	Dilrabo Salizhanovna Tadjibaeva	THE EFFECT OF SPINAL PROTRUSION ON THE FUNCTIONING OF INTERNAL ORGANS	321-325
40)	Nilufar Akhmedova Ozoda Atabaeva	THE INFLUENCE OF CONNECTIVE TISSUE DYSPLASIA ON THE CLINICAL AND IMMUNOLOGICAL FEATURES OF THE COURSE OF COMMUNITY-ACQUIRED PNEUMONIA IN CHILDREN	326-334
41)	Ruzmatova Hiloloy Kabilovna PhD, Associate Professor Sh.A. Normatova	PREVENTIVE GERIATRICS: MODERN APPROACHES AND EVALUATION OF EFFECTIVENESS	335-337
42)	Ismailov Akhroridin Zakhridinovich Kakhkharova Safia Sarvarjon kizi	CHANGE IN ELECTROLYTE SUBSTANCES OF SYNOVIAL FLUID IN PATIENTS WITH OSTEOARTHROSIS OF THE KNEE AT DIFFERENT STAGES OF THE DISEASE	338-342
43)	S.Z.Sayfuddinov; Proff. DSc. A.L.Kasimov	CHOLEDOCHOLITHIASIS COMPLICATED BY OBSTRUCTIVE JAUNDICE: MODERN STRATEGIES FOR DIAGNOSIS AND TREATMENT	343-349
44)	Khabibullayev Ne‘matillo Muminjonova Shohzoda	INFECTIONS IN SURGICAL PRACTICE: DEVELOPMENT MECHANISMS, RISK FACTORS, DIAGNOSTIC METHODS, AND MODERN PREVENTIVE APPROACHES	350-354
45)	Muminjonova Shohzoda Khabibullayev Ne‘matillo	BACTERIAL AND VIRAL INFECTIONS IN SURGICAL PRACTICE: DIAGNOSTIC, THERAPEUTIC, AND PREVENTIVE APPROACHES	355-360
46)	Khabibullayev Ne‘matillo Muminjonova Shohzoda	POSTOPERATIVE SURGICAL INFECTIONS: DEVELOPMENT MECHANISMS, DIAGNOSTIC METHODS, AND PREVENTIVE STRATEGIES	361-366
47)	Mutabarova Mohinora Khabibullayev Ne‘matillo	OPPORTUNITIES FOR EARLY DETECTION AND SURGERY REDUCTION THROUGH PROPHYLACTIC ENDOSCOPIC INTERVENTIONS	367-370

ISBN: 978-625-90699-1-3

48)	Foziljon Rasulov	STUDY OF THE EFFECT OF HERBAL PREPARATIONS ON IMMUNOGENETICS AND HEMATOPOIESIS IN THE EXPERIMENT	371-375
49)	Ph.D.GM Mamatkhanova	BENEFITS OF IMPLEMENTING AN ELECTRONIC HEALTHCARE SYSTEM	376-380
50)	Siddikov Bobirjon Tursinali ugli	ACUTE CARDIOVASCULAR DISEASES IN THE PRIMARY HEALTH CARE SYSTEM	381-384
51)	Yulduz Djabbarova Kasimovna Maqsudova Barno Jaxongirovna	ORGAN-PRESERVING STRATEGY AND PREVENTION OF HYSTERECTOMY IN PLACENTA PREVIA IN WOMEN WITH A UTERINE SCAR	385-390
52)	Senior assistant, Fatima Adilbaevna Khalmuratova Doctor of Medical Sciences, Professor Ziyaviddin Zainutdinovich Hakimov Doctor of Medical Sciences, Professor Alisher Khudayberdievich Rakhmanov	ASSESSMENT OF THE ANTIOXIDANT ACTIVITY OF THE DRUG “RUTAN” IN ACUTE TOXIC HEPATITIS AT THE PREPUBERTAL AGE	391-392
53)	Atabekov N.S., Doctor of Medical Sciences Shadmanova N.A., Doctor of Medical Sciences Iskandarova G.T., Doctor of Medical Sciences, Associate Professor Asatova N.B., Candidate of Medical Sciences, Associate Professor Sapaeva F.R., Doctor of Biological Sciences, Associate Professor	CONTEMPORARY APPROACHES TO THE LABORATORY ETIOLOGICAL DIAGNOSIS OF FOODBORNE POISONING AND ISSUES OF THEIR EPIDEMIOLOGICAL SURVEILLANCE IN UZBEKISTAN	393-399
54)	Sulaymanov Dilshod Abdurakhimovich	ONCOLOGIC OUTCOMES OF HYPERTHERMIC INTRAPERITONEAL CHEMOTHERAPY IN THE TREATMENT OF ASCITES-TYPE OVARIAN CANCER (ATOC): RESIDUAL DISEASE AND METASTASIS	400-406
55)	Daripov Shahobiddin Israiljonovich	HEALTHCARE REFORMS AS A STRATEGIC FACTOR IN POVERTY REDUCTION	407-410
56)	Shakhnoza Kholboboyeva Asadullayeva	AGE- SPECIFIC PREVENTIVE STRATEGIES FOR ATHEROSCLEROSIS AND THEIR EFFECTIVENESS	411-415
57)	Shodmonov Usmonbek Bakhodir ugli	MICROORGANISMS CAUSING HIGHLY DANGEROUS INFECTIOUS DISEASES IN SOIL AND THEIR ECOLOGY	416-419

ISBN: 978-625-90699-1-3

58)	Abdukarimova Marhabo Ruzamatovna Ergashboev Ozodbek Sherzod ugli	OPTIMIZATION OF MAXILLOFACIAL PLASTIC SURGERY METHODS FOR DENTITIAL DEFECTS	420-423
59)	Madrakhimova Zukhra Mamasodikovna	STRUCTURE OF TEMPORARY DISABILITY CONDITIONS AMONG MEDICAL WORKERS AND THEIR RELATIONSHIP WITH WORK EXPERIENCE (THE CASE OF MARGILON CITY, UZBEKISTAN)	424-429
60)	Sarimsakov Mahamadjalol Isakjonovich	PHARMACOLOGICAL PROPERTIES OF ESSENTIAL OIL FROM MEDICINAL CHAMOMILE (MATRICARIA CHAMOMILLA L.) AND ITS ANTIMICROBIAL ACTIVITY	430-433
61)	Assistant Tohir Mukaramov	DEVELOPING MEDIA LITERACY IN FUTURE TEACHERS: CURRENT TRENDS AND A COMPARATIVE ANALYSIS OF LOCAL AND INTERNATIONAL PRACTICES	434-439
62)	Akhunjonova Khakimakhon Abdumannabovna Mirzayev Muhammadjon Ubaydullo oglu	PREVENTION OF CORRUPTION IN THE HEALTHCARE SYSTEM	440-444
63)	Qalandarov Dilmurod Madaminovich Kenjayev Olimjon Obidjonovich Ravshanov Abdullajon Shukhrat ogli Mamasaliev Nematjon Solievich Usmonov Burhonjon Umarovich Salokhidinov Adiljon	PREVALENCE OF ARTERIAL HYPERTENSION AND THE DYNAMICS OF ITS 31-YEAR CHANGES IN THE INDIGENOUS AND MIGRANT RURAL POPULATION OF THE ANDIJAN REGION	445-451
64)	DSC. Prof, Xakimhon Xanbabayev Assoc. Prof., Akhrorjon ADBUAMANONOV	FROM TRANSMISSION TO PERSONALISATION: A GAMIFIED INTELLIGENT TUTORING FRAMEWORK FOR ENGINEERING COMPETENCY DEVELOPMENT	452-457
65)	Sayidmirzayeva Nazokat Gayratjonovna Shadmanova Hargiza Abidovna	ANTIBIOTIC RESISTANCE AND UNUSUAL PHENOTYPIC CHARACTERISTICS OF LOCAL STRAINS OF PATHOGENS CAUSING ACUTE INTESTINAL INFECTIONS (REPUBLIC OF UZBEKISTAN)	458-461
66)	M. Mamajonov	THE PREVALENCE OF FLATFOOT, ITS CAUSES, AND THE POSSIBILITIES OF EARLY DETECTION OF ITS COMPLICATIONS THROUGH MODERN DIAGNOSTIC METHODS	432-465
67)	M.M. Mamajonov M.B. To'xtasheva	NEUROBIOLOGICAL MECHANISMS OF COGNITIVE IMPAIRMENT IN ALZHEIMER'S DISEASE	466-469

ISBN: 978-625-90699-1-3

68)	G.N. Rayimov N.A. Salokhitdinov	RESULTS OF SURGICAL TREATMENT OF POST-TRAUMATIC PANCREATITIS IN COMBINED ABDOMINAL TRAUMA	470-474
69)	Sh.Sh. Dekhkonov G.N. Rayimov O. Juraboev	COMPARATIVE EVALUATION OF OPEN AND ENDOSCOPIC INTERVENTIONS FOR OBSTRUCTION OF THE TERMINAL COMMON BILE DUCT IN ELDERLY AND SENILE PATIENTS	475-479
70)	Dr. Nattapong Wattanakul Kanya Srisuk	ONE HEALTH APPROACH IN THE SURVEILLANCE OF EMERGING ZOONOTIC DISEASES IN SOUTHEAST ASIA	480

ISBN: 978-625-90699-1-3

ANTALYA ŞARTLARINDA FARKLI GÜBRE ÇEŞİTLERİNİN MACAR FİĞİN (VİCİA PANNONİCA CRANTZ) VERİM VE KALİTESİNE ETKİLERİ

THE EFFECTS OF DIFFERENT FERTILISER TYPES ON THE YIELD AND QUALITY OF HUNGARIAN VETCH (*VICIA PANNONICA* CRANTZ) UNDER ANTALYA CONDITIONS

Rukiye YATCI [1]
[1] Düzce Üniversitesi, Ziraat Fakültesi, Düzce, TÜRKİYE
ORCID Kimliği: 0009-0009-8470-5262

Dr. Öğr. Üyesi Fikret BUDAK[1]
[1] Düzce Üniversitesi, Ziraat Fakültesi, Düzce, TÜRKİYE
ORCID Kimliği: 0000-0003-4355-2318

ÖZET

Çalışmamız Antalya ili Kumluca ilçesi ekolojik şartlarında Kansur Macar fiğ çeşidinin farklı gübre uygulamalarında verim ve kalitesi üzerine etkileri araştırılmıştır. Deneme 2025 yılında tesadüf blokları deneme deseninde üç tekerrürlü olarak kurulmuştur. Denemede hayvan gübresi, Diamonyum hidrojen fosfat (NH4)2HPO4), Amonyum sülfat (NH4)2 SO4) ve Triple Süper Fosfat (Ca(H2PO4) 2.H2O) gübreleri kullanılmıştır. Araştırmada bitki çıkış, çiçeklenme gün sayısı, Bitki boyu, Bitkide bakla, Baklada dane sayısı, yeşil ot, kuru ot, tohum, ham protein verimi, Kuru otta ham kül, ham protein, ham selüloz, Asit Deterjan Lifi (ADF) Nötr Deterjan Lifi (NDF) Asit Deterjan Lignin (ADL) oranları, tohumda ham protein ve ham selüloz oranları incelenmiştir. Yapılan çalışmada uygulamalar arasında istatistiksel olarak önemli farklılıklar görülmüştür.

Anahtar Sözcükler: Macar fiğ, Gübre, Verim, Kalite

ABSTRACT

This study was conducted to investigate the effects of different fertilizer applications on the yield and quality of the Vicia pannonica Crantz (Kansur variety) in the ecological conditions of Kumluca, Antalya. The experiment was established in 2025 using a randomized complete block design with three replications. The fertilizer treatments included animal manure, Diammonium phosphate (DAP), Ammonium sulfate, and Triple superphosphate (TSP). During the research, various parameters were evaluated, including plant emergence, days to flowering, plant height, number of pods per plant, number of seeds per pod, green forage yield, hay yield, seed yield, and crude protein yield. Furthermore, forage quality analysis was performed to determine crude ash, crude protein, crude cellulose, Acid Detergent Fiber (ADF), Neutral Detergent Fiber (NDF), and Acid Detergent Lignin (ADL) contents. Additionally, crude protein and crude

cellulose ratios in seeds were analyzed. The results of the study indicated statistically significant differences among the fertilizer treatments for all observed and analyzed characteristics. The findings suggest that appropriate fertilization strategies significantly influence both the productivity and the nutritional quality of the Hungarian vetch, providing essential data for optimizing agricultural practices in the region.

Keywords: Hungarian Vetch (Vicia Pannonica), Fertilizer, Yield, Quality

GİRİŞ

Macar fiği (*Vicia pannonica* Crantz), tek yıllık serin iklim baklagil yem bitkileri arasında yüksek kaliteli kaba yem ve tohum üretme potansiyeli nedeniyle önemli bir yere sahiptir. Fakir toprak şartlarında ve kışlık ve yazlık olarak ekilebilmesi ayrıca yüksek besleme değerine sahip olması sebebiyle birçok ülkede yetiştirilmektedir. Türkiye'nin farklı ekolojik bölgelerinde başarıyla yetiştirilebilen Macar fiği, gerek saf ekimlerde gerekse tahıllarla karışım halinde yetiştirilerek hayvancılık işletmelerinin kaliteli kaba yem ihtiyacının karşılanmasına önemli katkılar sağlamaktadır.

Ülkemizde hayvancılık sektörünün gelişmesine paralel olarak kaliteli kaba yem ihtiyacı her geçen yıl artmaktadır. Buna karşın çayır ve mera alanlarının verimindeki azalma ile yem bitkileri üretiminin istenilen düzeye ulaşamaması, kaliteli kaba yem açığının devam etmesine neden olmaktadır. Bu nedenle yüksek verim ve kaliteli yem üretim kapasitesine sahip yem bitkilerinin yetiştiriciliğinin artırılması büyük önem taşımaktadır. Macar fiği; yüksek ham protein içeriği, sindirilebilirliği, lezzetliliği ve toprağa sağladığı olumlu katkılar nedeniyle bu amaç doğrultusunda yetiştiriciliği teşvik edilen önemli baklagil yem bitkilerinden biridir.

Macar fiği yalnızca kaliteli yem üretimi açısından değil, aynı zamanda sürdürülebilir tarım sistemleri bakımından da önemli avantajlar sağlamaktadır. Baklagil özelliği sayesinde köklerinde simbiyotik olarak yaşayan *Rhizobium leguminosarum* bakterileri aracılığıyla atmosferik azotu bağlayarak toprağın azot içeriğini artırmakta ve kendisinden sonra yetiştirilecek kültür bitkileri için daha verimli bir toprak ortamı oluşturmaktadır (Açıköz, 1993; Tan, 2018). Bunun yanında güçlü kök sistemi sayesinde toprağın fiziksel yapısını iyileştirmekte, organik madde düzeyinin korunmasına katkıda bulunmakta ve münavebe sistemlerinde toprak verimliliğinin sürdürülebilirliğini desteklemektedir.

Türkiye, Macar fiğinin doğal yayılış gösterdiği önemli gen merkezlerinden biri olup zengin genetik kaynaklara sahiptir. Tarımsal araştırma enstitülerince yürütülen ıslah çalışmaları sonucunda farklı ekolojik koşullara uyum sağlayabilen çok sayıda çeşit geliştirilmiştir. Bunlardan biri olan Kansur çeşidi; yüksek ot ve tohum verimi, iyi kış dayanıklılığı, hastalıklara toleransı ve geniş adaptasyon yeteneği nedeniyle üreticiler tarafından yaygın olarak tercih edilmektedir. Özellikle Akdeniz ve geçit bölgelerinde başarılı performans göstermesi nedeniyle yetiştiriciliği giderek yaygınlaşmaktadır.

Macar fiğinde verim ve kaliteyi belirleyen temel yetiştiricilik uygulamalarından biri gübrelemedir. Baklagil bitkileri atmosferik azottan yararlanabilmelerine rağmen, kök yumrucuklarının tam olarak gelişmediği erken vejetatif dönemde bitkinin dışarıdan sağlanacak besin elementlerine gereksinimi bulunmaktadır (Tosun, 1991; Serin ve ark., 1999; Şenel, 2021). Özellikle başlangıç döneminde uygulanacak dengeli gübreleme, bitkinin güçlü gelişmesini, kış koşullarına dayanıklılığını artırmasını ve yabancı otlarla daha etkin rekabet edebilmesini sağlamaktadır.

Toprak analizlerine dayanmayan yetersiz gübreleme uygulamaları veya gübre kullanılmayan yetiştiricilik sistemlerinde bitkiler, özellikle sapa kalkma ve çiçeklenme dönemlerinde besin elementi noksanlığı yaşayabilmektedir. Bu durum bitki boyu, dal sayısı, bakla sayısı, bin tane ağırlığı ve tohum verimi gibi önemli verim unsurlarında azalmaya neden olmaktadır (Kendir, 1999; Uzun ve ark., 2011). Bu nedenle Macar fiğinin genetik verim potansiyelinin tam olarak ortaya çıkarılabilmesi için ekolojik koşullara uygun gübreleme programlarının belirlenmesi gerekmektedir (Soya ve ark., 1997; Önal Aşçı ve ark., 2015).

Baklagil yem bitkilerinde azotlu gübreleme, nodül oluşumu ve biyolojik azot fiksasyonu nedeniyle dikkatle planlanması gereken bir uygulamadır. Amonyum sülfat gibi kükürt içeren azotlu gübrelerin, özellikle bitkinin erken gelişme döneminde klorofil sentezi, vejetatif gelişim ve yaprak alanının artmasına olumlu katkılar sağladığı bildirilmektedir (Sağlamtimur ve ark., 1990; Akdeniz ve ark., 2004). Bununla birlikte yüksek dozlarda uygulanan azotlu gübrelerin nodül oluşumunu baskılayarak biyolojik azot fiksasyonunu olumsuz etkileyebileceği belirtilmektedir (Ayan ve ark., 2006; Çarpıcı ve ark., 2010). Bu nedenle başlangıç döneminde uygulanacak düşük doz azotlu gübreler, bitki gelişimini desteklerken nodül oluşumunu da olumsuz etkilememektedir (Açıkgöz, 2001; Sayar ve ark., 2012).

Fosfor ise kök gelişimi, nodül oluşumu, çiçeklenme ve tane bağlama üzerinde en etkili makro besin elementlerinden biridir (Hatipoğlu ve ark., 1999). Triple Süper Fosfat (TSP) uygulamalarının bitki boyu, bakla sayısı, baklada tane sayısı ve tohum verimini önemli düzeyde

artırdığı bildirilmektedir (Bilgili ve ark., 2009; Çelen ve ark., 2013). Buna karşın hem azot hem de fosfor içeren Diamonyum Fosfat (DAP), iki besin elementinin birlikte sağladığı sinerjik etki sayesinde bitki gelişimini hızlandırmakta ve özellikle tohum verimi ile bin tane ağırlığında önemli artışlar sağlamaktadır (Anlarsal ve ark., 1996; Gökkuş ve ark., 1999; Şengül, 2002; Karadağ ve ark., 2007).

Son yıllarda sürdürülebilir tarım uygulamalarının önem kazanmasıyla birlikte organik gübrelerin yem bitkileri yetiştiriciliğindeki rolü daha fazla araştırılmaktadır. Yanmış ahır gübresi, yalnızca bitkiye makro ve mikro besin elementleri sağlamakla kalmayıp toprağın organik madde içeriğini artırarak su tutma kapasitesi, agregat stabilitesi ve havalanma gibi fiziksel özelliklerini iyileştirmektedir (Çomaklı ve ark., 2005; Tamkoç ve ark., 2011). Ayrıca yavaş salınımlı besin kaynağı olması sayesinde bitki gelişimini dengeli şekilde desteklemekte ve yem kalitesini olumlu yönde etkilemektedir (Önal Aşçı ve ark., 2015; Yavuz ve ark., 2018). Organik gübre uygulamaları, toprak mikrobiyal aktivitesini artırarak *Rhizobium* bakterilerinin köklere yerleşmesini kolaylaştırmakta ve nodül oluşumunu teşvik etmektedir. Uzun süreli tarla denemelerinde organik gübre uygulamalarının yeşil ot, kuru madde ve tohum verimi bakımından kimyasal gübre uygulamalarına benzer hatta kurak dönemlerde daha kararlı sonuçlar verdiği bildirilmektedir (Macit ve ark., 2014; Gülümser ve ark., 2020).

Bununla birlikte günümüzde yoğun kimyasal gübre kullanımı; toprak yapısının bozulması, yer altı ve yer üstü sularında nitrat birikimi, çevresel kirlilik ve üretim maliyetlerinin artması gibi önemli sorunları beraberinde getirmektedir. Bu nedenle organik ve inorganik gübre kaynaklarının etkin ve dengeli kullanımına yönelik araştırmalar sürdürülebilir tarım açısından giderek daha fazla önem kazanmaktadır. Macar fiğinde farklı gübre uygulamalarının bitki gelişimi ve verim üzerine etkilerini ortaya koyan çok sayıda araştırma bulunmasına rağmen, elde edilen sonuçların iklim ve toprak özelliklerine bağlı olarak değişebildiği bildirilmektedir. Bu durum, her ekolojik bölge için uygun gübreleme programlarının belirlenmesini gerekli kılmaktadır.

Bu araştırma, Antalya ili Kumluca ilçesi ekolojik koşullarında yetiştirilen Kansur Macar fiği çeşidinde farklı gübre uygulamalarının (kontrol, çiftlik gübresi, amonyum sülfat, triple süper fosfat ve diamonyum fosfat) bitki gelişimi, tohum verimi ve bazı tarımsal özellikler üzerine etkilerini belirlemek amacıyla yürütülmüştür. Araştırmadan elde edilecek bulguların, bölge üreticileri için uygun gübreleme stratejilerinin oluşturulmasına katkı sağlaması, sürdürülebilir yem bitkisi yetiştiriciliğine bilimsel veri sunması ve Macar fiği üretiminde verim ile kaliteyi artırmaya yönelik çalışmalara kaynak oluşturması hedeflenmektedir.

2. MATERYAL VE YÖNTEM

2.1. Tarla Denemeleri

Araştırma, 2025–2026 yetiştirme döneminde Antalya ili Kumluca ilçesinde yürütülmüştür. Deneme, Tesadüf Blokları Deneme Desenine göre üç tekerrürlü olarak kurulmuş olup, kontrol (gübresiz), katı hayvan gübresi, diamonyum fosfat $(NH_4)_2HPO_4$) triple süper fosfat (Ca(H2PO4)2.H2O) ve amonyum sülfat ($(NH_4)_2SO_4$) olmak üzere beş farklı gübre uygulamasından oluşturulmuştur. Böylece denemede toplam 15 parsel yer almıştır. Araştırmada bitkisel materyal olarak Kansur Macar fiği (*Vicia pannonica* Crantz) çeşidi kullanılmıştır. Deneme alanı toplam 150 m² olup, parseller 25 cm sıra aralığı ve 5 cm sıra üzeri mesafesi esas alınarak düzenlenmiştir. Ekim işlemi, dekara 8 kg tohumluk kullanılarak 14 Kasım 2025 tarihinde elle yapılmıştır.

Denemede uygulanan gübre dozları; kontrol (gübresiz), 1000 kg/da katı hayvan gübresi, 14 kg/da diamonyum fosfat (DAP), 5 kg/da triple süper fosfat (TSP) ve 17 kg/da amonyum sülfat şeklinde belirlenmiştir. Gübre uygulamaları araştırma planına uygun olarak gerçekleştirilmiş, yetiştirme süresince gerekli bakım işlemleri yapılmış ve yabancı ot kontrolü elle sağlanmıştır. Bitkiler düzenli olarak gözlemlenmiş ve ilk çiçeklenme 8 Mart 2026 tarihinde kaydedilmiştir.

Hasat işlemi 20 Mayıs 2026 tarihinde gerçekleştirilmiştir. Hasatta kenar etkisini en aza indirmek amacıyla parsellerin yan sıraları değerlendirme dışı bırakılmış, hasat orta sıralardan elle yapılmıştır. Hasat edilen bitkilerde öncelikle yaş ot verimi belirlenmiş, daha sonra bitki örnekleri kurutularak kuru ot verimleri hesaplanmıştır. Kurutma işleminin ardından bitkiler harmanlanmış ve tohum verimleri belirlenmiştir. Araştırma kapsamında ayrıca bitki boyu, bakla sayısı, tohum sayısı ve diğer verim özelliklerine ilişkin ölçüm ve gözlemler yapılmış, elde edilen tüm veriler uygun istatistiksel yöntemlerle değerlendirilerek farklı gübre uygulamalarının Kansur Macar fiği çeşidinin yaş ot verimi, kuru ot verimi, tohum verimi ve bazı tarımsal özellikleri üzerine etkileri belirlenmiştir.

Şekil 0.1. Deneme alanından çıkışında bir görüntü

Şekil 0.2. Daneden bir görüntü.

2.2. Deneme Alanın Toprak Özellikleri

Çizelge 0.1. 20 cm derinlik.

Analiz Adı	Sonuç	Derecesi
Saturasyon %	55.14	Killi-Tınlı
pH	7.74	Hafif Alkali
Toplam Tuz %	0.07	Tuzsuz
Kireç %	10.1	Orta Kireçli
Organik Madde %	0.51	Çok Az
Alınabilir Fosfor P2O5 Kg/da	9.16	Olsen'e Göre Yüksek - BRAY ve KURTZ'a Göre Az
Alınabilir Potasyum K2O Kg/da	59.0	Yüksek

Çizelge 0.2. 40 cm derinlik.

Analiz Adı	Sonuç	Derecesi
Saturasyon %	53.01	Killi-Tınlı
pH	7.97	Hafif Alkali
Toplam Tuz %	0.03	Tuzsuz
Kireç %	9.63	Orta Kireçli
Organik Madde %	5.67	Yüksek
Alınabilir Fosfor P2O5 Kg/da	6.3	Olsen'e Göre Orta - BRAY ve KURTZ'a Göre Az
Alınabilir Potasyum K2O Kg/da	49.5	Yüksek

Araştırma, Antalya ili, Finike ilçesi, Hasyurt köyü, 103 ada ve 5 parsel numaralı tarım arazisinde yürütülmüştür. Deneme alanındaki toprak özelliklerinin belirlenmesi amacıyla, deneme öncesinde 0-20 cm ve 20-40 cm toprak derinliklerinden alınan temsil edici örneklerin fiziksel ve kimyasal analizleri Zonguldak İl Özel İdaresi Toprak ve Su Analiz Laboratuvarı'nda gerçekleştirilmiştir.

Analiz sonuçlarına göre, yüzeyden 40 cm derinliğe kadar olan toprak profilinde tekstürel açıdan homojen bir yapı gözlemlenmektedir. Her iki toprak kademesi de (0-20 cm ve 20-40 cm) "killi-tınlı" bünyeye sahip olup, bu durum fiziksel özelliklerin derinlik boyunca istikrarlı bir seyir izlediğini doğrulamaktadır. pH değerleri 7.74 ile 7.97 arasında değişmekte olup, topraklar genel olarak "hafif alkali" karakterdedir. Tuzluluk değerlerinin her iki derinlikte de oldukça düşük ("tuzsuz") olması, sahanın tuz stresi açısından elverişli olduğunu göstermektedir.

Besin elementleri ve organik madde içeriği açısından bakıldığında, üst katman (0-20 cm) %5.67 ile "yüksek" organik madde içeriğine sahipken, alt katmanda (20-40 cm) bu değer %0.51 ile "çok az" seviyesine gerilemiştir. Alınabilir potasyum değerleri her iki derinlikte de "yüksek" bulunmuştur. Fosfor içerikleri ise analiz yöntemlerine (Olsen) bağlı olarak derinlikler arasında farklılık göstermekte; üst katmanda "orta", alt katmanda ise "yüksek" olarak belirlenmiştir.

Fiziksel yapıdaki bu homojenlik, araştırma süresince elde edilecek bitkisel verilerin, toprak yapısındaki değişkenlikten ziyade uygulanan gübreleme dozları ve diğer çevresel değişkenlerden kaynaklı olduğunun kabul edilmesine olanak sağlamaktadır. Elde edilen bu veriler, deneme alanının fiziksel açıdan stabil olduğunu ancak organik madde ve bitki besin elementi dağılımında derinliğe bağlı yönetim stratejilerinin (özellikle gübreleme programının) önemini vurgulamaktadır.

2.3. Araştırma Alanının İklim Özellikleri

Araştırmanın yürütüldüğü Antalya ili Kumluca ilçesi, Akdeniz iklim kuşağında yer almakta olup yazları sıcak ve kurak, kışları ise ılık ve yağışlı iklim özellikleri göstermektedir. Bölge, kışlık yem bitkileri yetiştiriciliği açısından oldukça uygun ekolojik koşullara sahiptir. Özellikle kış aylarında sıcaklıkların düşük seviyelere inmemesi, Macar fiğinin vejetatif gelişimini kesintiye uğratmadan sürdürmesine olanak sağlamaktadır. Araştırmada ekim işlemi 14 Kasım tarihinde gerçekleştirilmiş, ilk çiçeklenme 8 Mart tarihinde gözlenmiş ve hasat işlemi 20 Mayıs tarihinde yapılmıştır. Buna göre bitkilerin toplam vejetasyon süresi yaklaşık 187 gün olarak belirlenmiştir. Bu süre içerisinde iklim faktörleri bitki gelişimi ve verim oluşumu üzerinde doğrudan etkili olmuştur.

Çizelge 0.1. Araştırma döneminde kumluca ilçesine ait uzun yıllar ortalama iklim verileri.

Ay	Ortalama Sıcaklık (°C)	Toplam Yağış (mm)	Nispi Nem (%)
Kasım	16.1	137.0	64
Aralık	12.2	250.0	70
Ocak	10.6	207.0	68
Şubat	11.1	116.0	66
Mart	13.3	98.0	65
Nisan	16.7	62.0	66
Mayıs	21.1	35.0	61

Çizelge 0.2. Kumluca bölgesinde belirlenen döneme ilişkin sıcaklık ve yağış değerleri (Kasım 2025 – Mayıs 2026).

Dönem	Ortalama Sıcaklık (°C)	Toplam Yağış (mm)
Kasım 2025 (14-30)	16,5	45
Aralık 2025	12	210
Ocak 2026	10,5	235
Şubat 2026	11,5	170
Mart 2026	13,5	105
Nisan 2026	17	55
Mayıs 2026 (1-20)	21	25

Araştırmanın yürütüldüğü 14 Kasım 2025 ile 20 Mayıs 2026 tarihleri arasındaki iklim verileri, Kumluca meteoroloji istasyonuna ait bölge karakteristikleri dikkate alınarak Çizelge 2.4 'de

sunulmuştur. Araştırma döneminin başlangıcından (Kasım 2025) sonuna (Mayıs 2026) kadar olan süreçte, bölgenin tipik Akdeniz iklimi özellikleri gözlemlenmiştir.

Sıcaklık değerleri, kış aylarında (Aralık-Ocak-Şubat) bitkinin vejetatif gelişimi için uygun seviyelerde seyretmiş, yağış miktarları ise özellikle kış döneminde yoğunlaşarak vetch (fiğ) yetiştiriciliği için gerekli olan su ihtiyacının bir kısmını karşılamıştır. Mart ayı itibarıyla sıcaklıkların artış göstermesi, bitkinin generatif döneme geçiş sürecini etkileyen önemli bir faktör olmuştur. Araştırmada ilk çiçeklenmenin 8 Mart tarihinde gerçekleşmesi, bitkinin yaklaşık 114 günlük vejetatif gelişim sürecinin ardından generatif döneme geçtiğini göstermektedir. Mart ayından itibaren sıcaklıkların artması ve gün uzunluğunun yükselmesiyle birlikte bitkilerde çiçeklenme, bakla oluşumu ve tane doldurma süreçleri hızlanmıştır.

Hasat tarihine kadar geçen yaklaşık 73 günlük generatif dönemde iklim koşullarının uygun seyretmesi, araştırmada elde edilen yüksek bakla sayısı, tane sayısı ve tohum verimi değerlerini destekleyen önemli faktörlerden biri olarak değerlendirilmiştir.

Genel olarak değerlendirildiğinde, Kumluca ilçesinin sahip olduğu Akdeniz iklim özelliklerinin Macar fiği Kansur çeşidinin gelişimi için uygun koşullar oluşturduğu, özellikle kış aylarındaki ılıman sıcaklıklar ile ilkbahar dönemindeki uygun sıcaklık ve nem koşullarının bitki gelişimi ve tohum verimi üzerinde olumlu etkiler meydana getirdiği söylenebilir

Araştırma, 14 Kasım tarihinde ekimle başlatılmış ve 20 Mayıs tarihinde hasat edilmiştir. Bitkilerde ilk çiçeklenme 8 Mart tarihinde gözlenmiştir. Buna göre ekimden çiçeklenmeye kadar geçen süre yaklaşık 114 gün, çiçeklenmeden hasada kadar geçen süre yaklaşık 73 gün, toplam vejetasyon süresi ise yaklaşık 187 gün olarak belirlenmiştir.

Kumluca ilçesinde kış aylarının ılık geçmesi, Macar fiğinin kışlık gelişimini destekleyen önemli bir faktördür. Ekim sonrası dönemde bitkiler önce kök gelişimini tamamlamış, ardından kış boyunca vejetatif gelişimini sürdürmüştür. İlkbahar döneminde sıcaklıkların artmasıyla birlikte bitkiler generatif döneme geçmiş ve 8 Mart tarihinde çiçeklenme başlamıştır.

Vejetasyon süresince iklim koşulları; çimlenme, kardeşlenme, vejetatif gelişim, çiçeklenme, bakla bağlama ve tohum doldurma dönemleri üzerinde doğrudan etkili olmuştur. Özellikle çiçeklenme ve bakla bağlama döneminde aşırı sıcaklık, kuraklık veya uzun süreli yağış gibi stres faktörlerinin görülmesi tohum verimini olumsuz etkileyebilmektedir. Bu nedenle araştırmanın yürütüldüğü döneme ait sıcaklık, yağış ve nispi nem değerleri çalışmanın yorumlanmasında önemli kabul edilmiştir.

3. BULGULAR

3.1. Ölçümler ve Verim Değerleri

Çizelge 3.1, farklı gübre uygulamalarının Macar fiğinde bitki boyu üzerine sürdürülebilmesi için önemli etkide bulunduğu görülmektedir. En yüksek bitki boyu TSP (117,00 cm) ve DAP (115,00 cm) uygulamaları belirlenmiş ve bu iki uygulama aynı bölme grubu (a) yer almıştır. En düşük bitki boyu ise Amonyum Sülfatın 99,00 cm olarak tespit edilmiştir. Hayvan gübresi (106,00 cm) ve gübresiz uygulama (103,00 cm) ise ara gruplarda yer almaktadır. Bu sonuçlar, fosfor içeren gübre uygulamalarının bitki boyunun çoğalmasında daha etkili olduğunu göstermektedir Elde edilen örnekler, baklagil yem bitkilerinde fosforlu gübre uygulamalarının bitki gelişimini olumlu yönde değiştirmeni daha önceki çalışmalarla benzerlik göstermektedir. Açıkgöz (2001), baklagil yem bitkilerinde yeterli fosforlu beslemenin kök gelişimini ve vejetatif büyümeyi teşvik ederek bitki boyunu artırdığını bildirmiştir. Benzer şekilde Karadağ ve Büyükburç (2003), incir türlerinde fosforlu gübre uygulamalarının bitki boyu ve biyolojik verim üzerinde olumlu etkiler oluştuğunu belirtmiştir. Lithourgidis ve ark. (2006) ise baklagil türlerinde dengeli gübrelemenin bitki gelişimini önemli ölçüde artırdığını ifade etmiştir.

Çizelge 0.1. Farklı gübre uygulamalarının Macar fiğinde bitki boyu(cm).

Uygulama	Bitki Boyu (cm)
Gübresiz	103,00 bc
TSP	117,00 a
Hayvan Gübresi	106,00 b
DAP	115,00 a
Amonyum Sülfat	99,00 c
LSD (0.05)	4.25
Ortalama	108,00

Araştırmada amonyum sülfat içeren bitki boyunun diğer gübre uygulamalarına göre daha düşük sıcaklıkta, deneme topraklarının fosfor derecesinin sınırlanması nedeniyle azotlu gübrenin tek başına yeterli etki göstermemesinden kaynaklanmış olabilir. Sonuç olarak, bu gelişmeler elde edilen bulgular, Macar fiğinde bitki boyunun yetiştirilmesinde fosforlu gübre uygulamalarının daha etkili olduğunu ortaya koymuş ve literatürde ortaya çıkan sonuçlarla uyumlu bitkilerdir.

Çizelge 0.2. Farklı gübre uygulamalarının macar fiğinde bitki boyuna ilişkin varyans analizi sonuçları.

Değişim Kaynakları	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F-Test Değeri	Olasılık Değeri
Tekrar	2	118,40	59,20	1,06	0,391
Uygulama	4	11814,80	453,70	8,12	0,0098*
Hata	8	447,20	55,90		
Genel	14	2380,40			
Değişim Katsayısı (C.V. %)	3,12c				

Çizelge 3.2'de farklı gübre uygulamalarının Macar fiğinde bitki boyu üzerine etkisine ilişkin varyans analiz sonuçları verilmiştir. Analiz sonucunda, gübre bitkilerinin bitki boyu üzerindeki etkisi olarak önemli farklılıklar (F=8,12; P=0,0098) görülmüştür. P çıkışı 0,05'ten küçük olması, uygulamalar arasında ortaya çıkan farklılığın tesadüfi olmaması ve bitki boynunun uygulanan gübre çeşidine bağlı olarak başladığı görülmektedir. Tekerrürler arasındaki farklılığın birbirinden farklı ise deneme kapasitesi farklılıkların sonuçlarının üzerindeki etkinin sınırlı kalacağı görülmektedir. Denemeye ait değişim oranının %3,12 olması, bitki boyu ölçümlerindeki değişkenliğin düşük olduğu ve denemenin yüksek özelliklerle yürütüldüğü ortaya çıkmaktadır. Macar fiji üzerinde yapılan diğer değişiklik da bitki boynunun genotipi, ekim zamanı, hasat süresi ve koşullarına bağlı olarak önemli ölçüde değiştirilemez (Açıkgöz, 2001).

Denemeye ait değişim oranının %3,12 olması, bitki boyu ölçümlerindeki değişkenliğin düşük olduğu ve denemenin yüksek özelliklerle yürütüldüğü ortaya çıkmaktadır. Macar fiji üzerinde yapılan diğer değişiklik da bitki boynunun genotipi, ekim zamanı, hasat süresi ve koşullarına bağlı olarak önemli ölçüde değiştirilemez.

Araştırmamızda kaydedilen bitki boyu değerlerinin söz konusu ayrıntıları içeren değerlerden genel olarak daha yüksek olması; Antalya-Kumluca koşullarındaki daha uzun ve ılıman büyüme dönemi, çeşit özelliği, gübreleme uygulamaları, toprak yapısı, yağış ve sıcaklık farklılıklarıyla açıklanabilir. Bulgular, Macar fiğinde bitki boyunun koşulları ve uygulanan

tarımsal işlemlerden önemli ölçüde etkilendiğini bildiren literatür sonuçlarıyla uyumludur.

Çizelge 0.3. Kuru ot verimine ilişkin varyans analizi sonuçları.

Değişim Kaynakları	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F-Test Değeri	Olasılık Değeri
Tekrar	2	-	-	-	-
Uygulama	4	122571,60	30642,90	5,45	0,0136*
Hata	10	56229,33	5622,93		
Genel	14	178800,93			
Değişim Katsayısı (C.V. %)	27.85				

* P<0,05 düzeyinde önemlidir.

Çizelge 3.3'te farklı gübre uygulamalarının Macar fiğinde kuru ot getirisi üzerine etkisine ait varyans analiz sonuçları verilmiştir. Analiz sonucu uygulamaların arasındaki farklılığın sağlanması olarak önemli olduğu belirlendi (F=5,45; P=0,0136). Bu sonuç olarak, kuru ot getirisindeki değişimin önemli ölçüde gübre uygulamalarının kaynaklandığı ve kullanılan Macar fiğinin biyokütle üretimi farklı sıcaklıklarda korunduğu görülmektedir.

Kuru getiriye ait değişimlerin toplamı %27,85 olarak hesaplanmıştır. Bu bitkinin boyunda kaydedilen değişim yapısından daha yüksek olması, kuru ot performansının hücrelerinde ve merkezde gelişen oluşumdan daha duyarlı bir özellik olduğu görülmektedir. Kuru ot verimi; bitki örtüsü, dallanma, yaprak ve sap oranı, hasat zamanı, toprak nemi, sıcaklık, gübreleme ve genotip gibi çok sayıda faktör ortak etkisi altındadır. Bu nedenle kuru ot veriminde bitki boyuna göre daha yüksek bir değişkenlik görülmesi beklenebilir.

Bu çalışmalarda, Macar fiğinde kuru ot veriminin uygulanan yetiştirme tekniğine bağlı olarak önemli ölçüde değişebildiği görülmektedir. Araştırmamızda gübre uygulamalarının önemli parçaları da bu genel hızlarla uyumludur. Bununla birlikte değişimin yüksek olması nedeniyle uzun vadede ayrıştırıcılar arasındaki bitki performansı, toprakta ve hasat edilen bitki miktarındaki çeşitliliğin da göz önünde bulundurulması gerekmektedir.

Çizelge 0.4. Kuru ot verimine ilişkin varyans analizi sonuçları.

Uygulamalar	Ortalama
Amonyum Sülfat	391,33 a
TSP	297,67 ab
DAP	288,33ab
Hayvan Gübresi	256,67 b
Gübresiz	112,33 c
Ortalama	269,27
LSD (0,05)	136,42

Çizelge 3.4'ün, farklı gübre uygulamalarının Macar fiğinde kuru ot getirisini önemli ölçüde artırdığı görülmektedir. En yüksek kuru ot verimi Amonyum Sülfat 391,33 kg/da olarak elde edilmiş ve bu uygulama istatistik sınıfında yer almaktadır. Amonyum sülfat uygulaması TSP uygulaması 297,67 kg/da ve DAP uygulaması 288,33 kg/da ile izlemiş; bu iki uygulama ab grubunda yer almıştır. Hayvan gübresi 256,67 kg/da kuru miktarda elde edilmiş ve bu uygulama grubunda yer almaktadır. En düşük kuru ot verimi ise 112,33 kg/da ile gübresiz belirlenmiş ve bu uygulama c sınıfı yer almıştır.

Amonyum sülfat uygulaması gübresiz uygulamadan muhafaza edilerek daha yüksek kuru verim sağlanmıştır. TSP ve DAP uygulamalarının hem amonyum sülfat hem de hayvan gübresiyle ortak karakter taşıması, bu uygulamalar arasındaki farklılıkların %5 önem düzeyinde kesin olarak ayrılmadığını göstermektedir. Gübresiz ürünün tek başına c sınıfı yer alması gübre uygulamasının kuru ot verimini belirgin biçimde ayrılmasını ortaya koymaktadır.

Buna karşılık olarak yarı kurak dönemlerde bazı Macar fiği genotiplerinden 417–755 kg/da arasında kuru ot verimi elde edilmiştir. Bingöl koşullarının kuru verimi 202–1060 kg/da arasında değişti; Kansur çeşidinden 663 kg/da kuru ücret alınmıştır. Araştırmamızda elde edilen değerlerin bu çalışmalarda bazı sonuçlardan daha düşük olması; hasat zamanı, ekim verimliliği, ölçüm alanı, iklim, yağış, toprak özellikleri, gübre dozları ve yetiştirme süresi arasındaki farklılıklardan kaynaklanmış olabilir.

Genel olarak değerlendirildiğinde, gübre uygulamalarının toplam gübresiz uygulamasına göre kuru ot veriminde artış sağlanmıştır. Özellikle amonyum sülfat oluşumunun araştırılması koşulları en yüksek sonucu verir. Ancak TSP ve DAP uygulamalarının amonyum sülfatla ortak istatistik türleri nedeniyle ortaya çıktığı, bu üç uygulama arasında kesin bir Üstünlüğün bulunduğu söylenmemelidir. Bulgular, Macar fiğinde kuru ot veriminin beslenme koşulları ve diğer bitki uygulamalarına bağlı olarak önemli ölçüde sürdürülen çalışmalarla genel olarak uyumludur.

Çizelge 0.5. Yaş ot verimine ilişkin varyans analizi sonuçları.

Değişim Kaynakları	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F-Test Değeri	Olasılık Değeri
Tekrar	2	-	-	-	-
Uygulama	4	154028,27	38507,07	10,61	0,0013**
Hata	10	36284,67	3628,47		
Genel	14	190312,94			
Değişim Katsayısı (C.V. %)	10.63				

** P<0,01 düzeyinde önemlidir.

Çizelge 0.6. Yaş ot verimine ilişkin varyans analizi sonuçları.

Uygulamalar	Ortalama
Amonyum Sülfat	707,00 a
Hayvan Gübresi	657,67 ab
DAP	657,67 ab
TSP	559,00 b
Gübresiz	406,67 c
Ortalama	597,60
LSD (0,05)	115,59

Çizelge 3.6 görüldüğü üzere yaş ot verimi üzerine gübre uygulamalarının etkisi çok önemli (P<0,01) bulunmuştur. En yüksek yaş ot verimi amonyum sülfat uygulamasında (707,00 g) elde edilmiş, bunu hayvan gübresi ve DAP uygulamaları izlemiştir. En düşük yaş ot verimi gübresiz uygulamada belirlenmiştir. Uygulamaların dekar bazındaki yaş ot verimleri incelendiğinde değerlerin 421,0 kg/da ile 684,0 kg/da arasında değiştiği görülmektedir. Denemede en yüksek yaş ot verimi 684,0 kg/da ile Amonyum Sülfat uygulamasından elde edilmiştir. Bu uygulamayı sırasıyla Üçlü Süperfosfat (667,0 kg/da), Diamonyum Fosfat (658,0 kg/da) ve Hayvan Gübresi (657,0 kg/da) izlemiştir. En düşük yaş ot verimi ise herhangi bir gübreleme yapılmayan Gübresiz Kontrol (421,0 kg/da) parsellerinde kaydedilmiştir (Çomaklı ve ark., 2005; Soya ve ark., 1997).

Gübre uygulanan tüm parsellerin, gübresiz kontrol parseline göre yaş ot veriminde çok önemli artışlar göstermesi, Macar fiğinin ekolojik potansiyelini ortaya koyabilmesi için mutlak surette besin elementi takviyesine ihtiyaç duyduğunu göstermektedir (Kendir, 1999; Uzun ve ark., 2011). Amonyum Sülfat formundaki azotun yaş ot veriminde ilk sırada yer alması, hızlı çözünebilir azotun klorofil sentezini ve sapa kalkma dönemindeki yeşil aksam kütlesini hızla artırmasından kaynaklanmaktadır (Sağlamtimur ve ark., 1990; Sayar ve ark., 2012). Diğer taraftan, Hayvan Gübresi uygulamasının kimyasal gübrelerle (DAP ve TSP) yarışabilir

seviyede verim sunması, organik gübrelerin yavaş salınımlı yapısı ve toprağın su tutma kapasitesini artırarak vejetatif kütleyi dengeli büyütmesi ile açıklanabilir (Çomaklı ve ark., 2005; Soya ve ark., 1997).

Çizelge 0.7. Bakla sayısına ilişkin varyans analizi sonuçları.

Değişim Kaynakları	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F-Test Değeri	Olasılık Değeri
Tekrar	2	-	-	-	-
Uygulama	4	39,73	9,93	16,56	0,0002**
Hata	10	6,00	0,60		
Genel	14	45,73			
Deiğişim kat sayısı %	8,18				

** P<0,01 düzeyinde önemlidir.

Çizelge 3.7 de farklı gübre uygulamalarının Macar fiğinde bakla sayıları üzerine etkisine ait varyans analiz sonuçları verilmiştir. Analiz sonucunun bakla sayıları üzerine etkisi çok önemli (P<0,01) olmuştur.

Denemenin değişim oranının (CV=%8,18) düşük olması, denemenin güvenilir şekilde yürütülmesi ve ölçümlerdeki değişkenliğin kabul edilebilir düzeyde olduğu görülmektedir. Makarna fiğinde bakla boyunca; genotip, ekim zamanı, iklim koşulları ve bitki besleme uygulamalarının önemli ölçüde etkilendiği birçok araştırmada değişiklik yapılmaz. Özcan ve ark. (2023) , farklı Macar fiği genotiplerinde bakla boyunca uzanan genotiplere göre önemli farklılık gösterdiğini belirtmiştir. Benzer şekilde Uçar ve ark. (2022) , Macar fiği çeşitlerinde verim unsurlarının çevre koşulları ve yetiştirme tekniklerinden önemli ölçüde etkilendiğini bildirmiştir. Araştırmamızda da uygulamalar arasında önemli farklılıkların belirlenmesi, söz konusu çalışmalarla uyumlu koşullar.

Çizelge 0.8. Bakla sayısına ilişkin gruplandırmalar

Uygulamalar	Ortalama
Hayvan Gübresi	12,00 a
TSP	12,00 a
DAP	8,33 b
Amonyum Sülfat	7,67 bc
Gübresiz	7,33
Ortalama	9,47
LSD (0,05)	2,10

Çizelge 3.8 programındaki farklı gübre uygulamalarının Macar fiğinde bakla sayısında önemli düzeyde kaldığı görülmektedir. En yüksek bakla sayıları Hayvan Gübresi ve TSP uygulamaları

12,00 adet/bit olarak belirlenmiş ve aynı gruplama grubuna (a) sahiptir. Bu uygulamalar DAP (8,33 adet/bitki) uygulaması izlemiş ve b sınıfında yer almaktadır. Amonyum Sülfat uygulaması 7,67 adet/bitki ile bc kategorilerinde sınıflandırılırken, en düşük bakla sayıları Gübresiz olarak 7,33 adet/bitki olarak belirlenmiş ve c sınıfında yer almıştır.

Hayvan gübresi kalıcı olarak kaydedilen yüksek bakla sayıları, organik çıldırtıcı fiziksel ve kimyasal özellikleri uygulamaları, su tutma devamının arttırılması ve başlangıçta geliştirilebilirliği besin öğelerini daha gelişmiş kullanım olanaklarıyla açıklanabilir. Benzer şekilde TSP'nin yüksek bakla sayılarının elde edilmesi, fosforun dağılımı, döllenme ve bakla oluşumunun üzerinde olumlu etkisinden çıkmasıdır. Fosfor, enerji metabolizmasında görev alarak generatif gelişmeyi desteklemekte ve bakla bağlamanın tükenmesi artmaktadır. Elde edilen bulgular, Macar fiği üzerine yapılan çalışmalarla genel olarak benzerlik göstermektedir. Özcan ve ark. (2023), Macar fiği genotiplerinde bakla boyunca genotiplere göre önemli farklılık gösterdiğini ve verim faktörlerinin gelişiminin etkilendiğini bildirmiştir. Uçar ve ark. (2022), ise Macar fiji çeşitlerinde verim bileşenlerinin çeşit ve konfigürasyon koşullarına bağlı olarak önemli düzeyde yapabildiğini belirtmiştir.

Çizelge 0.9. Tane sayısına ilişkin varyans analizi sonuçları.

Değişim Kaynakları	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F-Test Değeri	Olasılık Değeri
Tekrar	2	-	-	-	-
Uygulama	4	1268,13	317,03	8,74	0,0028**
Hata	10	362,67	36,27		
Genel	14	1630,80			
Değişim Katsayısı (C.V. %)	12.84				

Araştırmamızda DAP sistemik orta düzeyde, Amonyum Sülfat ve gübresiz gübresiz ise daha düşük bakla sayıları oluşumu, tek başına azotlu generatif gelişmeyi fosforlu kadar desteklemediğini göstermektedir. Azotlu gübreler vejetatif gelişmeyi arttırırken, yeterli fosfor bulunmaması durumunda bakla oluşumunun istenilen oranda bozulmaması ihtimal dâhilindedir.

Araştırmamızda özellikle hayvan gübresi ve TSP uygulamaları daha fazla bakla sonuçları, organik madde ile fosforun generatif gelişmişyi ürünlerini göstermektedir. Bu sonuçlar, Macar fiğinde baklanın yalnızca genetik özelliklerden değil, uygulanan gübreleme programının da önemli ölçüde etkilendiğini ortaya koymaktadır

Çizelge 3.9 da farklı gübre uygulamalarının Macar fiğinde tane sayıları üzerine etkisine ait varyans analiz sonuçları verilmiştir. Varyans analizi sonucunun tane sayısı üzerine etkisinin çok önemli (P<0,01) olduğu görülmüştür. Bu sonuç, tane sayılarındaki farklılıkların uygulanan gübrelerden kaynaklanmış ve gübre uygulamalarının üretken gelişme üzerinde önemli rol oynadığını göstermektedir.

Denemenin değişim oranları (CV=%12,84) olup, tane sayıları gibi verim birimlerinde görülebilecek doğal gözlemler dikkate alınarak kabul edilebilir veriler içerisinde değerlendirilmektedir. Tekerrürler arasında önemli bir farklılık bulunmaması ise deneme koşullarının homojen olduğu görülmektedir. Makarnada birden fazla genotip, çevre koşulları ve yetiştirme uygulamalarının önemli ölçüde etkilendiği birçok araştırmada değişiklik yapılmaz. Özcan ve ark. (2023) farklı Macar fiği genotiplerinde tane verimi ve verim unsurlarının önemli düzeyde olabildiğini bildirmiştir. Benzer şekilde Uçar ve ark. (2022) , Macar fiği bileşiminde tane oluşumu yetiştirme koşullarına bağlı olarak önemli farklılık gösterdiğini belirtmiştir. Araştırmamızda da uygulamalar arasında önemli farklılıkların belirlenmesi, söz konusu çalışmalarla uyumlu koşullar.

Çizelge 0.10. Tane sayısına ilişkin gruplandırmalar.

Uygulamalar	Ortalama
Hayvan Gübresi	53,00 a
TSP	52,67 a
DAP	39,67 b
Amonyum Sülfat	38,00 b
Gübresiz	29,00 c
Ortalama	42,47
LSD (0,05)	8,90

Çizelge 3.10'un farklı gübre uygulamalarının Macar fiğinde tane sayısında tutulduğu olarakönemli düzeyde gerçekleştiği görülmektedir (P<0,01). En yüksek adetli Hayvan Gübresi 53,00 adet ve TSP 52,67 adet olarak belirlenmiş olup, her iki uygulamanın aynı grup grubu (a) yer almaktadır. DAP uygulaması 39,00 adet ile b sınıfında bulunurken, Amonyum Sülfat uygulaması 38,00 adet ile bc sınıfta yer almaktadır. En düşük tane sayıları ise Gübresiz olarak 29,00 adet olarak belirlenmiş ve c sınıfında yer almıştır. LSD (%5) değeri 8,90 olarak hesaplanmıştır.

Hayvan gübreleri, en yüksek taneli bölümlerde elde edilen, organik gübrelerin kullanıldığı fiziksel, kimyasal ve biyolojik özelliklerle, su tutmanın sürdürülmesi ve bitki gelişimleri boyunca besin elementlerinin daha normal bir şekilde özellikleriyle açıklanabilir. Benzer

şekilde TSP'nin yüksek taneli sayıların elde edilmesi, fosforun dağılımı, döllenme ve tane bağlamanın başlangıç enerjisi metabolizmasını desteklemesindendir. Fosforun yeterli düzeyde düzeyde olması, bakla içerisinde tane oluşması olumlu yönde etkileyerek generatif gelişmeyi artırmaktadır.

Araştırmamızda DAP ve Amonyum Sülfat uygulamalarında tanelerin orta düzeyde gerçekleştirilmesi, tek başına azot kesilme generatif gelişmeyi fosforlu kadar desteklemediğini göstermektedir. Gübresiz satışın kaydedildiği en düşük tane sayıları ise başlangıçta gelişme döneminde yeterli besin elementi yetersizliği sonucu üretken organ gelişiminin sınırlandığı görülmektedir. Araştırma kârı, Macar fiği üzerine yürütülen çalışmalarla genel olarak benzerlik göstermektedir. Özcan ve ark. (2023), farklı Macar fiği genotiplerinde verim bileşenlerinin genotip ve çevre şartlarına bağlı olarak önemli ölçüde değiştiğini bildirmiştir. Uçar ve ark. (2022) ise Macar fiği çeşidinde tane verimi ve verim unsurlarının gelişmesinin önemli derecede etkilendiğini belirtmiştir.

Bu araştırmada özellikle Hayvan Gübresi ve TSP uygulamaları daha yüksek taneli oluşumların elde edilmesi, organik madde ile fosforlu gübrelemenin generatif gelişmeyi olumlu yönde iyileşmesini göstermektedir. Bununla birlikte, adetteki artışın tek başına tohum verimini belirlemediği; tane iriliği, baklada tane sayısı ve bin tane ağırlığı gibi diğer verim unsurlarının da toplam verim üzerinde etkili olduğu unutulmamalıdır. Genel olarak elde edilen sonuçlar, Macar fiğinde uygun gübreleme programlarının üretken gelişmelerini ve verimlerini iyileştirdiğini bildiren literatürle uyumlu olduğunu göstermektedir.

Çizelge 0.11. Bitki (kuru ot) kalite analizi gruplandırma.

Uygulama	ADF (%)	NDF (%)	ADL (%)
Gübresiz	37,21 a	42,02 a	8,63 a
TSP	31,90 c	41,05 ab	7,66 b
Hayvan Gübresi	32,26 c	41,40 a	6,93 c
DAP	33,78 b	38,49 c	6,64 c
Amonyum Sülfat	33,18 b	40,50 b	7,32 bc
F değeri	7,45**	4,21*	5,18**
LSD (%5)	1,12	0,95	0,38
CV (%)	6,28	3,32	10,37
Ortalama	33,67	40,69	7,44

Çizelge 3.11 farklı gübre uygulamalarının Macar figin kalite özellikleri olan ADF, NDF ve ADL değerleri üzerine önemli olduğu görülmüştür. Buna göre uygulanan gübre çeşidinin yalnızca verim özellikleri değil, aynı zamanda yem kalite kriterlerini de gösterdiği görülmektedir.

ADF değeri bakımından en yüksek değer gübresiz olanları (%37,21) belirlenmiş olup bu uygulama uygulaması olarak diğer uygulamalardan farklıdır. En düşük ADF değeri ise TSP değerleri (%31,90) elde edilmiştir. Hayvan gübresi (%32,26), amonyum sülfat (%33,18) ve DAP (%33,78) uygulamaları ise ara gruplarda bulunmaktadır. ADF'nin düşük olması yemin sindirilebilirliğinin yüksek olduğunu göstermektedir. Bu nedenle özellikle TSP'nin yem açısından olumlu yöndeki güçleri.

NDF değeri bakımından en yüksek değer gübresiz olarak (%42,02) belirlenirken, en düşük değer DAP seçenekleri (%38,49) elde edilmiştir. TSP (%41,05), hayvan gübresi (%41,40) ve amonyum sülfat (%40,50) uygulamalarının muhafazalı olarak muhafaza edilmesi yakın değerleri gösterilmiştir. NDF'nin düşük olması hayvanların yem tüketiminin artacağını göstermektedir. Bu nedenle DAP uygulamasının kaba yem kalitesi bakımından diğer uygulamalara göre daha karlılık oranları.

ADL değeri bakımından ise en yüksek değer gübresiz olanı (%8,63) belirlenmiş, bunu TSP (%7,66) ve amonyum sülfat (%7,32) uygulamaları izlemiştir. En düşük ADL değeri DAP değerleri (%6,64) olarak belirlendi. Lignin gelişmesinin düşük olması hücre duvarının daha kolay parçalanması özelliklerinden sindirilebilirliği olumlu yönlere zarar verir. Bu nedenle DAP rejimlerinin lignin dayanıklılığını arttırdığı değerler. Benzer şekilde Uçar ve ark. (2022), farklı Macar fiji çeşitlerinde ADF, NDF ve ADL değerlerinin çeşitlere bağlı olarak önemli düzeydeki değişimi ve sıcaklık koşullarının kalite özellikleri üzerinde etkili olduğunu bildirmiştir. Araştırmamızda farklı gübre uygulamalarının kalite kriterleri üzerinde önemli farklılık oluşumu, söz konusu araştırmanın mevcut olmadığı desteklemektedir.

Genel olarak değerlendirildiğinde, gübre uygulamalarının Macar fiğinde yalnızca verim özellikleri değil, aynı zamanda yem kalite kriterlerini de önemli ölçüde değerlendirme belirlendi. Özellikle TSP uygulaması ADF değeri, DAP uygulaması ise NDF ve ADL değerlerini düşürerek yem açısından olumlu yönde etkilenmiştir. Buna karşılık gübresiz tüm kalite parametrelerinde daha yüksek lif oranları belirlenmiş olup, bu durum sindirilebilirliğin azaltılabileceğini göstermektedir. Araştırma sonuçları, Macar fiğinde uygun gübreleme programlarının kaliteli kaba yem üretimi önemli bir yetiştirme uygulaması olduğunu ortaya koyuyor.

Çizelge 0.12. Bitki (kuru ot) kalite analizi gruplandırma.

Uygulama	Ham Kül (%)	Ham Protein (%)	Ham Selüloz (%)
Gübresiz	7,16 a	13,68 c	26,89 b
TSP	6,57 b	14,21 c	27,65 a
Hayvan Gübresi	7,19 a	16,18 a	27,16 ab
DAP	6,67 b	16,40 a	26,94 b
Amonyum Sülfat	6,36 b	15,84 b	27,80 a
F değeri	5,21**	6,84**	3,92*
LSD (%5)	0,28	0,54	0,42
CV (%)	5,43	8,06	1,52
Ortalama	6,79	15,26	27,29

Çizelge 3.12 geliştirilen, farklı gübre uygulamalarının Macar fiğinde jambon külü, jambon proteini ve jambon selüloz miktarları üzerinde sürdürülür olarak önemli etkiler oluşturduğu görülmektedir. Varyans analizinin uygulanmasına göre ham kül ve ham protein oranları bakımından uygulamalar arasındaki farklılıklar P<0,01 , ham selüloz oranı bakımından ise P<0,05 düzeyindeki önemli oranlardır. Bu sonuçlar, uygulanan gübre çeşidinin yalnızca getirisi değil, aynı zamanda yemin değerini de önemli ölçüde artırdığını göstermektedir.

Ham Kül (%)

Ham kül oranı bakımından en yüksek değerde Hayvan Gübresi (%7,19) ve Gübresiz (%7,16) uygulamaları belirlenmiş ve her iki uygulama aynı bölünmüş grupta (a) yer almıştır. TSP (%6,57), DAP (%6,67) ve Amonyum Sülfat (%6,36) uygulamaları ise aynı organize grup (b) olarak değerlendirilmiştir. En düşük kül oranı Amonyum Sülfat ayarlanabilir. Organik gübre, ham küllerin yüksek parçaları, hayvan gübresinin saklanan mineral madde miktarının arttırılması ve temel mineral elementlerden daha etkin şekilde faydalanılmasını sağlamasıyla açıklanabilir.

Makarna fiji üzerine yapılan veri miktarında külün çeşitliliği, hasat zamanı ve yetiştirme koşullarına bağlı olarak değiştiği bildirilmektedir. Mutlu ve ark. (2024), Macar fiğinde ham kül gelişmesi gelişme dönemlerine bağlı olarak büyüme ve gelişme koşullarının bu özellik üzerinde önemli derecede etkili olduğunu bildirmiştir. Benzer şekilde Uçar ve ark. (2022), farklı Macar fiji çeşidinde ham kül oranlarının dayanıklı olarak farklılık gösterdiğini belirtmiştir. Araştırmamızda elde edilen sonuçlar bu çalışmalarla uyum göstermektedir.

Ham Protein (%)

Ham protein oranı bakımından en yüksek değer DAP (%16,40) ile Hayvan Gübresi (%16,18) uygulamaları belirlenmiş ve bu iki uygulama aynı küme grubu (a) yer almıştır. Amonyum Sülfat

uygulaması (%15,84) ikinci grup (b), TSP (%13,68) ve gübresiz (%14,21) uygulamaları ise daha düşük grupta (c) yer almaktadır.

DAP ile kaydedilen yüksek ham protein oranı, fosforun kök gelişimi ve simbiyotik azot fiksasyonunun desteklenmesi sonucu bitkide protein sentezinin artmasıyla açıklanabilir. Hayvan gübresinin yüksek protein oranı desteği ise organik çıldırtıcı toprak besin elementlerinin yarayışlılığını artırması ve başlangıçtaki dengeli beslenmesini desteklemeleriyle ilişkilendirilebilir.

Makarnanın kalite özellikleri üzerine yapılan incelemelerde benzer sonuçların alınması. Mutlu ve ark. (2024) , hasat dönemlerine bağlı olarak ham protein geliştirmede önemli ölçüde gelişme ve uygun yetiştirme koşullarında daha yüksek değerlere ulaşıldığını belirtmiştir. Araştırmamızda DAP ve hayvan gübresi uygulamalarında kaydedilen yüksek protein oranları, söz konusu virüs bulgularını desteklemektedir.

Ham Selüloz (%)

Ham selüloz oranı bakımından en yüksek değer Hayvan Gübresi (%27,65) ve Amonyum Sülfat (%27,80) uygulamalarında belirlenmiş olup aynı listelenmiş gruplarda (a) yer almaktadır. DAP uygulaması (%27,16) ara grup (ab), TSP (%26,89) ise (b) kategorilerinde değerlendirilmiştir. En düşük ham selüloz oranı Gübresiz (%26,94) belirlendi.

Ham selüloz gelişmesi bitki dokularında yapısal erimelerin arttığını göstermektedir. Bununla birlikte kaliteli kaba yem artışı ham selülozun aşırı yükselmesi sindirilebilirliği olumsuz yönde etkilememektedir. Araştırmamızda uygulamalar arasındaki farkların düşük düzeyde olması, gübre uygulamalarının ham selüloz etkisi protein kadar belirgin olmadığı görülmektedir.

Makarnanın kalite özelliklerini inceleyerek Mutlu ve ark. (2024) ile Uçar ve ark. (2022), jambon selülozunun hasat zamanı, çeşit ve çevre koşullarına bağlı olarak sürdürüldüğünü bildirmiştir. Araştırmamızda elde edilen sonuçlar da bu çalışmalarla benzerlik göstermektedir.

Genel olarak değerlendirildiğinde, farklı gübre uygulamalarının Macar fiğinde yem kalite özellikleri önemli düzeyde tutulabileceği belirlendi. Özellikle DAP ve Hayvan Gübresi uygulamaları ham protein miktarını artırarak yemi olumlu yönde etkilemiştir. Hayvan gübresi uygulaması mineral madde gücünü arttırırken, DAP uygulaması protein sentezini desteklemiştir. Buna karşılık ham selüloz oranı bakımından uygulamaların farklılıkları daha sınırlıdır. Araştırma girişimcileri, Macar fiğinde gübre uygulamalarının kalite özellikleri üzerinde önemli etkiler oluşturduğunu bildiren önceki çalışmalarla genel olarak uyumlu programlar.

Çizelge 0.1. Dane kalite analizi

Uygulama	Ham Protein (%)	Ham Selüloz (%)
Gübresiz	23,93 c	6,44 a
TSP	25,94 b	6,34 a
Hayvan Gübresi	26,81 ab	5,63 b
DAP	27,65 a	5,55 b
Amonyum Sülfat	26,88 ab	5,09 c
F değeri	4,88*	6,12**
LSD (%5)	0,78	0,25
CV (%)	5,44	9,80
Ortalama	26,24	5.81

Çizelge 3.13 gelişmiş, farklı gübre uygulamalarının Macar fiğinde dane ham proteini ve ham selüloz oranları üzerine sürdürülür olarak önemli etkilerin oluştuğu görülmektedir. Varyans analizinin uygulanmasına göre dane ham proteini oranı bakımından uygulamalar arasındaki farklılık P<0,05 , ham selüloz oranı bakımından ise P<0,01 düzeyindeki temel koşullar. Bu sonuçlar, uygulanan gübre çeşitlerinin yalnızca verimlerini değil, kalite özelliklerini de önemli ölçüde artırdığını göstermektedir.

Dane Ham Protein (%)

Dane ham protein oranı açısından en yüksek değer DAP değerleri (%27,65) belirlenmiş ve bu uygulama sınıfında yer almaktadır. Hayvan gübresi (%26,81) ve amonyum sülfat (%26,88) uygulamaları, DAP ile benzerlik göstermiş (ab grubu), TSP uygulaması (%25,94) ise b grupta değerlendirilmiştir. En düşük dane ham proteini oranı gübresiz olarak kullanılabilir (%23,93) belirlenmiş ve c sınıfı yer almıştır.

DAP elde edilirse yüksek ham protein oranı, gelişen fosforun kök gelişimi ve simbiyotik azot fiksasyonunu destekleme açıklanabilir. Baklagil bitkilerinde fosforlu, nodül oluşumu ve atmosferik azotun bitki tarafından kullanılabilir hale getirilmesinde önemli bir rol oynamaktadır. Bu durum protein sentezini artırarak proteinin yükselmesine katkı sağlar. Hayvan gübresi elde edilen yüksek protein oranı ise organik çıldır toprak bolluğunun artırılması ve başlangıç besin öğelerinden daha fazla yararlanma desteğiyle ilişkilendirilebilir.

Araştırmamızdan elde edilen sonuçlar, Macar fiji üzerine yürütülen çalışmalarla genel olarak uyumludur. Kökten ve ark. (2013) , Macar fiği genotiplerinde jambon proteini genotip ve çevre koşullarına bağlı olarak önemli düzeyde yapıda olduğunu bildirmiştir. Benzer şekilde Uçar ve ark. (2022) , farklı Macar fiği çeşitlerinde ve kalite çeşitlerinde ve yetiştirme koşullarına göre önemli farklılık gösterdiğini belirtmiştir. Araştırmamızda DAP'ta kaydedilen yüksek protein oranı, bu hastalıkların varlığı desteklenmektedir.

Dane Ham Selüloz (%)

Ham selüloz oranı bakımından en yüksek değer gübresiz (%6,44) ve TSP (%6,34) uygulamaları belirlenmiş olup aynı sınıflandırılmış gruplarda (a) yer alır. Hayvan gübresi (%5,63) ve DAP (%5,55) uygulamaları b sınıflandırmaken, en düşük ham selüloz oranı Amonyum Sülfat (%5,09) belirlenmiş ve c sınıfında yer almaktadır.

Ham selülozun düşük olması, yemin sindirilebilir bakış açısı olumlu kabul edilmektedir. Bu nedenle amonyum sülfatlardan oluşan kalite özelliklerine göre olumlu yönlere sahiptir. Azotlu gübre uygulamalarının bitkiselde protein sentezini arttırırken, bileşenlerin parçalarını azaltabilmesi, ham selüloz oranındaki tükenmenin temel nedenlerinden biri olabilir.

Genel olarak değerlendirildiğinde, farklı gübre uygulamalarının Macar fiğinde kalite özellikleri önemli düzeyde tutulabileceği belirlendi. DAP uygulaması dane ham proteini geliştirmeyi artırarak dane besleme değerini yükseltirken, amonyum sülfat uygulaması ham selüloz değişimini düşürerek sindirilebilirliği olumlu yönlere etki etmiştir. Buna karşılık gübresiz satılan protein büyümeleri düşük, ham selüloz gelişimleri ise yüksek sıcaklıkta, merkezde yeterli beslenmemesinin sağlanması dane sağlıklın olumsuz etkilenmektedir. Araştırma sonuçları, uygun gübreleme programlarının Macar fiğinde yalnızca getirisi değil, aynı zamanda kalite özelliklerinin de iyileştirildiğini bildiren literatürdeki gelişmelerle genel olarak uyumlu programlar. Makarnanın kalite özellikleri üzerine yapılan incelemelerde benzer sonuçların alınması.

SONUÇ

Bu araştırmada, Antalya ili Kumluca ilçesi ekolojik koşullarında yetiştirilen Macar fiği (Vicia pannonica Crantz.) Kansur çeşidinde farklı gübre uygulamalarının bitki gelişimi, verim ve yem kalite özellikleri üzerine etkileri değerlendirilmiştir. Elde edilen bulgular, gübre uygulamalarının incelenen tüm özellikler üzerinde önemli düzeyde etkili olduğunu göstermiştir. Deneme alanı topraklarının organik madde ile alınabilir fosfor ve potasyum bakımından yetersiz olması, gübre uygulamalarının bitki gelişimi ve verim üzerindeki etkilerini daha belirgin hâle getirmiştir.

Araştırmada bitki boyu bakımından en yüksek değerler triple süper fosfat (TSP) uygulamasında belirlenirken, yaş ot ve kuru ot veriminde amonyum sülfat uygulaması öne çıkmıştır. Bu sonuç, fosforun kök gelişimi ve vejetatif büyümeyi desteklediğini, azotun ise biyokütle üretimini artırdığını göstermektedir. Buna karşılık kontrol parsellerinde bitki boyu ile yaş ve kuru ot

verimlerinin en düşük düzeylerde gerçekleşmesi, uygun gübreleme yapılmaksızın ekonomik düzeyde verim elde edilmesinin güç olduğunu ortaya koymuştur.

Generatif gelişim özellikleri bakımından hayvan gübresi ve TSP uygulamaları daha başarılı sonuçlar vermiştir. Bakla ve tane sayısındaki artış, bu uygulamaların bitkinin generatif gelişimini desteklediğini göstermektedir. Özellikle hayvan gübresinin sağladığı organik madde katkısı sayesinde toprak yapısının iyileşmesi, su tutma kapasitesinin artması ve besin elementlerinin daha dengeli kullanılabilmesi, bu sonuçların temel nedenleri arasında değerlendirilmektedir.

Yem kalite özellikleri incelendiğinde, ham protein oranının en yüksek hayvan gübresi uygulamasında elde edildiği belirlenmiştir. Buna karşın ADF, NDF ve ADL değerlerinin TSP ve DAP uygulamalarında daha düşük olması, bu uygulamaların sindirilebilirliği ve yem kalitesini olumlu yönde etkilediğini göstermektedir. Kontrol uygulamasında lif fraksiyonlarının yüksek, ham protein oranının ise düşük olması, gübrelemenin yalnızca verimi değil aynı zamanda yemin besleme değerini de önemli ölçüde artırdığını ortaya koymaktadır.

Araştırma sonuçları genel olarak değerlendirildiğinde, tek bir gübre uygulamasının tüm verim ve kalite özelliklerinde üstünlük sağlamadığı belirlenmiştir. Üretim amacına bağlı olarak farklı gübre kaynaklarının tercih edilmesi daha uygun görünmektedir. Yüksek yaş ve kuru ot verimi hedeflenen işletmelerde amonyum sülfat, yüksek ham protein oranı ile generatif özelliklerin artırılmasının amaçlandığı üretim sistemlerinde hayvan gübresi, yem kalite kriterlerinin iyileştirilmesinin hedeflendiği işletmelerde ise TSP ve DAP uygulamaları ön plana çıkmaktadır.

Araştırma bulguları doğrultusunda, Macar fiği yetiştiriciliğinde gübreleme programlarının standart uygulamalar yerine toprak analiz sonuçlarına göre planlanması önerilmektedir. Organik madde bakımından yetersiz topraklarda organik gübre uygulamalarının mineral gübrelerle birlikte değerlendirilmesi, hem toprak verimliliğinin korunması hem de sürdürülebilir yem bitkisi üretiminin sağlanması açısından önem taşımaktadır. Ayrıca gelecekte yapılacak araştırmalarda organik ve mineral gübrelerin farklı doz ve kombinasyonlarının, farklı ekolojik koşullar ile farklı Macar fiği çeşitlerinde verim ve yem kalite özellikleri üzerine etkilerinin incelenmesi, daha kapsamlı gübreleme stratejilerinin geliştirilmesine katkı sağlayacaktır.

KAYNAKÇA

Açıkgöz, E. (1993). *Yem bitkileri I: Buğdaygil ve baklagil yem bitkileri tarımı*. Uludağ Üniversitesi Ziraat Fakültesi Ders Notları No: 33.

Açıkgöz, E. (2001). *Yem bitkileri* (3. bs.). Uludağ Üniversitesi Güçlendirme Vakfı Yayınları No: 182.

Akdeniz, H., Keskin, B., Yılmaz, İ., & Oral, E. (2004). Farklı azot ve fosfor dozlarının Macar fiği (*Vicia pannonica* Crantz.) verim ölçütlerine etkileri. *Yüzüncü Yıl Üniversitesi Ziraat Fakültesi Tarım Bilimleri Dergisi, 14*(1), 25–31.

Anlarsal, A. E., Çakmakçı, S., & Karadağ, Y. (1996). Çukurova kıraç koşullarında farklı gübre formlarının kışlık fiğ yetiştiriciliğinde tane verimi ve kalitesine etkileri. *Türkiye 3. Çayır-Mera ve Yem Bitkileri Kongresi Bildirileri*, 412–418.

Ayan, İ., Mut, H., Önal Aşçı, Ö., & Başaran, U. (2006). Azotlu gübrelemenin fiğ türlerinde nodülasyon ve bitki büyümesine etkileri. *Ondokuz Mayıs Üniversitesi Ziraat Fakültesi Dergisi, 21*(3), 290–295.

Bilgili, U., Uzun, A., & Sancak, C. (2009). Macar fiği (*Vicia pannonica* Crantz.) genotiplerinde fosforlu gübrelemenin tane verimi ve verim öğelerine etkileri. *Tarım Bilimleri Dergisi, 15*(3), 214–220.

Çarpıcı, E. B., Çelik, N., & Budaklı Çarpıcı, E. (2010). Farklı azot dozlarının kışlık ekilen Macar fiğinin verim ve bazı kalite özelliklerine etkileri. *Uludağ Üniversitesi Ziraat Fakültesi Dergisi, 24*(2), 45–53.

Çelen, A. E., Önal Aşçı, Ö., & Çakmakçı, S. (2013). Fosforlu gübrelemenin Macar fiğinde ot ve tohum verim bileşenlerine etkileri. *Ege Üniversitesi Ziraat Fakültesi Dergisi, 50*(1), 15–22.

Çomaklı, B., Daşcı, M., & Tan, M. (2005). Ahır gübresi ve kimyasal gübrelerin Macar fiğinin yeşil ot ve kuru madde verimine etkileri. *Atatürk Üniversitesi Ziraat Fakültesi Dergisi, 36*(2), 145–151.

Gökkuş, A., Tan, M., & Daşcı, M. (1999). Erzurum taban koşullarında kışlık ve yazlık olarak ekilen Macar fiğinin verim ve kalitesi üzerine bir araştırma. *Türkiye 3. Tarla Bitkileri Kongresi Bildirileri*, 184–189.

Gülümser, E., Mut, H., & Doğrusöz, M. C. (2020). Organik ve kompoze gübrelerin Macar fiğinde (*Vicia pannonica* Crantz.) tohum verimi ve kalitesine etkileri. *Black Sea Journal of Agriculture, 3*(2), 115–122.

Hatipoğlu, R., Tükel, T., & Çınar, S. (1999). Farklı fosfor dozlarının Macar fiği (*Vicia pannonica* Crantz.) verim ve verim unsurlarına etkileri. *Türkiye 3. Tarla Bitkileri Kongresi Bildirileri*, 224–229.

Karadağ, Y., Özköse, A., & Kır, B. (2007). Tokat koşullarında bazı Macar fiği (*Vicia pannonica* Crantz.) hat ve çeşitlerinin tane verimi performansları. *Gaziosmanpaşa Üniversitesi Ziraat Fakültesi Dergisi, 24*(1), 35–41.

Macit, M., Koç, A., & Karaoğlu, M. (2014). Organik tarımda ahır gübresi uygulamalarının Macar fiği verim bileşenleri üzerine etkileri. *Doğu Anadolu Tarım Araştırma Dergisi, 12*(1), 18–25.

Önal Aşçı, Ö., Ayan, İ., & Başaran, U. (2015). Organik ve inorganik gübrelemenin Macar fiği yetiştiriciliğinde bitki gelişimi ve yem kalitesine etkileri. *Karadeniz Tarım Bilimleri Dergisi, 28*(2), 110–117.

Sağlamtimur, T., Tansa, V., & Baytekin, H. (1990). Azotlu gübrelemenin baklagil yem bitkilerinde nodülasyon ve yeşil ot verimi ile ilişkileri. *Çukurova Üniversitesi Ziraat Fakültesi Dergisi, 5*(2), 85–94.

Serin, Y., Tan, M., & Gökkuş, A. (1999). *Kışlık yem bitkileri tarımı*. Atatürk Üniversitesi Yayınları.

Soya, H., Avcıoğlu, R., & Geren, H. (1997). Organik ve inorganik gübre kaynaklarının tek yıllık baklagil yem bitkilerinde yem kalitesi ve verimliliğe etkileri. *Ege Üniversitesi Ziraat Fakültesi Dergisi, 34*(1), 45–52.

Şenel, M. (2021). Farklı gübre kaynaklarının Macar fiği (*Vicia pannonica* Crantz.) yetiştiriciliğinde verim, verim unsurları ve nodülasyon üzerine etkileri. *Tarla Bitkileri Merkez Araştırma Enstitüsü Dergisi, 30*(1), 12–21.

Şengül, S. (2002). Farklı gübre dozlarının Macar fiği + buğday karışımlarında verim ve botanik kompozisyona etkileri. *Atatürk Üniversitesi Ziraat Fakültesi Dergisi, 33*(1), 55–61.

Tamkoç, A., Sade, B., & Soylu, S. (2011). Organik gübrelemenin yem bitkilerinde tohum verimi ve çimlenme gücü üzerine etkileri. *Selçuk Tarım ve Gıda Bilimleri Dergisi, 25*(3), 32–38.

Tan, M. (2018). *Yem bitkileri tarımı*. Atatürk Üniversitesi Yayınları.

Tosun, F. (1991). *Yem bitkileri kültürü*. Ondokuz Mayıs Üniversitesi Yayınları.

Uzun, A., Bilgili, U., Sancak, C., & Filya, İ. (2011). Macar fiği yetiştiriciliğinde gübreleme ve yabancı ot kontrolünün verim bileşenlerine etkileri. *Uludağ Üniversitesi Ziraat Fakültesi Dergisi, 25*(1), 67–76.

Yavuz, M., Gürsoy, E., & Ertekin, İ. (2018). Farklı gübreleme rejimlerinin Macar fiğinde kalite parametreleri ve sindirilebilirlik üzerine etkileri. *Süleyman Demirel Üniversitesi Ziraat Fakültesi Dergisi, 13*(2), 89–97.

ZONGULDAK EKOLOJİK KOŞULLARINDA BAZI YEM BEZELYESİ [*Pisum sativum ssp. arvense* (L.) Poir.] ÇEŞİTLERİNİN TARIMSAL ÖZELLİKLERİNİN BELİRLENMESİ

DETERMINATION OF AGRICULTURAL CHARACTERISTICS OF SOME FORAGE PEA [Pisum sativum ssp. arvense (L.) Poir.] CULTIVARS UNDER ZONGULDAK ECOLOGICAL CONDITIONS

Edanur SÖNMEZ[1]
[1]Düzce Üniversitesi, Ziraat Fakültesi, Düzce, TÜRKİYE
ORCID ID: https://orcid.org/0009-0009-8470-5262

Dr. Öğr. Üyesi Fikret BUDAK[1]
[1]Düzce Üniversitesi,Ziraat Fakültesi, Düzce, TÜRKİYE
ORCID ID: https://orcid.org/0000-0003-4355-2318

ÖZET

Bu çalışma, Zonguldak ili Merkez ilçesi Sofular köyünün ekolojik koşullarında beş farklı yem bezelyesi çeşidinin (Arda, Furkan, Assas, Atlas ve Mahir) adaptasyonu ve tarımsal performanslarını belirlemek amacıyla yapılmıştır. Çalışmada bitki boyu yaş ot, kuru ot ve dane verimleri, ham protein oranı, ADF (asit deterjanda çözünmeyen lif) NDF (Nötral deterjanda çözünmeyen lif), ADL (asit deterjanda çözünmeyen lignin) oranları incelenmiştir.

Yapılan çalışma ve gözlemler sonucunda bitki boyu bakımdan Atlas çeşidi, 123,67 cm ortalama ile en yüksek, Assas çeşidi 84,60 cm ile en düşük değere sahip olmuştur. Yaş ot verimi bakımdan yine Atlas çeşidi 2428,33 kg/da ile ilk sırada yer alırken, Mahir çeşidi 1968,33 kg/da ile en düşük yaş ot verimi vermiştir. Yaş ot verimleri 1968,33 kg/da'dan 2428,33 kg/da'ya kadar değişmiştir. Kuru ot veriminde Atlas çeşidi 490,00 kg/da verimle yine en yüksek verime sahip olmuş Furkan çeşidi 353,33 kg/da en düşük kuru ot verimine sahip olmuştur. Yapılan analiz sonucunda, kuru otta ham protein oranı %16,04-17,60, ham kül oranı %7,28-9,35 ve ham selüloz oranı %24,88-26,70 olmuştur. Sindirilebilirlik ve lif yapısını ortaya koyan parametrelerden (NDF) oranı %38,19–39,47, asit deterjanda çözünmeyen lignin (ADL) oranı %6,66-7,53 ve (ADF) oranı %29,55–30,88 olarak gözlemlenmiştir. Dane analizleri, ham protein oranlarının %26,17-26,68 ve ham selüloz oranlarının %5,20-6,05 gibi küçük ancak istatistiksel olarak önemli olduğunu göstermiştir. Çalışma sonucuna göre, Atlas çeşidi hem kaba yem verimi hem de boylanma açısından bölgedeki diğer tüm çeşitleri geride bırakmıştır.

Anahtar Kelimeler: Yem bezelyesi [*Pisum sativum ssp. arvense* (L) Poir.], yaş ot, kuru ot, dane, verim kalite

ABSTRACT

This study was conducted to determine the adaptation and agricultural performance of five different forage pea cultivars (Arda, Furkan, Assas, Atlas, and Mahir) under the ecological conditions of Sofular village in the Central district of Zonguldak province. In the study, plant height, fresh forage, dry matter, and seed yields, crude protein ratio, ADF (acid detergent fiber), NDF (neutral detergent fiber), and ADL (acid detergent lignin) ratios were investigated.

As a result of the study and observations, the Atlas cultivar had the highest value in terms of plant height with an average of 123.67 cm, while the Assas cultivar had the lowest value with 84.60 cm. In terms of fresh forage yield, the Atlas cultivar again ranked first with 2428.33

kg/da, whereas the Mahir cultivar gave the lowest fresh forage yield with 1968.33 kg/da. Fresh forage yields ranged from 1968.33 kg/da to 2428.33 kg/da. In dry matter yield, the Atlas cultivar once again had the highest yield with 490.00 kg/da, while the Furkan cultivar had the lowest dry matter yield with 353.33 kg/da. As a result of the analysis, the crude protein ratio in dry matter was 16.04–17.60%, the crude ash ratio was 7.28–9.35%, and the crude fiber ratio was 24.88–26.70%. Among the parameters revealing digestibility and fiber structure, the (NDF) ratio was observed as 38.19–39.47%, the acid detergent lignin (ADL) ratio as 6.66–7.53%, and the (ADF) ratio as 29.55–30.88%. Seed analyses showed that crude protein ratios of 26.17–26.68% and crude fiber ratios of 5.20–6.05% were small yet statistically significant. According to the study results, the Atlas cultivar outperformed all other cultivars in the region in terms of both roughage yield and plant height.

Keywords: Forage pea, [*Pisum sativum ssp. Arvense* (L) Poir.] fresh forage, dry matter, seed, yield, quality

1.GİRİŞ

Dünya genelinde ve Türkiye'de nüfus artışına paralel olarak, hayvansal kökenli proteinlere olan talep her geçen gün artmaktadır. Sağlıklı ve dengeli bir beslenmenin temel taşlarından biri olan hayvansal üretimin artırılması, doğrudan hayvancılık sektörünün sürdürülebilirliğine ve karlılığına bağlıdır. (Açıkgöz, 2001)

Hayvancılık işletmelerinde ekonomik başarının en önemli kriteri, girdi maliyetlerinin minimize edilmesidir. Modern hayvancılık işletmelerinde toplam üretim maliyetlerinin yaklaşık %60 ile %70 gibi büyük bir kısmını yem giderleri oluşturmaktadır. (Açıkgöz, 2001) Bu maliyetin büyük bir bölümü, rasyonların protein ihtiyacını karşılamak amacıyla kullanılan kesif yemlerden kaynaklanmaktadır. Kaliteli kaba yemlerin rasyondaki payının artırılması, hem hayvan sağlığının ve süt/et kalitesinin korunması hem de kesif yem kullanımının azaltılarak üretim maliyetlerinin düşürülmesi açısından zorunludur. (Sancak ve ark., 2008)

Ülkemizde uzun yıllardır devam eden kaliteli kaba yem açığının kapatılmasında ve hayvancılığın karlı bir yapıya kavuşturulmasında tek yıllık baklagil yem bitkileri üretimi stratejik bir öneme sahiptir. (Sancak ve ark., 2008)

Tek yıllık baklagil yem bitkileri, yüksek besin değerleri, sindirilebilirlik oranları ve hayvanlar tarafından iştahla tüketilmeleri nedeniyle kaba yem üretimi içerisinde özel bir yere sahiptir. (Uzun ve ark., 2005) Bu grup içerisinde yer alan yem bezelyesi (*Pisum sativum* L. ssp. *arvense*), serin iklim koşullarına mükemmel adaptasyon yeteneği ve yüksek tarımsal potansiyeli ile öne çıkan en önemli baklagil bitkilerinden biridir. (Uzun ve ark., 2005)

Yem bezelyesi, hem yeşil ot hem kuru ot hem de silaj yapımı için oldukça elverişlidir. Ayrıca, tanelerindeki yüksek protein içeriği sayesinde kesif yem karışımlarında protein kaynağı olarak da yaygın biçimde kullanılmaktadır. (Bilgili ve ark., 2010) Yem bezelyesinin kuru otundaki ham protein oranı, yetiştirilme, biçim dönemi ve çeşit özelliklerine bağlı olarak %18 ile %25 arasında değişirken; tanelerindeki ham protein oranı %20 ile %30 seviyelerine kadar ulaşabilmektedir. (Bilgili ve ark., 2010) Bu yüksek besleyici değer, özellikle süt sığırcılığı ve besi hayvancılığında rasyonların dengelenmesinde kritik bir avantaj sağlamaktadır.

Yem bezelyesinin tarımsal ekosistemlere ve sürdürülebilir tarım uygulamalarına sağladığı katkılar, sadece hayvan besleme ile sınırlı değildir.

Baklagiller familyasının genel bir özelliği olarak yem bezelyesi, köklerinde konukçu olarak barındırdığı *Rhizobium leguminosarum* bakterileri vasıtasıyla havanın serbest azotunu

toprağa bağlama (biyolojik azot fiksasyonu) yeteneğine sahiptir. (Anlarsal ve ark., 2001) Bu biyolojik süreç sayesinde tek bir vejetasyon döneminde dekara önemli miktarda saf azot kazandırabilmektedir. (Anlarsal ve ark., 2001) .

Bu durum, kendisinden sonra ekilecek olan çapa veya endüstri bitkilerinin (mısır, buğday, ayçiçeği vb.) azotlu gübre ihtiyacını önemli ölçüde azaltarak üretim maliyetlerini düşürür ve aşırı kimyasal gübre kullanımının çevreye verdiği zararları engeller. (Tan ve ark., 2012)

Aynı zamanda, yem bezelyesinin erken dönemde toprak üstü aksamının toprağa karıştırılmasıyla yapılan yeşil gübreleme uygulamaları, toprağın organik madde miktarını artırmakta, fiziksel yapısını gevşetmekte, havalanma ve su tutma kapasitesini iyileştirmektedir. (Tan ve ark., 2012) Bu yönleriyle yem bezelyesi, ekim nöbeti (münavebe) sistemlerinin verimliliğini ve sürdürülebilirliğini artıran ekolojik bir sigorta işlevi görmektedir.

Türkiye'de ve dünyada yürütülen tarımsal ıslah çalışmaları sonucunda, farklı morfolojik, fizyolojik ve tarımsal karakterlere sahip çok sayıda yem bezelyesi çeşidi tescil edilerek üretime kazandırılmıştır. (Sayar ve ark., 2011)

Bu çeşitler arasında yaprak yapısı (normal yapraklı veya yarı yapraksız), büyüme formu (dik, yarı dik veya yatık gelişen), çiçeklenme zamanı (erkenci, orta veya geççi) ve soğuğa dayanıklılık gibi agro-morfolojik özellikler yönünden büyük farklılıklar bulunmaktadır. (Sayar ve ark., 2011) Örneğin, afila (yarı yapraksız) sülük yapılı çeşitler, sülükleri vasıtasıyla birbirlerine tutunarak bitkinin yatmasını engellemekte, bu da makineli hasadı kolaylaştırarak çürüme kaynaklı verim ve kalite kayıplarını minimuma indirmektedir. (Sayar ve ark., 2011)
Buna karşın, geniş yapraklı çeşitler ise fotosentez alanlarının genişliği sayesinde belirli ekolojilerde daha yüksek yeşil ot verimi potansiyeline sahip olabilmektedir. (Aydın ve ark., 2007)

Ancak bir genotipin sahip olduğu bu genetik potansiyel, yetiştirildiği bölgenin iklim, toprak ve kültürel uygulamaları ile doğrudan etkileşim halindedir. (Aydın ve ark., 2007) Genotip x çevre etkileşimi, tarımsal üretimde çeşit seçimini belirleyen en temel unsurdur. (Başaran ve ark., 2013)

Bu nedenle, tarımsal üretimde sürdürülebilir yüksek verim ve kaliteyi yakalamanın yolu, bölge ekolojisine en iyi uyum sağlayan (adaptasyon kabiliyeti yüksek) çeşitlerin bilimsel yöntemlerle belirlenmesinden geçmektedir. (Başaran ve ark., 2013) Ülkemizin farklı tarımsal bölgelerinde mevcut tescilli çeşitlerin performanslarının yerel ölçekte test edilmesi, üreticilere doğru çeşidin önerilebilmesi ve milli sermayenin doğru yönlendirilmesi açısından bilimsel bir zorunluluktur. (Başaran ve ark., 2013).

2.BULGULAR VE TARTIŞMA

2.1 Bitki Boyu (cm)

Denemede yer alan yem bezelyesi çeşitlerinin bitki boyu değerlerine ait varyans analizi sonucunda, çeşitler arasındaki fark istatistiksel olarak ileri düzeyde önemli ($p<0.01$) bulunmuştur (Çizelge 2.1). Çeşitlerin bitki boyu ortalamaları 86,9 cm ile 123,2 cm arasında değişim göstermiştir. Yapılan LSD çoklu karşılaştırma testi sonucunda tüm çeşitler istatistiksel olarak farklı gruplarda yer almıştır. En yüksek bitki boyu ortalaması 123,2 cm ile "a" grubunda yer alan ATLAS çeşidinden elde edilirken, bunu sırasıyla ARDA (113,7 cm, b grubu) ve MAHİR (104,1 cm, c grubu) çeşitleri takip etmiştir. En düşük bitki boyu ise 86,9 cm ile "e" grubunda yer alan ASSAS çeşidinde ölçülmüştür.

Yem bezelyesinde bitki boyunun çeşitlere göre bu denli keskin farklılıklar göstermesi, genotiplerin genetik yapılarının ve büyüme habituslarının deneme ekolojisine (zonguldak şartlarına) verdiği farklı morfolojik tepkilerden kaynaklanmaktadır.

Çizelge 2.1. Yem bezelyesi çeşitlerinde bitki boyu değerlerine ait varyans analizi

Varyasyon Kaynağı	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F Değeri
Tekerrür	2	24,15	12,075	0,425
Çeşit	4	2642,84	660,71	23,2547**
Hata	8	227,3	28,4125	
Genel	14	2894,29		
Değişim Katsayısı (C.V. %)	5,18			

** $p < 0.01$ seviyesinde ileri düzeyde önemli

Denemenin en yüksek bitki boyu değerine Atlas (123.2 cm) çeşidi ulaşmış, onu yakın bir performansla Arda (113.7 cm) çeşidi takip etmiştir. Her iki çeşidin de 110 cm sınırını aşarak yüksek boy performansı göstermesi, Zonguldak'ın nemli ve bol yağışlı ilkbahar periyodunu morfolojik açıdan en iyi şekilde değerlendirdiklerini kanıtlamaktadır. Özellikle bu iki çeşidin yüksek boy potansiyeli, birim alandan elde edilecek kuru ot ve yeşil ot verimi açısından bölge üretimi için oldukça büyük bir avantaj sunmaktadır.

Denemede orta boylu grup içerisinde Mahir Bey (104.1 cm) çeşidi yer almıştır. Mahir Bey geççi bir çeşit olmasına rağmen, vejetatif gelişimini boy uzatmak yerine dallanma ve yaprak ayasını genişletme yönünde kullanmış olabileceğini göstermektedir. Bu boy seviyesiyle hem tatminkar bir ot verimi potansiyeline sahiptir hem de yatmaya karşı nispeten dengeli bir duruş sergilemektedir. En Araştırmada en düşük bitki boyu değerleri ise Furkan (92.5 cm) ve Assas (86.9 cm) çeşitlerinde kaydedilmiştir. Özellikle Assas çeşidinin 90 cm'nin altında kalarak denemenin en kısa boylu çeşidi olması, doğrudan bu genotipin yatık/yarı yatık gelişim gösteren bodur genetik yapısından kaynaklanmaktadır. Bu çeşitlerin kısa boylu olması ot verimi potansiyellerini sınırlandırsa da Karadeniz'in ani ve yoğun ilkbahar yağışlarında yatmaya karşı yüksek direnç gösterebileceklerini ve mekanize hasada (biçime) daha uygun olabileceklerini düşündürmektedir.

2.2 Bitkilerin Çıkış Süresi

Çizelge 2.2. Yem bezelyesi çeşitlerinde çıkış süresine (gün) ait varyans analizi

Varyasyon Kaynağı	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F Değeri
Tekerrür	2	0,73	0,37	104,57**
Çeşit	4	70,64	17,66	5046,00**
Hata	8	0,03	0,0035	
Genel	14	71,40		
Değişim Katsayısı(C.V. %)	0,33			

$p < 0.01$ seviyesinde ileri düzeyde önemli

Çizelge 2.2 incelendiğinde çeşitler arasında çıkış süresi bakımından tespit edilen farklılıklar $p < 0.01$ düzeyinde ileri derecede önemli bulunmuştur. Bu durum, çeşitlerin çimlenme ve toprak yüzeyine çıkış yeteneklerinin genetik olarak birbirinden tamamen farklı olduğunu kanıtlar.

En hızlı çıkış yapan çeşitler Arda (ortalama 15,3 gün) ve Atlas (ortalama 16,0 gün) olmuştur. Küçük tohum yapısına sahip veya çimlenme enerjisi yüksek olan bu erkenci çeşitler, Zonguldak'ın nispeten serin toprak şartlarında bile hızlıca çimlenerek toprak yüzeyine çıkmıştır.

Çıkış süresi en uzun olan çeşitler ise Assas (ortalama 20,1 gün) ve Mahir Bey (ortalama 20,8 gün) olarak saptanmıştır. Bu çeşitlerin tohumlarının daha kalın kabuklu veya bin dane ağırlığının daha yüksek olması, topraktan su alıp şişme ve çimlenme sürecini (imbibisyon) uzatarak çıkışı yaklaşık 5 gün geciktirmiştir. Furkan çeşidi ise 18,2 gün ile orta grupta yer almıştır.

2.3 Çiçeklenme Tarihleri

Çizelge 2.3 incelendiğinde Zonguldak ekolojik koşullarında yürütülen bu araştırmada, yem bezelyesi çeşitlerinin çiçeklenme tarihleri arasında istatistiki olarak $p < 0.01$ seviyesinde ileri düzeyde önemli farklılıklar tespit edilmiştir. Denemede en erkenci karakteri Arda (27 Nisan) ve Atlas (28 Nisan) çeşitleri gösterirken, en geççi karakteri Assas (15 Mayıs) ve Mahir Bey (16 Mayıs) çeşitleri sergilemiştir.

Çeşitler arasında saptanan bu 19 günlük geniş çiçeklenme penceresi, bölge çiftçisinin ekim nöbeti sistemine (ikinci ürün planlamasına) veya erken ilkbahar don risklerine göre en uygun genotipi seçmesine olanak tanıyacak zengin bir varyasyon sunmaktadır. Düşük C.V. (%0.81) değeri ise elde edilen bu fenolojik verilerin güvenilirliğini ve çeşitlerin çevreye adaptasyon kararlılığını açıkça ortaya koymaktadır."

Çizelge 2.3. Yem bezelyesi çeşitlerinde çiçeklenme tarihlerine ait varyans analiz

Varyasyon Kaynağı	**Serbestlik Derecesi**	**Kareler Toplamı**	**Kareler Ortalaması**	**F Değeri**
Tekerrür	2	4,80	2,40	2,25
Çeşit	4	971,07	242,77	227,59**
Hata	8	8,53	1,07	
Genel	14	984,40		
Değişim Katsayısı (C.V. %)	0,81			

$p < 0.01$seviyesinde ileri düzeyde önemli

Çizelge 2.4. Bitki boyu (cm), çeşitlerin çıkış süresi ve çiçeklenme tarihlerine ilişkin oluşan gruplar

Çeşit Adı	**Bitki Boyu (cm)**	**Çıkış Süresi (gün)**	**Çiçeklenme Tarihi**
Arda	113,70 b	15,30 d	117 c
Atlas	123,20 a	16,00 cd	118 c
Furkan	92,50 d	18,20 b	130 b

Assas	86,90 e	20,10 a	135 a
Mahir Bey	104,10 c	20,80 a	136 a
Genel Ortalama	104,08	18,08	127,2

Aynı sütunda farklı harflerle gösterilen ortalamalar arasındaki farklar istatistiki olarak önemlidir.
($p < 0.01$).

Çizelge 2.4 incelendiğinde Zonguldak ekolojik koşullarında yürütülen bu araştırmada, yem bezelyesi çeşitlerinin incelenen tüm karakterleri (bitki boyu, çıkış süresi ve çiçeklenme tarihi) arasında istatistiki olarak ileri düzeyde önemli $p < 0.01$ farklılıklar tespit edilmiştir. Çeşitlerin bu parametreler yönünden gösterdiği performanslar genel olarak iki farklı gelişim karakterini ortaya koymaktadır:

Bu grupta yer alan çeşitler, ekim sonrasında en hızlı çıkış yapan (ortalama 15-16 gün) ve ilkbaharda en erken çiçeklenen (Nisan sonu) genotipler olmuştur.

Erken ve hızlı bir vejetatif gelişim göstermeleri, bu çeşitlerin Zonguldak'ın ilkbahar nemini çok iyi değerlendirerek en yüksek bitki boyu değerlerine (113,7 - 123,2 cm) ulaşmalarını sağlamıştır. Hızlı gelişimleri ve yüksek boylanma kapasiteleriyle yüksek ot verimi ve erken hasat için en uygun çeşitler olarak öne çıkmışlardır.

Bu gruptaki çeşitlerin topraktan çıkış süreleri daha uzun sürmüş (18-21 gün) ve çiçeklenmeleri Mayıs ortasına kadar gecikmiştir. Vejetasyon süreleri (toprakta kalma süreleri) daha uzun olmasına rağmen, Assas ve Furkan çeşitleri 90 cm civarında kalarak denemenin en kısa boylu çeşitlerini oluşturmuşlardır. Bu durum, geççi çeşitlerin enerjilerini boy uzatmak yerine yatay dallanma ve yaprak aksamı oluşturmaya harcadıklarını; kısa boyları sayesinde de bölgenin yoğun yağışlarında yatmaya karşı yüksek direnç gösterebileceklerini kanıtlamaktadır.

2.4 Yaş Ot Verimi

Çizelge 2.5. Yem bezelyesi çeşitlerinin yeşil ot verimlerine ilişkin varyans analiz sonuçları.

Varyasyon Kaynağı	**Serbestlik Derecesi**	**Kareler Toplamı**	**Kareler Ortalaması**	**F Değeri**
Tekerrür	2	21450,4	10725,2	0,8122
Çeşit	4	368421	92105,125	6,9749*
Hata	8	105644	13205,513	-
Genel	14	495515	-	-
Değişim Katsayısı (C.V. %)	5,32			

**$p < 0.05$ seviyesinde önemli

Çeşitlerin Verim Performanslarının Karşılaştırılması: Gerçek çeşit ortalamaları (Çizelge 2.5) dikkate alındığında, denemenin en yüksek yeşil ot verimi 2428.33 kg/da ile Atlas çeşidinden elde edilmiştir. Atlas çeşidini sırasıyla 2188.33 kg/da ile Arda, 2110.00 kg/da ile Furkan ve 2108.33 kg/da ile Assas çeşitleri takip etmiştir. Denemenin en düşük yeşil ot verimi performansı ise 1968.33 kg/da ile Mahir çeşidinde kaydedilmiştir.

En yüksek verimli çeşit (Atlas) ile en düşük verimli çeşit (Mahir) arasında 460 kg/da gibi tarımsal açıdan oldukça büyük bir verim farkı meydana gelmiştir. Denemenin genel yeşil ot verimi ortalaması ise 2160.66 kg/da olarak hesaplanmıştır.

2.5 Kuru Ot Verimi

Çeşitlerin Kuru Ot Verim Performanslarının Karşılaştırılması: Gerçek çeşit ortalamaları (Çizelge 2.6) incelendiğinde, denemenin en yüksek kuru ot verimi 490.00 kg/da ile Atlas çeşidinden elde edilmiştir. Kuru ot verimi parametresinde Atlas çeşidini sırasıyla 423.33 kg/da ile Arda, 408.33 kg/da ile Mahir ve 385.00 kg/da ile Assas çeşitleri takip etmiştir. Denemenin en düşük kuru ot performansı ise 353.33 kg/da ile Furkan çeşidinde kaydedilmiştir.

En yüksek kuru ot verimi veren çeşit (Atlas) ile en düşük çeşit (Furkan) arasında 136.67 kg/da gibi çok belirgin bir verim farkı ortaya çıkmıştır. Deneme genelinde elde edilen kuru ot verimi ortalaması ise 412.00 kg/da olarak saptanmıştır

Çizelge 2.6. Yem bezelyesi çeşitlerinin kuru ot verimlerine ilişkin varyans analiz sonuçları.

Varyasyon Kaynağı	**Serbestlik Derecesi**	**Kareler Toplamı**	**Kareler Ortalaması**	**F Değeri**
Tekerrür	2	840,15	420,075	0,9416
Çeşit	4	31845,6	7961,4	17,8443**
Hata	8	3569,2	446,15	-
Genel	14	36255	-	-
Değişim Katsayısı (C.V. %)	5,14			

$P \leq 0{,}01$ düzeyinde önemlidir

Çizelge 2.7. Yem bezelyesi çeşitlerinin yaş ot ve kuru ot verimlerine ilişkin gruplandırmalar.

Çeşitler	**Yaş Ot Verimi (kg/da)**	**Kuru Ot Verimi (kg/da)**
Atlas	2428.33a	490.00a
Arda	2188.33b	423.33b
Furkan	2110.00bc	353.33c
Assas	2108.33bc	385.00bc
Mahir	1968.33c	408.33b
Genel Ortalama	2160.66	412.00

En yüksek yaş ot verimi 2428.33 kg/da ile Atlas çeşidinden elde edilmiş ve bu çeşit tek başına "a" üst grubunu oluşturmuştur. Atlas çeşidinin, en yakın takipçisi olan Arda çeşidine göre birim alandan daha fazla yaş ot kütlesi ürettiği istatistiksel olarak da tescillenmiştir.

Orta Verim Grubu: Arda çeşidi 2188.33 kg/da yaş ot verimi ile genel ortalamanın üzerinde bir performans göstererek tek başına "b" grubunda yer almıştır. Furkan (2110.00 kg/da) ve Assas

(2108.33 kg/da) çeşitleri ise birbirine son derece yakın değerler sergilemiş ve aralarındaki fark istatistiksel olarak önemsiz bulunarak aynı alt grupta ("bc" grubu) yer almışlardır.

Denemede en düşük yaş ot verimi performansı 1968.33 kg/da ile Mahir çeşidinde saptanmış ve bu çeşit "c" grubunu oluşturmuştur. Mahir çeşidi, lider durumdaki Atlas çeşidinden yaklaşık %19 daha az yaş ot verimi sağlamıştır.

2.6 Bakla Sayıları

Çizelge 2.8. Çeşitlerin bitki başına bakla sayılarına ait varyans analizi

Varyasyon Kaynağı	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F Değeri
Tekerrür	2	0,4333	0,2167	0,6842 ns
Çeşit	4	5,9440	1,4860	4,6926 *
Hata	8	2,5333	0,3167	
Genel	14	8,9107		
Değişim Katsayısı (CV %)	4,50			

* : %5 düzeyinde istatistiksel olarak önemli (P < 0.05), ns: Önemli değil

Çeşitlerin performansları birbiriyle kıyaslandığında, bitki başına bakla sayısı bakımından en yüksek başarıyı ASSAS çeşidi (13,1 adet/bitki) göstermiştir. ASSAS çeşidini sırasıyla FURKAN (12,8 adet/bitki) ve ATLAS (12,6 adet/bitki) çeşitleri yakından takip etmiştir. Bu üç çeşidin bakla tutma kapasitelerinin birbirine oldukça yakın ve yüksek seyretmesi, Batı Karadeniz (Zonguldak/Sofular) ekolojik şartlarında generatif organ (çiçek ve bakla) taslağı oluşturma kabiliyetlerinin benzer düzeyde başarılı olduğunu ortaya koymaktadır.

ARDA çeşidi ise 12,5 adet/bitki ile deneme ortalamasına (12,48 adet/bitki) en yakın değeri sergileyerek orta sıralarda yer almıştır. Buna karşılık, denemenin en düşük bitki başına bakla sayısı 11,4 adet/bitki ile MAHİR çeşidinde saptanmıştır. MAHİR çeşidinin bakla sayısı, lider durumdaki ASSAS çeşidinden yaklaşık %13 daha az gerçekleşmiştir.

2.7 Dane Verimleri

Çizelge 2.9. Çeşitlerin m² dane verimlerine (g/m²) ait varyans analizi tablosu

Varyasyon Kaynağı	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F Değeri
Tekerrür	2	312,4	156,2	0,452 ns
Çeşit	4	5834,0	1458,5	4,221 *
Hata	8	2764,0	345,5	
Genel	14	8910,4		
Değişim Katsayısı (CV %)	3,72			

** : %5 düzeyinde istatistiksel olarak önemli (P < 0.05), ns: Önemli değil*

Çizelge 2.10. Çeşitlerin bitki başına bakla sayıları ve m^2 dane verimlerine ilişkin Duncan gruplandırmaları

Çeşitler	Bitki Başına Bakla Sayısı (adet/bitki)	Dane Verimi (g/m^2)
Assas	13.10a	198.50a
Furkan	12.80a	184.20ab
Atlas	12.60ab	165.40bc
Arda	12.50ab	152.10cd
Mahir	11.40b	124.80a
Genel Ortalama	12.48	165.00

** Aynı sütunda benzer harfle gösterilen ortalamalar arasındaki fark istatistiksel olarak önemli değildir (P<0.05).*

Bitki başına bakla sayısı değerleri incelendiğinde, çeşitlerin 11.40 adet/bitki ile 13.10 adet/bitki arasında değişim gösterdiği görülmektedir. En yüksek bakla sayısı ASSAS (13.10 adet/bitki) çeşidinden elde edilmiş olup, bu genotipi istatistiksel olarak aynı grupta (a) yer alan FURKAN (12.80 adet/bitki) çeşidi takip etmiştir.

ATLAS (12.60 adet/bitki) ve ARDA (12.50 adet/bitki) çeşitleri ise "ab" grubunda yer alarak deneme ortalamasına (12.48 adet/bitki) yakın bir performans sergilemişlerdir. Denemede en düşük bakla sayısı ise 11.40 adet/bitki ile MAHİR çeşidinde saptanmış ve bu çeşit tek başına "b" grubunu oluşturmuştur.

Çeşitlerin m^2 dane verimi performansları değerlendirildiğinde, genotiplerin sıralamasının bakla sayısı ile büyük oranda paralellik gösterdiği dikkat çekmektedir. En yüksek dane verimi 198.50 g/m^2 ile ASSAS çeşidinde saptanmış ve "a" grubunda yer almıştır.

FURKAN çeşidi 184.20 g/m^2 dane verimi ile üst sıralardaki yerini koruyarak "ab" grubuna dahil olmuştur. Deneme genel ortalamasının (165.00 g/m^2) hemen üzerinde yer alan ATLAS (165.40 g/m^2) çeşidi "bc" grubunu oluştururken, ARDA çeşidi 152.10 g/m^2 ile alt-orta grupta ("cd") kendine yer bulmuştur. Araştırmada en düşük dane verimi performansı ise 124.80 g/m^2 ile MAHİR çeşidinden elde edilmiş ve istatistiksel olarak "d" grubunda sınıflandırılmıştır.

Sonuç olarak; lider konumda bulunan ASSAS çeşidinin dane verimi, en düşük performansı gösteren MAHİR çeşidine kıyasla yaklaşık %37.1 daha yüksek gerçekleşmiştir. Bu durum, bitki başına bakla sayısı gibi verim komponentlerinin yüksek olmasının, doğrudan birim alan dane verimine yansıdığını ve ASSAS ile FURKAN çeşitlerinin bölge ekolojisine agronometrik adaptasyon yeteneğinin diğer çeşitlere göre daha üstün olduğunu göstermektedir.

2.8 Ham Protein Oranı

Çizelge 2.11. Yem bezelyesi çeşitlerinin ham protein oranlarına ilişkin varyans analiz sonuçları.

Varyasyon Kaynağı	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F Değeri
Tekerrür	2	0,114	0,057	0,7308
Çeşit	4	4,9252	1,2313	15,7859**

Hata	8	0,624	0,078	-
Genel	14	5,6632	-	-
Değişim Katsayısı (C.V. %)	1,65			

**$P \leq 0{,}01$ düzeyinde önemlidir.

Çizelge 2.12. Yem bezelyesi çeşitlerinin ham protein oranlarına ilişkin gruplandırmalar.

Çeşit Adı	Ham Protein Oranı (%)
Arda	17,60 a
Atlas	17,31 a
Mahir	17,28 a
Furkan	16,53 b
Assas	16,04 b
Genel Ortalama	16,95

Çizelge 2.12'da sunulan varyans analizi (ANOVA) sonuçları incelendiğinde, denemeye alınan yem bezelyesi çeşitlerinin ham protein oranları arasındaki farkın istatistiksel olarak $P \leq 0.01$ önem seviyesinde ileri düzeyde önemli olduğu görülmektedir. Araştırmada hesaplanan %1,65'lik oldukça düşük Değişim Katsayısı (C.V. %), deneme hatasının minimize edildiğini ve elde edilen verilerin yüksek güvenilirlik düzeyinde olduğunu ortaya koymaktadır.

Çizelge 2.12'da yer alan Duncan çoklu karşılaştırma testi sonuçlarına göre, çeşitlerin ham protein oranları %16,04 ile %17,60 arasında değişim göstermiş ve çeşitler istatistiksel olarak iki farklı grupta toplanmıştır. Bu doğrultuda, ARDA (%17,60), ATLAS (%17,31) ve MAHİR (%17,28) çeşitleri, istatistiksel olarak aynı grupta (a) yer alarak en yüksek protein içeriğine sahip olan çeşitler olmuşlardır. Protein oranı bakımından daha düşük değerler gösteren FURKAN (%16,53) ve ASSAS (%16,04) çeşitleri ise istatistiksel olarak farklı bir grupta (b) yer almıştır.

Yem bezelyesi gibi baklagil yem bitkilerinde ham protein oranı, besleme değerini belirleyen en kritik kalite parametresidir; çeşitler arasında tespit edilen bu farklılıklar, genetik yapının besin maddesi sentezleme ve biriktirme kapasitesindeki varyasyonların bir sonucudur.

2.9 Ham Kül Oranı

Çizelge 2.13. Yem bezelyesi çeşitlerinin ham kül oranlarına ilişkin varyans analiz sonuçları.

Varyasyon Kaynağı	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F Değeri
Tekerrür	2	0,0842	0,0421	0,9152
Çeşit	4	7,1248	1,7812	38,7217**
Hata	8	0,368	0,046	-
Genel	14	7,577	-	-
Değişim Katsayısı (C.V. %)	2,66			

** $P \leq 0{,}01$ düzeyinde önemlidir.

Yem bezelyesi çeşitlerinin ham kül oranlarına ilişkin varyans analizi sonuçları Çizelge 2.13'da verilmiştir. Yapılan varyans analizi sonucunda, çeşitlerin ham kül oranları arasındaki

farklılıklar %1 (P ≤ 0,01) düzeyinde önemli bulunmuştur (Çizelge 2.13). Denemenin varyasyon katsayısı (C.V.) %2,66 olarak hesaplanmış olup, elde edilen sonuçların istatistiksel olarak yüksek güvenilirlikte olduğu belirlenmiştir.

2.10 Ham Selüloz Oranı

Çizelge 2.14. Yem bezelyesi çeşitlerinin ham selüloz oranlarına ilişkin varyans analiz sonuçları

Varyasyon Kaynağı	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F Değeri
Tekerrür	2	0,1422	0,0711	0,4257
Çeşit	4	6,3124	1,5781	9,4497**
Hata	8	1,336	0,167	-
Genel	14	7,7906	-	-
Değişim Katsayısı (C.V. %)	1,58			

$**P \leq 0{,}01$ düzeyinde önemlidir.

Yem bezelyesi çeşitlerinin ham selüloz oranlarına ait varyans analizi sonuçları Çizelge 2.14'de sunulmuştur. Gerçekleştirilen varyans analizi neticesinde, çeşitlerin ham selüloz oranları arasındaki farklılıkların %1 (P ≤ 0,01) önem düzeyinde istatistiksel olarak anlamlı olduğu tespit edilmiştir (Çizelge 2.14). Araştırmanın varyasyon katsayısı (C.V.) %1,58 olarak belirlenmiş olup, bu değer denemenin yüksek güvenilirlikte yürütüldüğünü ortaya koymaktadır.

Çizelge 4.15 Yem bezelyesi çeşitlerinin ham kül ve ham selüloz oranlarına ilişkin gruplandırmalar.

Çeşit Adı	Ham Kül Oranı (%)	Ham Selüloz Oranı (%)
Arda	9,35a	24,88c
Mahir	8,2b	26,43a
Furkan	7,87bc	26,2ab
Atlas	7,61cd	26,7a
Assas	7,28d	25,64bc
Genel Ort.	8,06	25,97

Yem bezelyesi çeşitlerinin ham kül ve ham selüloz oranlarına ilişkin ortalama değerler ile Duncan grupları Çizelge 2.15'de verilmiştir. Yapılan analizler sonucunda çeşitler arasında her iki parametre bakımından da istatistiksel olarak önemli farklılıklar saptanmıştır. Ham kül oranı en yüksek %9,35 ile Arda çeşidinden, en düşük ise %7,28 ile Assas çeşidinden elde edilmiştir. Ham selüloz oranı bakımından en yüksek değerler Atlas (%26,7) ve Mahir (%26,43) çeşitlerinde görülürken, en düşük değer %24,88 ile Arda çeşidinde belirlenmiştir. Elde edilen bulgular, çeşitlerin besin içeriği bakımından farklı adaptasyon ve kompozisyon özelliklerine sahip olduğunu ortaya koymaktadır.

2.11 ADF Oranı

Çizelge 2.16. Yem bezelyesi çeşitlerinin ADF oranlarına ilişkin varyans analiz sonuçları.

Varyasyon Kaynağı	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F Değeri
Tekerrür	2	0,214	0,107	0,5515
Çeşit	4	3,1245	0,7811	4,0263*
Hata	8	1,552	0,194	-
Genel	14	4,8905	-	-
Değişim Katsayısı (C.V. %)	1,46			

$** P \leq 0{,}05$ düzeyinde önemlidir.

Yem bezelyesi çeşitlerinin ADF oranlarına ilişkin varyans analizi sonuçları Çizelge 2.16'te sunulmuştur. Yapılan varyans analizi sonucunda, çeşitlerin ADF oranları arasındaki farklılıkların %5 ($P \leq 0{,}05$) önem düzeyinde istatistiksel olarak anlamlı olduğu belirlenmiştir (Çizelge 2.16). Araştırmanın varyasyon katsayısı (C.V.) %1,46 olarak hesaplanmış olup, elde edilen sonuçların istatistiksel olarak güvenilir olduğu görülmektedir.

2.12 NDF Oranı

Çizelge 2.17. Yem bezelyesi çeşitlerinin NDF oranlarına ilişkin varyans analiz sonuçları.

Varyasyon Kaynağı	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F Değeri
Tekerrür	2	0,341	0,1705	0,5982
Çeşit	4	2,845	0,7113	2,4958^ns
Hata	8	2,28	0,285	-
Genel	14	5,466	-	-
Değişim Katsayısı (C.V. %)	1,38			

$** P \leq 0{,}05$ düzeyinde önemlidir.

Yem bezelyesi çeşitlerinin NDF oranlarına ilişkin varyans analizi sonuçları Çizelge 2.17'te verilmiştir. Yapılan analiz sonucunda, çeşitlerin NDF oranları arasındaki farklılıklar istatistiksel olarak önemsiz ($P > 0{,}05$) bulunmuştur (Çizelge 2.17). Araştırmanın varyasyon katsayısı (C.V.) %1,38 olarak hesaplanmış olup, denemenin yüksek güvenilirlikte yürütüldüğü belirlenmiştir.

2.13 ADL Oranı

Çizelge 2.18. Yem bezelyesi çeşitlerinin ADL oranlarına ilişkin varyans analiz sonuçları.

Varyasyon Kaynağı	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F Değeri
Tekerrür	2	0,0312	0,0156	0,4875
Çeşit	4	1,4425	0,3606	11,2688**
Hata	8	0,256	0,032	-
Genel	14	1,7297	-	-

Değişim Katsayısı (C.V. %)	2,56			

$^{**}P \leq 0{,}01$ düzeyinde önemlidir.

Yem bezelyesi çeşitlerinin ADL oranlarına ilişkin varyans analizi sonuçları Çizelge 2.18'da sunulmuştur. Gerçekleştirilen varyans analizi neticesinde, çeşitlerin ADL oranları arasındaki farklılıkların %1 (P ≤ 0,01) önem düzeyinde istatistiksel olarak anlamlı olduğu tespit edilmiştir. Araştırmanın varyasyon katsayısı (C.V.) %2,56 olarak belirlenmiş olup, bu değer denemenin yüksek güvenilirlikte yürütüldüğünü ortaya koymaktadır.

Çizelge 2.19 Yem bezelyesi çeşitlerinin ADF, NDF ve ADL oranlarına ilişkin gruplandırmalar

Çeşit Adı	ADF Oranı (%)	NDF Oranı (%)	ADL Oranı (%)
Arda	30,22ab	38,94a	6,66c
Mahir	29,73bc	39,27a	7,14ab
Furkan	30,88a	38,19a	7,53a
Atlas	30,72a	38,45a	6,8bc
Assas	29,55c	39,47a	6,85bc
Genel Ort.	30,22	38,86	6,99

Yem bezelyesi çeşitlerinin ADF, NDF ve ADL oranlarına ilişkin ortalama değerler ve Duncan grupları Çizelge 2.19'da verilmiştir. Yapılan istatistiksel analizler sonucunda ADF ve ADL oranları bakımından çeşitler arasında istatistiksel olarak önemli farklılıklar (P≤0,01) saptanırken, NDF oranları bakımından çeşitler arası farkın istatistiksel olarak önemsiz olduğu belirlenmiştir. ADF oranı bakımından en yüksek değer %30,88 ile Furkan çeşidinde, en düşük değer %29,55 ile Assas çeşidinde ölçülmüştür. Lignin (ADL) içeriği açısından Furkan çeşidi en yüksek değeri verirken, Arda çeşidi en düşük değere sahip olmuştur. NDF oranları ise tüm çeşitlerde istatistiksel olarak benzer bulunmuştur.

2.14 Dane Ham Protein Oranları

Çizelge 2.20. Çeşitlerin dane ham protein oranlarına ait varyans analizi

Varyasyon Kaynağı	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F Değeri
Tekerrür	2	0,1124	0,0562	0,9590 ns
Çeşit	4	0,6953	0,1738	2,9667 ns
Hata	8	0,4688	0,0586	-
Genel	14	1,2765	-	-
Değişim Katsayısı (C.V. %)	0,92			

ns: İstatistiksel olarak önemsiz (P > 0.05)

Çizelge 2.20'da sunulan yem bezelyesi çeşitlerinin dane ham protein oranları \%26,17 ile \%26,76 arasında değişirken deneme ortalaması \%26,53 olarak tespit edilmiştir. En düşük dane

ham protein oranı \%26,17 ile ASSAS çeşidinde belirlenirken, en yüksek dane ham protein oranı ise \%26,76 ile ATLAS çeşidinden tespit edilmiştir.

2.15 Dane Ham Selüloz Oranları

Çizelge 2.21. Çeşitlerin dane ham selüloz oranlarına ait varyans analizi

Varyasyon Kaynağı	Serbestlik Derecesi	Kareler Toplamı	Kareler Ortalaması	F Değeri
Tekerrür	2	0,0645	0,0323	0,6525 ns
Çeşit	4	1,1762	0,2941	5,9414 ns
Hata	8	0,3961	0,0495	-
Genel	14	1,6368	-	-
Değişim Katsayısı (C.V. %)	3,96			

*: İstatistiksel olarak önemli (P < 0.05)

Yem bezelyesi çeşitlerinin dane ham selüloz oranları ise \%5,20 ile \%6,05 arasında değişirken deneme ortalaması \%5,60 olarak tespit edilmiştir. En düşük dane ham selüloz oranı \%5,20 ile MAHİR çeşidinde belirlenirken, en yüksek dane ham selüloz oranı ise \%6,05 ile ASSAS çeşidinden tespit edilmiştir.

Çizelge 2.22. Çeşitlerin dane ham protein ve ham selüloz oranları ile oluşan Duncan grupları

Çeşit Adı	Ham Protein (%)	Ham Selüloz (%)
Arda	26,68a	5,59b
Mahir	26,67a	5,20c
Furkan	26,36a	5,61b
Atlas	26,76a	5,56b
Assas	26,17a	6,05a
Genel Ort.	26,53	5,60

Yem bezelyesi çeşitlerinin dane ham selüloz oranlarına ait oluşan Duncan grupları incelendiğinde ise, en düşük ham selüloz oranına sahip olan MAHİR çeşidi %5,20 ile tek başına c grubunu oluşturmuştur. En yüksek ham selüloz oranına sahip olan ASSAS çeşidi %6,05 ile a grubunda yer alırken; ATLAS (%5,56), ARDA (%5,59) ve FURKAN (%5,61) çeşitleri istatistiksel olarak benzer grupta bulunarak yan yana b grubunu oluşturmuşlardır.

3.SONUÇLAR VE ÖNERİLER

2025-2026 üretim sezonunda Zonguldak (Merkez-Sofular Köyü) ekolojik koşullarında yürütülen bu araştırma kapsamında; tescilli 5 farklı yem bezelyesi çeşidinin (Atlas, Arda, Furkan, Assas, Mahir Bey) agronomik, verim ve kalite performansları kapsamlı bir şekilde incelenmiş ve elde edilen sonuçlar aşağıda özetlenmiştir:

Agronomik ve Ot Verimi Performansları: Atlas çeşidi; 123,67 cm bitki boyu, 2428,33 kg/da yeşil ot verimi ve 490,00 kg/da kuru ot verimi ile incelenen tüm agronomik ve ot verim unsurlarında ilk sırada yer alarak bölge ekolojisine mükemmel bir uyum göstermiştir. Denemede en düşük ot verimleri ise Mahir (1968,33 kg/da yeşil ot) ve Furkan (353,33 kg/da kuru ot) çeşitlerinden elde edilmiştir. Elde ettiğimiz bu ot verimi bulguları, yem bezelyesinde yeşil ot verimini 1800–2500 kg/da, kuru ot verimini ise 350–520 kg/da arasında bildiren araştırmacılarla (Sayar ve ark., 2011; Bilgili ve ark., 2010; Seydoşoğlu, 2021) büyük oranda paralellik göstermektedir. Buna karşın verimlerin 1332–3418 kg/da yeşil ot veya 250–763 kg/da kuru ot gibi daha geniş aralıklarda seyrettiği çalışmalarla da (Anlarsal ve ark., 2001; Karaköy ve ark., 2014; Taş ve ark., 2022) iklimsel farklılıklara bağlı sapmalar tespit edilmiştir. Çeşitlerin verim sıralamalarındaki bu değişimler; genotiplerin vejetatif büyüme potansiyelleri, dokularındaki su tutma kapasiteleri, yetişme dönemindeki sıcaklık-yağış dağılımı ve biçim dönemi fizyolojisinden kaynaklanmaktadır.

Generatif (Dane) Performansı: Bitki başına bakla sayısı (13,10 adet/bitki) ve birim alan dane verimi (198,50 g/m²) bakımından Assas çeşidi en yüksek performansı sergilemiş, bunu istatistiksel olarak aynı grupta yer alan Furkan çeşidi takip etmiştir. Bu durum, Assas ve Furkan çeşitlerinin generatif gelişme yönünün Batı Karadeniz iklim koşullarında ön plana çıktığını göstermektedir. Dane analizlerinde ise en düşük dane selülozu \%5,20 ile Mahir çeşidinde ölçülmüştür.

Ot Kalitesi ve Besin Değeri Parametreleri: Yem bezelyesi çeşitlerinin kalite parametreleri genelinde ortalamalar ham protein için \%16,95, ham kül için \%8,06, ham selüloz için \%25,97, ADF için \%29,55-30,88 ve ADL (lignin) için \%6,66-7,53 arasında değişim göstermiştir. NDF oranları (\%38,19-39,47) arasındaki farklılıklar ise istatistiksel olarak önemsiz saptanmıştır. Çeşitler arasında Arda çeşidi; \%17,60 ham protein ve \%9,35 ham kül oranı ile en yüksek değerleri verirken, hayvancılıkta sindirilebilirliği kısıtlayan en kritik parametre olan ADL (lignin) içeriğinde \%6,66 ve ham selülozda \%24,88 ile en düşük değerleri sunarak ot kalitesi bakımından ilk sırada yer almıştır. Elde edilen kalite bulguları literatürdeki benzer çalışmalarla (Bilgili ve ark., 2010; Sayar ve ark., 2011; Taş ve ark., 2022) uyum sergilerken; farklı biçim dönemleri ve çevre koşullarına bağlı olarak proteinin gerilediği veya selülozun arttığı araştırmalardan (Özveren ve Mut, 2018) ayrılmaktadır.

Yüksek Ot Verimi İçin: Zonguldak ve kışları serin, ilkbaharı bol yağışlı benzer ekolojilerde yüksek ot tonajı (yeşil ve kuru ot) hedefleyen üreticilere; vejetatif büyüme performansı ve adaptasyon kabiliyeti en üst düzeyde tespit edilen ATLAS çeşidinin ekilmesi öncelikli olarak tavsiye edilmelidir.

Yüksek Kaliteli Kaba Yem İçin: Yüksek verimli süt sığırcılığı gibi rasyonlarında yüksek protein ve yüksek sindirilebilirlik (düşük lignin) aranan hayvancılık işletmelerinde, ot kalitesi parametreleriyle öne çıkan ARDA çeşidinin üretimi teşvik edilmelidir.

Tohum ve Dane Üretimi İçin: Bölgede kaba yem üretimi yerine dane üretimi veya kesif yem karışımları için tohum üretimi planlanıyorsa; bakla tutma potansiyeli ve birim alan dane verimi en yüksek saptanan ASSAS çeşidi tercih edilmelidir.

Münavebe ve Yatma Riskine Karşı: Bölgenin nemli yapısından kaynaklanabilecek taban çürümelerini ve yatmadan kaynaklı makineli hasat kayıplarını önlemek adına; Atlas ve Arda gibi uzun boylu çeşitlerin yulaf veya tritikale gibi dik duruşu destekleyen hububatlar ile karışım halinde ekilmesi ya da hasadın tam çiçeklenme-bakla başlangıcı evresinde yapılması önerilir.

Gelecek Çalışmalar İçin: Bölgede öne çıkan çeşitlerin farklı ekim sıklıkları ve gübreleme dozlarındaki tepkileri incelenmeli; ayrıca silaj kalitesi parametreleri (fermantasyon özellikleri, silaj pH'ı ve organik asit kompozisyonları) de araştırma kapsamlarına dahil edilmelidir.

4.KAYNAKÇA

Açıkgöz, E. (2001). Yem Bitkileri (3. Baskı). Uludağ Üniversitesi Güçlendirme Vakfı Yayın No:182, VIPAŞ A.Ş. Yayın No: 58, Bursa, 584 s.

Anlarsal, A. E., Yücel, C., & Yılmaz, İ. (2001). Çukurova koşullarında farklı yem bezelyesi (Pisum sativum L.) hat ve çeşitlerinin verim ve verim öğelerinin saptanması. Çukurova Üniversitesi Ziraat Fakültesi Dergisi, 16(2), 33-42.

AOAC. (1990). Official Methods of Analysis (15th ed.). Association of Official Analytical Chemists, Washington, DC, USA.

Aydın, İ., Uzun, F., & Karadağ, Y. (2007). Farklı yem bezelyesi (Pisum arvense L.) hat ve çeşitlerinin Samsun ekolojik koşullarında ot verimi ve kalitelerinin belirlenmesi. Gaziosmanpaşa Üniversitesi Ziraat Fakültesi Dergisi, 24(2), 25-31.

Başaran, U., Acar, Z., Mut, H., & Aşcı, Ö. O. (2013). Karadeniz Bölgesi ekolojik koşullarında yem bezelyesi genotiplerinin adaptasyon kabiliyetleri ve verim performansları. Anadolu Tarım Bilimleri Dergisi, 28(1), 12-19.

Bilgili, U., Hanioğlu, S., & Açıkgöz, E. (2010). Yem bezelyesi (Pisum sativum L.) hat ve çeşitlerinin ot ve dane verimleri ile kalite özelliklerinin saptanması. Uludağ Üniversitesi Ziraat Fakültesi Dergisi, 24(1), 45-56.

Bilgili, U., Uzun, A., & Sancak, C. (2010). Yem Bezelyesi Çeşitlerinin Verim ve Kalite Özelliklerinin Belirlenmesi. Tarım Bilimleri Dergisi, 16(3), 167-174.

Duncan, D. B. (1955). Multiple range and multiple F tests. Biometrics, 11(1), 1-42.

Karaköy, T., Saruhan, V., & Toklu, F. (2014). Sivas Ekolojik Koşullarında Ekim Zamanlarının Bazı Yem Bezelyesi Genotiplerinde Verim ve Verim Unsurlarına Etkisi. Tarım Bilimleri Dergisi, 20(2), 112-123.

Özveren, G., & Mut, H. (2018). Yem Bezelyesi Çeşitlerinde Ot Verimi, Kalitesi ve Kuruma Davranışlarının Belirlenmesi. Karadeniz Tarım Araştırmaları Dergisi, 8(1), 45-54.

Sancak, C., Kendir, H., & Ekiz, H. (2008). Ankara koşullarında yetiştirilen bazı yem bezelyesi (Pisum sativum L.) çeşitlerinin verim ve kalite özelliklerinin belirlenmesi. Tarım Bilimleri Dergisi, 14(3), 215-222.

Sayar, M. S., Anlarsal, A. E., & Han, İ. (2011). Güneydoğu Anadolu Bölgesi Koşullarında Bazı Yem Bezelyesi (Pisum sativum L. var. arvense) Hat ve Çeşitlerinin Ot Verimi ve Kalitesi Yönünden Değerlendirilmesi. Tarım Bilimleri Dergisi, 17(3), 211-221.

Sayar, M. S., Han, Y., Yolcu, H., & Yücel, H. (2011). Güneydoğu Anadolu Bölgesi koşullarında bazı yem bezelyesi (Pisum sativum L.) hat ve çeşitlerinin ot verimi ve adaptasyon yetenekleri. TÜRKTOB Dergisi, 1(2), 38-44.

Seydoşoğlu, S. (2021). Batman Ekolojik Koşullarında Bazı Yem Bezelyesi (Pisum sativum L. var. arvense) Çeşitlerinin Kaba Yem Verimi ve Kalite Özelliklerinin Belirlenmesi. Dicle Üniversitesi Veteriner Fakültesi Dergisi, 14(1), 32-38.

Steel, R. G. D., & Torrie, J. H. (1980). Principles and Procedures of Statistics: A Biometrical Approach (2nd ed.). McGraw-Hill, New York.

Tan, M., Temel, S., & Şahin, M. (2012). Erzurum şartlarında yem bezelyesi çeşitlerinin ot ve dane verimleri ile toprağa sağladıkları organik madde ve azot katkıları. Atatürk Üniversitesi Ziraat Fakültesi Dergisi, 43(1), 15-22.

Taş, N., Erkovan, H. İ., & Kuşvuran, A. (2022). Yem Bezelyesi Çeşitlerinin Farklı Biçim Zamanlarında Verim ve Nispi Yem Değerlerinin Belirlenmesi. Tarla Bitkileri Merkez Araştırma Enstitüsü Dergisi, 31(1), 78-89.

Uzun, A., Gül, İ., & Ertekin, İ. (2005). Serin iklim yem bitkilerinden yem bezelyesinde (Pisum arvense L.) farklı biçim dönemlerinin ot verimi ve besin değerine etkisi. Türkiye VI. Tarla Bitkileri Kongresi, 5-9 Eylül, Antalya, Cilt 2, s. 781-786.

Van Soest, P. J., Robertson, J. B., & Lewis, B. A. (1991). Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. Journal of Dairy Science, 74(10), 3583-3597.

BALIKESİR EKOLOJİK KOŞULLARINDA BAZI İTALYAN ÇİMİ (*LOLIUM MULTIFLORUM LAM.*) ÇEŞİTLERİNİN VERİM VE KALİTE ÖZELLİKLERİNİN İNCELENMESİ

INVESTIGATION OF THE YIELD AND QUALITY CHARACTERISTICS OF SOME ITALIAN RYEGRASS (*LOLIUM MULTIFLORUM LAM.*) CULTIVARS UNDER BALIKESİR ECOLOGICAL CONDITIONS

Hilal Özge DEMİR
Düzce Üniversitesi
ORCID ID: : https://orcid.org/0009-0009-8718-6197

Dr. Öğretim Üyesi Fikret BUDAK
ORCID ID: https://orcid.org/0000-0003-4355-2318

ÖZET

Bu araştırma, Balıkesir ekolojik koşullarında bazı İtalyan çimi (*Lolium multiflorum Lam.*) çeşitlerinin verim ve kalite özelliklerini belirlemek amacıyla yürütülmüştür. Deneme, 2025 yılında tesadüf blokları deneme desenine göre üç tekerrürlü olarak kurulmuştur. Araştırmada bitki çıkış süresi, çiçeklenme gün sayısı, bitki boyu, yeşil ot verimi, kuru ot verimi ve ham protein verimi belirlenmiştir. Ayrıca kuru otta ham kül, ham protein, ham selüloz, Asit Deterjan Lifi (ADF) ve Nötr Deterjan Lifi (NDF) oranları ile tohumda ham protein ve ham selüloz oranları incelenmiştir. Elde edilen bulgular, gözlemlenen ve analiz edilen özellikler bakımından çeşitler arasında istatistiksel olarak önemli farklılıklar bulunduğunu göstermiştir.

Anahtar Sözcükler: İtalyan çimi, *Lolium multiflorum*, çeşit, verim, kalite

ABSTRACT

This study was conducted to determine the yield and quality characteristics of selected Italian ryegrass (*Lolium multiflorum Lam.*) cultivars under the ecological conditions of Balıkesir. The field experiment was established in 2025 in a randomized complete block design with three replications. The traits evaluated were days to emergence, days to flowering, plant height, green forage yield, dry matter yield, and crude protein yield. In addition, crude ash, crude protein, crude cellulose, Acid Detergent Fiber (ADF), and Neutral Detergent Fiber (NDF) contents of the dry forage, together with crude protein and crude cellulose contents of the seed, were analyzed. The results revealed statistically significant differences among the cultivars for the observed and analyzed characteristics.

Keywords: Italian ryegrass, *Lolium multiflorum*, cultivar, yield, quality

STUDY OF THE HYDROLYSIS OF SODIUM BOROHYDRIDE IN THE PRESENCE OF A GRAPHITIC CARBON NITRIDE-SUPPORTED BIMETALLIC CATALYST

Assoc. Prof. Dr. Erhan ONAT[1]
[1]OrganizedIndustrial Zone Vocational School, Department of Electrical and Energy, Bitlis Eren University, Bitlis, Turkey
[1]ORCID ID: https://orcid.org/0000-0003-1638-0151

Gökay ÜSTYOL[2]
[2]OrganizedIndustrial Zone Vocational School, Department of Electrical and Energy, Bitlis Eren University, Bitlis, Turkey

Prof. Dr. Mehmet Sait İZGİ[3]
[3]Department of Chemical Engineering, Faculty of Engineering, Siirt University, Siirt, Turkey;
[3]Faculty of Health Sciences, Bitlis Eren University, Bitlis, Turkey
[3]ORCID ID: https://orcid.org/0000-0003-3685-3219

ABSTRACT

Hydrogen is considered one of the most promising alternatives for overcoming both the supply limitations associated with fossil fuels and the environmental problems arising from their utilization. Accordingly, research on the use of hydrogen in energy-related applications has been rapidly advancing. Sodium borohydride ($NaBH_4$) is one of the most important compounds used for solid-state hydrogen storage, as it offers a potential solution to the challenges associated with hydrogen transportation and storage.

In this study, the development of a novel bimetallic supported catalyst for hydrogen production through the hydrolysis of sodium borohydride ($NaBH_4$) was aimed. A two-dimensional material, graphitic carbon nitride (g-C_3N_4), was employed as the catalyst support. Within the scope of the study, g-C_3N_4 (GCN) was synthesized from urea by thermal treatment in a muffle furnace at 550 °C, in accordance with procedures reported in the literature. Subsequently, cobalt (Co) and copper (Cu) metal salts were decorated onto the GCN support using the impregnation method. The synthesized supported catalysts were characterized using FT-IR, ICP, SEM, and EDX analyses to elucidate their structural and morphological properties. The structurally characterized catalysts were then evaluated for hydrogen generation via the hydrolysis of sodium borohydride. The effects of NaOH concentration, catalyst amount, and $NaBH_4$ concentration on the hydrolysis performance were systematically investigated. The optimum conditions were determined as an 8 wt.% total metal loading, 7.5 wt.% NaOH concentration, 150 mg of catalyst, and 3 wt.% $NaBH_4$ concentration. Under these optimum conditions, the hydrogen generation rate was determined to be 4443 mL min^{-1} g^{-1}.

Keywords: Graphitic carbon nitride, hydrogen production, catalyst, sodium borohydride

INTRODUCTION

The increasing environmental pollution resulting from the excessive consumption of fossil fuels since the onset of globalization has encouraged the development and utilization of renewable energy sources such as wind, wave, solar, and hydrogen energy. Among these renewable energy sources, hydrogen has attracted considerable attention from researchers due to its

environmentally friendly nature and the fact that water vapor (H_2O) is its primary oxidation product (Arutyunov and Lisichkin, 2017; Gadgil et al., 2022). However, challenges associated with the storage and transportation of hydrogen have led to increased interest in metal hydrides, such as $NaBH_4$, KBH_4, and $LiBH_4$, as chemical hydrogen storage materials. Among these materials, sodium borohydride ($NaBH_4$) has emerged as a promising candidate for chemical hydrogen storage because of its high hydrogen content of approximately 10.8 wt.%, environmental compatibility, potential recyclability, and high stability in alkaline solutions. Previous studies on hydrogen generation from $NaBH_4$ have demonstrated that, in addition to providing a clean energy source, its hydrolysis enables controllable hydrogen production kinetics in the presence of suitable alkaline solutions (Kirk et al., 2023; Gülbahça, 2024). The hydrolysis of sodium borohydride in the presence of a catalyst produces four moles of hydrogen per mole of $NaBH_4$ according to the following reaction:

$$NaBH_4 + 2H_2O \rightarrow NaBO_2 + 4H_2 \uparrow$$

Importantly, two moles of the generated hydrogen originate from water, highlighting the significance of this reaction for efficient hydrogen production. Therefore, the catalyst employed in this process is expected to exhibit high catalytic activity and excellent stability. Noble metals, such as Ru, Rh, and Pt, are commonly used as catalysts for $NaBH_4$ hydrolysis; however, their high cost and limited availability restrict their widespread application (Al-Shaikh et al., 2022; Farrag and Ali, 2022). Consequently, considerable attention has been directed toward the development of inexpensive, stable, and highly active non-noble metal catalysts based on Ni, Co, Cr, Cu, and Fe. Among these transition metals, Co-based catalysts have received particular attention because of their relatively low cost, facile synthesis, and high catalytic activity (Bulut et al., 2016; Karami and Fathirad, 2023; Sun et al., 2023; Xu et al., 2024; Demirci and Miele, 2010). However, the use of metal nanoparticles alone may lead to agglomeration, resulting in a reduction in the number of accessible active sites and, consequently, decreased catalytic performance. To overcome this limitation, metal nanoparticles are commonly immobilized on suitable support materials to improve their dispersion and catalytic activity. Various support materials, including nickel foam, graphene, activated carbon, carbon nanotubes (CNTs), and metal–organic frameworks (MOFs), have been employed to suppress nanoparticle aggregation and enhance catalytic performance (Alshammari et al., 2023; Cong et al., 2023).

In the present study, a novel supported bimetallic catalyst was developed for hydrogen generation through the catalytic hydrolysis of sodium borohydride ($NaBH_4$). Graphitic carbon nitride (g-C_3N_4, GCN), an emerging two-dimensional material, was employed as the catalyst support. GCN was synthesized from urea through thermal treatment at 550 °C in a furnace, following procedures reported in the literature. After the synthesis of GCN, with an approximate yield of 4%, cobalt (Co) and copper (Cu) metal salts were deposited onto the GCN support using the impregnation method. The bimetallic catalyst was prepared using a 1:1 Co/Cu molar ratio, selected based on previously reported optimum compositions.

Following catalyst synthesis, comprehensive characterization studies were conducted to elucidate the structural and morphological properties of the prepared material. FT-IR, SEM, and EDX analyses were performed to investigate the interaction of the metal species with the GCN support, as well as the surface morphology and elemental distribution of the catalyst. The structurally characterized supported catalyst was subsequently evaluated for hydrogen generation via the hydrolysis of sodium borohydride. The effects of several reaction parameters, including the reaction medium, $NaBH_4$ concentration, and catalyst amount, were systematically investigated to determine the optimum conditions for hydrogen production.

METHODOLOGY

In this study, graphitic carbon nitride (g-C_3N_4) was synthesized from urea in accordance with procedures reported in the literature. The precursor was thermally treated in a muffle furnace at 550 °C using a heating rate of 10 °C min^{-1}. Following the synthesis of g-C_3N_4, which was obtained with an approximate yield of 4%, cobalt (Co) and copper (Cu) metal salts were deposited onto the graphitic carbon nitride (GCN) support using the impregnation method. The bimetallic catalyst was synthesized using a 1:1 Co/Cu molar ratio, based on the optimum ratio reported in the literature. After the dropwise addition of the metal precursor solution over approximately 2–3 h, the resulting composite mixture was stirred at 750 rpm for 24 h under ambient conditions. Subsequently, 20 mL of a 10 M sodium borohydride ($NaBH_4$) solution was added dropwise to the mixture to reduce the Co and Cu metal ions present on the surface of the support material. Following the reduction process, the resulting supported catalyst was collected by filtration, thoroughly washed with deionized water, and dried in an oven at 80 °C for 12 h under a nitrogen atmosphere. The dried catalyst was then used in the subsequent hydrolysis experiments.



Figure 1. Experimental setup used for the hydrolysis experiments.

The hydrolysis experimental setup consisted of a temperature-controlled circulating water bath, a jacketed reaction tube, a magnetic stirrer, a graduated burette, and connecting tubes. The jacketed reaction tube provided the reaction environment, while the magnetic stirrer and circulating water bath ensured that each hydrolysis experiment was conducted under controlled and predetermined reaction conditions. The volume of hydrogen gas generated during the reaction was measured using a graduated burette. Hydrogen production as a function of time was determined using the water displacement method.

Catalyst Characterization and Hydrolysis Experiments

Following the synthesis of the supported catalyst, several characterization techniques were employed to elucidate its structural, compositional, and morphological properties. Fourier-transform infrared spectroscopy (FT-IR), inductively coupled plasma (ICP) analysis, scanning electron microscopy (SEM), and energy-dispersive X-ray spectroscopy (EDX) were performed to investigate the interaction and incorporation of the Co and Cu species with the g-C_3N_4 support, as well as the surface morphology and elemental composition of the synthesized catalyst.

The structurally characterized supported catalyst was subsequently evaluated for hydrogen generation through the hydrolysis of sodium borohydride ($NaBH_4$). The effects of several reaction parameters, including the reaction medium, $NaBH_4$ concentration, and catalyst amount, were systematically investigated to determine the optimum conditions for hydrogen production. As a result, a supported bimetallic catalyst containing 92 wt.% g-C_3N_4 and 8 wt.% total metal content was successfully developed and evaluated for catalytic hydrogen generation.

CONCLUSION AND DISCUSSION

Caracterization

The total metal loading of the synthesized material was determined to be 8 wt.% by ICP-OES analysis**.** This value was consistent with the EDX results and was therefore used in the catalyst efficiency calculations based on the amount of H_2 gas generated in the present study. The morphological properties and elemental composition of the catalyst were analyzed using scanning electron microscopy coupled with energy-dispersive X-ray spectroscopy (SEM-EDS)**,** and the corresponding SEM images are presented in Figure 1. Examination of the SEM images obtained at two different magnifications revealed that the Co–Cu species were homogeneously dispersed over the g-C_3N_4 support**.** In addition, the images show a highly rough and irregular surface morphology, indicating a potentially high specific surface area and the presence of a large number of accessible active sites. As observed in the SEM analysis, the g-C_3N_4 support effectively inhibited the agglomeration of the metal species and thereby contributed positively to the catalytic activity of the synthesized catalyst.



a magnification of 10,000 a magnification of 20,000

Figure 2. SEM image of the synthesized supported catalyst.

Another analysis performed within the scope of this study was energy-dispersive X-ray spectroscopy (EDS)**.** The EDS elemental mapping results presented in Figure 3, clearly demonstrate the homogeneous distribution of the principal elements, including Co, Cu, C, N, and O, throughout the catalyst structure. The presence of oxygen-containing functional groups may be associated with carbonaceous oxygen species and/or N–Cu–O and N–Co–O

configurations. Since the distribution patterns of Co, Cu, and N are highly similar, the close interaction between the Co and Cu atoms anchored on the g-C_3N_4 support is expected to play an important role in the catalytic activity. This effect is also supported by the hydrolysis results obtained in this study. Overall, the EDS analysis confirms that the Co–Cu bimetallic species were successfully anchored and homogeneously dispersed on the g-C_3N_4 support**,** indicating the successful formation of the Co–Cu/g-C_3N_4 catalyst.



Figure 3. EDX analysis of the GCN-supported Co–Cu catalyst.

Another important characterization technique, Fourier-transform infrared spectroscopy (FT-IR)**,** was employed to confirm the successful synthesis of GCN and to investigate the incorporation of Co and Cu species onto the GCN support. As shown in Figure 4**,** the FT-IR spectrum confirms the successful formation of GCN in accordance with the literature. In addition, consistent with the EDX results, characteristic bands associated with the Co–Cu species can be observed around 1500 cm^{-1}**,** although their intensities are relatively weak. The weak intensity of these bands can be attributed to the high GCN content (92 wt.%**)** in the catalyst, which may mask the characteristic signals of the relatively low metal loading (Yan et al., 2009).



Figure 4. FT-IR analysis of the g-C_3N_4@Co–Cu catalyst.

Hydrolysis Experiments

Since sodium borohydride can undergo spontaneous hydrolysis, hydrolysis experiments were conducted in the presence of the catalyst by adding NaOH at different concentrations ranging from 0 to 10 wt.%. The corresponding experimental results are presented in Figure 5. As shown in the figure, hydrogen production increased with increasing NaOH concentration and reached a maximum at 7.5 wt.% NaOH. Therefore, a NaOH concentration of 7.5 wt.% was selected for the subsequent experiments. The decrease in hydrogen production observed at 10 wt.% NaOH may be attributed to the combined effect of the excessively high NaOH concentration and the deactivation of the catalyst surface by the metaborate species formed during the reaction, as illustrated in Equation 1 (Gülbahça, 2024).

Following the determination of the optimum reaction medium, hydrolysis experiments were performed using different amounts of catalyst. The corresponding results are presented in Figure 6b. As shown in the figure, the overall reaction rate decreased with increasing catalyst amount. However, the more relevant parameter for evaluating catalyst performance is the hydrogen generation rate normalized to the catalyst mass. The hydrogen generation rates obtained using 75, 100, 150, and 200 mg of catalyst were 2876, 3677, 4206, and 4102 mL g^{-1} min^{-1}, respectively. Among the investigated catalyst amounts, the highest hydrogen generation rate was achieved using 150 mg of catalyst. This result indicates that 150 mg provides the most favorable balance between the substrate and catalyst, resulting in the most efficient catalyst–substrate interaction under the investigated reaction conditions.



Figure 5. g-C_3N_4@Co–Cu catalyst-assisted SBH hydrolysis: (a) effect of NaOH concentration and (b) effect of catalyst amount.

Following the optimization of the reaction medium and catalyst amount, the effect of $NaBH_4$ concentration and reaction temperature on the hydrolysis performance was investigated. For the $NaBH_4$ concentration experiments, concentrations ranging from 1 to 4 wt.% were examined, and the corresponding hydrolysis results are presented in Figure 6a. As shown in the figure, the amount of hydrogen produced increased with increasing $NaBH_4$ concentration. This trend can be attributed to the higher availability of $NaBH_4$ molecules in the reaction medium, which provides more reactant molecules for the hydrolysis reaction and consequently enhances hydrogen generation.



Figure 6a. Hydrolysis of SBH using the g-C_3N_4@Co–Cu catalyst at different concentrations.

Figure 6-b shows the hydrogen generation rates obtained at different $NaBH_4$ concentrations. As shown in the figure, the highest hydrogen generation rate was achieved at a $NaBH_4$ concentration of 3 wt.%, reaching 4443 mL g^{-1} min^{-1}**.** This concentration was therefore identified as the optimum $NaBH_4$ concentration under the investigated reaction conditions.



Figure 6b. Hydrogen generation rate of the g-C_3N_4@Co–Cu catalyst at different SBH concentrations.

CONCLUSION

In this study, a comprehensive investigation was conducted on hydrogen generation through the hydrolysis of sodium borohydride ($NaBH_4$, SBH) using a Co–Cu@g-C_3N_4 catalyst. The g-C_3N_4 support was synthesized from urea at 550 °C, followed by the immobilization of Co and Cu species onto the support via the impregnation method. The resulting Co–Cu@g-C_3N_4 catalyst exhibited a total metal loading of 8 wt.%**.** The catalytic performance of the synthesized catalyst for SBH hydrolysis was systematically investigated by varying the NaOH concentration, catalyst amount, and SBH concentration. The optimum reaction conditions were determined to be 7.5 wt.% NaOH, 150 mg catalyst, and 3 wt.% SBH, respectively. Under these conditions, the maximum hydrogen generation rate was determined to be 4443 mL g^{-1} min^{-1}**.**

Furthermore, the synthesized catalyst was characterized using ICP-OES, SEM-EDS, and FT-IR techniques to elucidate its elemental composition, surface morphology, and structural properties. The characterization results confirmed the successful incorporation and dispersion of the Co–Cu species on the g-C_3N_4 support. Overall, the obtained results demonstrate that the Co–Cu@g-C_3N_4 catalyst is a promising candidate for efficient hydrogen generation from $NaBH_4$ hydrolysis**.**

REFERENCES

Aksoy, M., Korkut, S. E., & Metin, Ö. (2023). The rational design of gCN/a-WOx/Pt heterostructured nanophotocatalysts for boosting the hydrogen generation from the hydrolysis of ammonia borane under visible light. International Journal of Hydrogen Energy, 48(60), 22921-22933.

Al-shaikh, H., Lasri, J., Knight, J. G., & Al-Goul, S. T. (2022). Palladium mesoporous nanoparticles Pd NPs@[KIT-6] and Pd NPs@[KIT-6]-PEG-imid as efficient heterogeneous catalysts for H2 production from NaBH4 hydrolysis. Fuel, 325, 124962.

Alshammari, K., Alotaibi, T., Alshammari, M., Alhassan, S., Alshammari, A. H., & Taha, T. A. M. (2023). Synthesis of sulfur@ g-C3N4 and CuS@ g-C3N4 catalysts for hydrogen production from sodium borohydride. Materials, 16(12), 4218.

Arutyunov, V. S., & Lisichkin, G. V. (2017). Energy resources of the 21st century: problems and forecasts. Can renewable energy sources replace fossil fuels. Russian Chemical Reviews, 86(8), 777.

Bulut, A., Yurderi, M., Ertas, İ. E., Celebi, M., Kaya, M., & Zahmakiran, M. (2016). Carbon dispersed copper-cobalt alloy nanoparticles: A cost-effective heterogeneous catalyst with exceptional performance in the hydrolytic dehydrogenation of ammonia-borane. Applied Catalysis B: Environmental, 180, 121-129.

Cong, H., Ren, J., Zhang, D., Xu, F., Wang, X., Wang, Y., ... & Wu, S. (2023). Catalytic hydrolysis of sodium borohydride for hydrogen generation using g-C3N4/Co–W–B/Ni foam composite catalyst. Journal of Materials Science, 58(2), 787-801.

Demirci, U. B., & Miele, P. (2010). Cobalt in NaBH 4 hydrolysis. Physical Chemistry Chemical Physics, 12(44), 14651-14665.

Farrag, M., & Ali, G. A. (2022). Hydrogen generation of single alloy Pd/Pt quantum dots over Co3O4 nanoparticles via the hydrolysis of sodium borohydride at room temperature. Scientific Reports, 12(1), 17040.

Gadgil, A., Tomich, T. P., Agrawal, A., Allouche, J., Azevedo, I. M., Bakarr, M. I., ... & Wang, Y. (2022). The great intergenerational robbery: A call for concerted action against environmental crises. Annual Review of Environment and Resources, 47(1), 1-4.

Gülbahça, F. (2024) Sodyum borhidrür hidrolizi için yeni nesil nanokatalizörlerin sentezi, karakterizasyonu ve biyokimyasal uygulamaları, Kimya Ana Bilim Dalı, Fen Bilimleri Enstitüsü, Kütahya Dumlupınar Üniversitesi

Karami, M., & Fathirad, F. (2023). Cobalt ferrite nanoparticles anchored on reduced graphene oxide nanoribbons (0D/1D CoFe2O4/rGONRs) as an efficient catalyst for hydrogen generation via NaBH4 hydrolysis. Inorganic Chemistry Communications, 150, 110552.

Kirk, J., Kim, Y., Lee, Y. J., Kim, M., Min, D. S., Kim, P. S., ... & Jeong, H. (2023). Pushing the limits of sodium borohydride hydrolysis for on-board hydrogen generation systems. Chemical Engineering Journal, 466, 143233.

Sun, L., Gao, X., Ning, X., Qiu, Z., Xing, L., Yang, H., ... & Meng, Y. (2023). Cobalt-nickel bimetal carbon sphere catalysts for efficient hydrolysis of sodium borohydride: The role of synergy and confine effect. International Journal of Hydrogen Energy, 48(9), 3413-3428.

Xu, F., Ren, J., Ma, J., Wang, Y., Zhang, K., Cao, Z., ... & Bai, S. (2024). A review of hydrogen production kinetics from the hydrolysis of NaBH4 solution catalyzed by Co-based catalysts. International Journal of Hydrogen Energy, 50, 827-844.

Yan, S. C., Li, Z. S., & Zou, Z. G. (2009). Photodegradation performance of g-C3N4 fabricated by directly heating melamine. Langmuir, 25(17), 10397-10401.

STUDY OF THE CATALYTIC HYDROLYSIS KINETICS OF SODIUM BOROHYDRIDE IN THE PRESENCE OF A SUPPORTED BIMETALLIC CATALYST

Gökay ÜSTYOL[1]
[1]OrganizedIndustrial Zone Vocational School, Department of Electrical and Energy, Bitlis Eren University, Bitlis, Turkey

Prof. Dr. Mehmet Sait İZGİ[2]
[2]Department of Chemical Engineering, Faculty of Engineering, Siirt University, Siirt, Turkey;
[2]Faculty of Health Sciences, Bitlis Eren University, Bitlis, Turkey
[2]ORCID ID: https://orcid.org/0000-0003-3685-3219

Assoc. Prof. Dr. Erhan ONAT[3]
[3]OrganizedIndustrial Zone Vocational School, Department of Electrical and Energy, Bitlis Eren University, Bitlis, Turkey
[3]ORCID ID: https://orcid.org/0000-0003-1638-0151

Abstract

In this study, the kinetic behavior of hydrogen production through the hydrolysis of sodium borohydride ($NaBH_4$) was investigated in the presence of a Co-Cu@g-C_3N_4 catalyst prepared by impregnating Co and Cu metals onto graphitic carbon nitride (g-C_3N_4) synthesized from urea. Catalytic dehydrogenation reactions were conducted at different temperatures of 25, 30, 35, and 40 °C to determine the reaction order and activation parameters. The kinetic data obtained at different temperatures were evaluated according to zero-order, first-order, and general *n*th-order kinetic models. The results demonstrated that the hydrolysis of $NaBH_4$ in the presence of the Co-Cu@g-C_3N_4 catalyst was best described by a zero-order kinetic model, with a high degree of correlation. The rate constants obtained from the zero-order kinetic plots at different temperatures were subsequently evaluated using the Arrhenius equation, and the activation energy for $NaBH_4$ hydrolysis over the Co-Cu@g-C_3N_4 catalyst was determined to be 38.18 kJ mol^{-1}.

In addition, successive recycling experiments were conducted to evaluate the catalytic stability and reusability of the catalyst. The results revealed that the catalyst retained its catalytic activity even after the fifth reaction cycle, achieving 100% conversion. These findings demonstrate that the synthesized Co-Cu@g-C_3N_4 catalyst is a promising catalytic system for hydrogen production via $NaBH_4$ hydrolysis, exhibiting high catalytic activity, a relatively low activation energy, and good reusability.

Keywords: Activation energy, graphitic carbon nitride, hydrogen production, Co-Cu@g-C_3N_4, sodium borohydride.

INTRODUCTION

Chemical hydrides have attracted considerable attention as potential hydrogen storage materials because of their high hydrogen storage capacities and several advantageous properties. Among

them, potassium borohydride (KBH_4) and sodium borohydride ($NaBH_4$) are among the most widely investigated borohydrides due to their ability to readily generate hydrogen through methanolysis and hydrolysis reactions under relatively mild conditions, as well as their potential practical applications in fuel cell systems. However, sodium borohydride ($NaBH_4$) is generally preferred because of its relatively high hydrogen content (10.8 wt.%). Since $NaBH_4$ can undergo spontaneous hydrolysis at room temperature, alkaline conditions are generally required to improve the stability of the $NaBH_4$ solution. Therefore, NaOH is commonly added to the reaction medium to maintain a sufficiently high pH, suppress uncontrolled hydrolysis, and enable the effective evaluation of catalytic activity (Arutyunov and Lisichkin, 2017; Alshammari et al., 2023).

Previous studies have demonstrated that both homogeneous and heterogeneous catalysts can significantly accelerate the hydrolysis and methanolysis of $NaBH_4$. Noble metals such as Pd, Ru, and Pt generally exhibit high catalytic activity and stability toward $NaBH_4$ hydrolysis. However, their limited availability and high cost restrict their widespread practical application. Consequently, the development of alternative catalysts based on inexpensive transition metals has become highly desirable. Among these metals, cobalt (Co) has been considered a promising catalyst for promoting $NaBH_4$ hydrolysis because it can exhibit catalytic reactivity comparable to that of noble metals while offering a substantially lower cost. Furthermore, numerous studies have investigated the incorporation of secondary metals, such as Cu, Fe, Cr, and Mo, into Co-based catalysts to generate synergistic effects and enhance catalytic activity (Karami et al., 2023; Ekinci and Onat, 2024; Onat et al., 2026).

Various support materials have been employed to prevent the agglomeration of synthesized metal nanoparticles and improve catalyst stability. In this context, suitable structural supports or templates capable of promoting the uniform dispersion of cobalt-containing nanoparticles are particularly important. Metal–organic frameworks (MOFs), which constitute a stable family of highly modular functional materials assembled from metal ions and organic linkers, can be used as precursors or templates for the preparation of porous oxides, TiO_2, and metal oxide/carbon composites. In addition, carbon-based materials have received considerable attention as catalyst supports because of their large surface areas, favorable structural properties, and ability to promote the dispersion of active metal species.

Graphitic carbon nitride (g-C_3N_4) has emerged as a particularly attractive support material due to its high nitrogen content, chemical stability, and ability to interact strongly with metal nanoparticles. Nevertheless, only a limited number of studies have reported the application of ultrafine Co–Cu nanocrystals confined or supported on g-C_3N_4-derived materials as catalysts for $NaBH_4$ hydrolysis. Moreover, the durability and reusability of such catalytic systems have not yet been comprehensively investigated or systematically discussed in the existing literature. Various experimental approaches have been developed to determine the hydrogen generation rate during the catalytic hydrolysis of $NaBH_4$. In recent years, particular attention has been directed toward understanding the kinetic behavior of metal-based catalysts in $NaBH_4$ hydrolysis. The reaction rate can be influenced by several parameters, including temperature, pH, catalyst composition and loading, the formation of the $NaBO_2$ by-product and its associated hydration, and water availability. Among these parameters, reaction temperature has a substantial influence on both the hydrogen generation rate and the overall kinetics of the hydrolysis process. An increase in reaction temperature generally results in a higher hydrogen generation rate and can promote the nearly complete chemical conversion of $NaBH_4$. However, because $NaBH_4$ hydrolysis is an exothermic reaction, heat accumulation during the reaction may result in an uncontrolled increase in temperature. Such temperature fluctuations can lead

to unpredictable reaction behavior and an unstable hydrogen generation rate (Cong et al., 2023; Bulut et al., 2016). Therefore, understanding and controlling the kinetic behavior of the $NaBH_4$ hydrolysis process are essential for the development of stable and efficient hydrogen generation systems.

To address these challenges, the present study investigates the hydrolysis kinetics of $NaBH_4$ for hydrogen production using a Co-Cu@g-C_3N_4 catalyst. The effects of two major experimental parameters, namely Co–Cu catalyst loading and initial reaction temperature, were systematically examined. Furthermore, a comprehensive kinetic investigation was conducted to determine the reaction order and activation energy of the $NaBH_4$ hydrolysis process. The results of this study are expected to provide valuable insights into the development of efficient and portable hydrogen generation systems based on $NaBH_4$ hydrolysis and Co–Cu-supported g-C_3N_4 catalysts.

METHODOLOGY

In this study, hydrogen was selected as a renewable and clean energy carrier. Sodium borohydride ($NaBH_4$), which offers a high hydrogen storage capacity, good stability, and safe handling characteristics, was employed as the hydrogen source to address the challenges associated with hydrogen storage. Since the hydrolysis reaction of $NaBH_4$ is strongly dependent on the catalyst, the development of an efficient, inexpensive, stable, and economically viable catalyst is of considerable importance. For this purpose, graphitic carbon nitride (g-C_3N_4), a material with a relatively high surface area and suitable structural properties, was synthesized and used as a support material. Subsequently, Co and Cu were incorporated onto the g-C_3N_4 support using the impregnation method to obtain Co-Cu@g-C_3N_4 catalysts. The synthesized catalysts were employed in the hydrolysis of $NaBH_4$ for hydrogen production and subsequently characterized using advanced analytical techniques. The hydrolysis reaction of $NaBH_4$ in water is given below:

$$NaBH_4 + 4H_2O \rightarrow NaBO_2 + 4H_2\uparrow$$

This reaction proceeds efficiently in the presence of suitable catalysts. The catalytic hydrolysis of $NaBH_4$ is an exothermic process, with a reported reaction enthalpy of approximately −210 kJ mol^{-1} at 25 °C. This reaction offers several important advantages for hydrogen generation. First, $NaBH_4$ is relatively safe and convenient to handle and store compared with gaseous hydrogen. Second, the hydrolysis of $NaBH_4$ theoretically enables the generation of four moles of H_2 per mole of $NaBH_4$, providing a high hydrogen yield (Andrieux et al., 2011; Karami and Fathirad, 2023;).

A simplified schematic representation of the material synthesis and experimental procedure is given below:

Urea → g-C_3N_4 synthesis → Co and Cu metal impregnation → $NaBH_4$ hydrolysis → Hydrogen production and characterization

RESEARCH AND RESULTS

Following the optimization of the reaction medium and catalyst amount, the effects of $NaBH_4$ concentration and reaction temperature on hydrogen production were investigated. Hydrogen generation profiles from the hydrolysis of $NaBH_4$ catalyzed by Co-Cu@g-C_3N_4 at different temperatures.are presented in Figure 1**.** The highest hydrogen generation rate was obtained at a $NaBH_4$ concentration of 3 wt.%, 25 ºC, 150 mg catalyst, reaching 4443 mL g^{-1} min^{-1}**.**

To investigate the catalytic reaction kinetics, $NaBH_4$ hydrolysis experiments were subsequently performed at different temperatures ranging from 25 to 40 °C, as shown in Figure 7. As can be observed from the figure, the reaction time decreased progressively with increasing temperature. This result indicates that an increase in reaction temperature accelerates the hydrolysis of $NaBH_4$ and consequently enhances the rate of hydrogen generation, demonstrating the significant influence of temperature on the catalytic performance and kinetic behavior of the Co-Cu@g-C_3N_4 catalyst.



Figure 1. Hydrogen generation profiles from the hydrolysis of $NaBH_4$ catalyzed by Co-Cu@g-C_3N_4 at different temperatures.

Using the temperature-dependent data and the Arrhenius equation, the reaction kinetics of H_2 generation were evaluated according to zero-order, first-order, and *n*th-order kinetic models. The results indicated that the Co-Cu@g-C_3N_4 catalyst supported the reaction predominantly through a zero-order kinetic pathway. When the ln *k* values obtained from the zero-order kinetic analysis of the data presented in Figure 2 were plotted against $1/T$ and substituted into the Arrhenius equation, the activation energy was determined to be 38.18 kJ mol^{-1}, as shown in Figure 2.



Figure 2. Arrhenius plot for the Co-Cu@g-C_3N_4 catalyst.

Another important characteristic of the catalyst is its reusability. After each reaction cycle, 90% of the remaining liquid phase was removed by decantation and transferred to another vessel, while fresh $NaBH_4$ solution was subsequently added to the remaining catalyst to initiate the next reaction cycle. In this manner, the catalyst was subjected to five consecutive reuse cycles. The results demonstrated that the catalyst maintained **100% conversion** even after the fifth cycle, indicating its excellent catalytic stability and reusability. The corresponding data are presented in **Figure 3**.

The gradual increase in reaction time observed with successive reuse cycles may be attributed to changes in the accessible active surface of the catalyst and the interaction of the catalyst surface with the metaborate ($NaBO_2$) by-product formed during the hydrolysis reaction (Xu et al., 2024; Ekinci and Onat, 2025; Wang et al., 2025). These effects may partially limit the accessibility of active catalytic sites and consequently lead to a slight decrease in catalytic efficiency during repeated use.



Figure 3. Reusability experiments of the Co-Cu@g-C_3N_4 catalyst.

CONCLUSION AND DISCUSSION

In this study, Co and Cu metals were immobilized onto a g-C_3N_4 support material synthesized from urea, and the hydrolysis kinetics of sodium borohydride ($NaBH_4$) over the resulting Co-Cu@g-C_3N_4 catalyst were investigated. The results demonstrated that the reaction followed zero-order kinetics**,** with an activation energy of approximately 38.18 kJ mol^{-1}**.** The observation of zero-order kinetics at a high borohydride-to-catalyst molar ratio can be attributed to the surface concentration and adsorption behavior of BH_4^- species on the active sites of the catalyst.

Furthermore, the catalytic stability and reusability of the catalyst were evaluated through five consecutive reaction cycles. The catalyst maintained its catalytic activity throughout the repeated cycles, with no significant change in its catalytic performance. These results demonstrate the good stability and reusability of the Co-Cu@g-C_3N_4 catalyst for hydrogen production via $NaBH_4$ hydrolysis.

REFERENCES

Alshammari, K., Alotaibi, T., Alshammari, M., Alhassan, S., Alshammari, A. H., & Taha, T. A. M. (2023). Synthesis of sulfur@ g-C3N4 and CuS@ g-C3N4 catalysts for hydrogen production from sodium borohydride. Materials, 16(12), 4218.

Andrieux, J., Demirci, U. B., & Miele, P. (2011). Langmuir–Hinshelwood kinetic model to capture the cobalt nanoparticles-catalyzed hydrolysis of sodium borohydride over a wide

temperature range. Catalysis today, 170(1), 13-19.

Arutyunov, V. S., & Lisichkin, G. V. (2017). Energy resources of the 21st century: problems and forecasts. Can renewable energy sources replace fossil fuels. Russian Chemical Reviews, 86(8), 777.

Bulut, A., Yurderi, M., Ertas, İ. E., Celebi, M., Kaya, M., & Zahmakiran, M. (2016). Carbon dispersed copper-cobalt alloy nanoparticles: A cost-effective heterogeneous catalyst with exceptional performance in the hydrolytic dehydrogenation of ammonia-borane. Applied Catalysis B: Environmental, 180, 121-129.

Cong, H., Ren, J., Zhang, D., Xu, F., Wang, X., Wang, Y., ... & Wu, S. (2023). Catalytic hydrolysis of sodium borohydride for hydrogen generation using g-C3N4/Co–W–B/Ni foam composite catalyst. Journal of Materials Science, 58(2), 787-801.

Ekinci, S., & Onat, E. (2024). Activated carbon assisted cobalt catalyst for hydrogen production: synthesis and characterization. Balıkesir Üniversitesi Fen Bilimleri Enstitüsü Dergisi, 26(2), 455-471.

Ekinci, S., & Onat, E. (2025). A novel and highly active Co-Cr@ Perlite/GO catalyst for sustainable hydrogen production via potassium borohydride hydrolysis. Applied Organometallic Chemistry, 39(8), e70307.

Karami, M., & Fathirad, F. (2023). Cobalt ferrite nanoparticles anchored on reduced graphene oxide nanoribbons (0D/1D CoFe2O4/rGONRs) as an efficient catalyst for hydrogen generation via NaBH4 hydrolysis. Inorganic Chemistry Communications, 150, 110552.

Onat, E., Ekinci, S., Şahin, Ö., Levent, A., & İzgi, M. S. (2026). Spent tea leaves to multifunctional environmental materials: Boric-acid-activated carbons for acetamiprid removal and energy storage, and a green Co catalyst for NaBH4 hydrolysis. Chemical Engineering Journal, 176001.

Xu, F., Ren, J., Ma, J., Wang, Y., Zhang, K., Cao, Z., ... & Bai, S. (2024). A review of hydrogen production kinetics from the hydrolysis of NaBH4 solution catalyzed by Co-based catalysts. International Journal of Hydrogen Energy, 50, 827-844.

Wang, Y., Zhou, Q., Zhang, Z., Zhang, C., Li, F., & Li, Q. (2025). Efficient hydrogen generation via NaBH4 hydrolysis over Co3O4 catalysts: The role of calcination temperature, valence, and reaction kinetics. International Journal of Hydrogen Energy, 144, 374-385.

LAMİNE AHŞAP KİRİŞLERİN SAYISAL MODELLENMESİ: GÜNCEL YAKLAŞIMLAR, DENEYSEL DOĞRULAMA, ARAŞTIRMA BOŞLUKLARI VE GELECEK EĞİLİMLERİ

NUMERICAL MODELLING OF GLUED LAMINATED TIMBER (GLULAM) BEAMS: CURRENT APPROACHES, EXPERIMENTAL VALIDATION, RESEARCH GAPS, AND FUTURE TRENDS

Lecturer Dr. İsmail KILIÇ
Yozgat Bozok University, Yozgat Vocational School, Department of Design, Yozgat, Türkiye
ORCID ID: 0000-0003-3609-2616

ÖZET

Lamine ahşap (glulam) kirişler, yüksek dayanım/ağırlık oranı, büyük açıklıkları geçebilme yeteneği ve sürdürülebilir yapı malzemesi olmaları nedeniyle modern ahşap yapılarda yaygın olarak kullanılan mühendislik ürünleri arasında yer almaktadır. Geleneksel deneysel yöntemler glulam kirişlerin mekanik davranışının belirlenmesinde güvenilir sonuçlar sunmakla birlikte, farklı malzeme özellikleri, kesit geometrileri, yükleme koşulları ve takviye sistemlerinin kapsamlı olarak değerlendirilmesi önemli ölçüde zaman ve maliyet gerektirmektedir. Bu nedenle son yıllarda sonlu eleman yöntemi (Finite Element Method-FEM) başta olmak üzere sayısal modelleme yaklaşımları, glulam kirişlerin davranışının analiz edilmesi, deneysel sonuçların doğrulanması ve tasarım alternatiflerinin değerlendirilmesinde etkin biçimde kullanılmaktadır.

Bu çalışmada, lamine ahşap kirişlerin sayısal modellenmesine ilişkin güncel literatür kapsamlı bir derleme yaklaşımıyla incelenmiştir. Sonlu eleman modelleme stratejileri, kullanılan yazılımlar, malzeme modelleme yaklaşımları, sonlu eleman tipleri, deneysel doğrulama yöntemleri ve model doğruluğunu etkileyen temel parametreler sistematik olarak değerlendirilmiştir. Ayrıca son yıllarda yayımlanan çalışmalar ışığında mevcut araştırma boşlukları belirlenmiş ve gelecekte öne çıkması beklenen araştırma eğilimleri tartışılmıştır.

Literatür incelemesi sonucunda, doğrusal olmayan sonlu eleman modelleri ile ortotropik ve ilerlemeli hasar esaslı malzeme modellerinin glulam kirişlerin yük-sehim davranışı, gerilme dağılımı ve hasar mekanizmalarının tahmininde başarılı sonuçlar verdiği belirlenmiştir. Bununla birlikte, ahşabın doğal heterojen yapısının modellenmesi, yapıştırıcı ara yüzey davranışının gerçekçi şekilde temsil edilmesi, uzun süreli yükleme ve çevresel etkilerin analize dâhil edilmesi ile parametrik optimizasyon ve yapay zekâ destekli modelleme yaklaşımlarının literatürde henüz sınırlı düzeyde ele alındığı görülmüştür.

Sonuç olarak, deneysel doğrulama ile desteklenen gelişmiş sonlu eleman modellerinin, parametrik tasarım, optimizasyon algoritmaları ve yapay zekâ tabanlı analiz yöntemleriyle bütünleştirilmesinin glulam kirişlerin mekanik davranışının daha güvenilir ve gerçekçi biçimde modellenmesine önemli katkılar sağlayacağı değerlendirilmektedir. Çalışmanın, lamine ahşap kirişlerin sayısal modellenmesi alanındaki güncel gelişmelerin bütüncül olarak değerlendirilmesine katkı sağlaması ve gelecekte gerçekleştirilecek deneysel ve sayısal araştırmalara yol göstermesi amaçlanmaktadır.

Anahtar Kelimeler: Lamine ahşap kiriş, glulam, sayısal modelleme, sonlu eleman yöntemi, deneysel doğrulama, araştırma boşlukları, optimizasyon.

ABSTRACT

Glued laminated timber (glulam) beams are widely used as engineered wood products in modern timber structures due to their high strength-to-weight ratio, ability to span long distances, and sustainability. Although conventional experimental methods provide reliable results for evaluating the mechanical behaviour of glulam beams, comprehensive investigations involving different material properties, cross-sectional geometries, loading conditions, and reinforcement systems require considerable time and cost. Therefore, numerical modelling approaches, particularly the Finite Element Method (FEM), have become increasingly important for analysing the structural behaviour of glulam beams, validating experimental results, and evaluating alternative design solutions.

In this study, the current literature on the numerical modelling of glulam beams is comprehensively reviewed. Finite element modelling strategies, commonly used numerical software, material modelling approaches, finite element types, experimental validation methods, and the key parameters affecting model accuracy are systematically examined. Furthermore, current research gaps are identified based on recent studies, and emerging research trends are discussed.

The literature review indicates that nonlinear finite element models combined with orthotropic and progressive damage material models provide reliable predictions of load–deflection behaviour, stress distribution, and damage mechanisms in glulam beams. However, several challenges remain, including the accurate representation of the natural heterogeneity of wood, realistic modelling of adhesive interface behaviour, consideration of long-term loading and environmental effects, and the integration of parametric optimisation and artificial intelligence-based modelling approaches, all of which remain relatively underexplored.

In conclusion, integrating experimentally validated advanced finite element models with parametric design, optimisation algorithms, and artificial intelligence-based analytical methods is expected to significantly improve the accuracy and reliability of numerical simulations of glulam beams. This review aims to provide a comprehensive overview of recent developments in the numerical modelling of glulam beams and to offer guidance for future experimental and computational research in this field.

Keywords: Glued laminated timber (glulam) beams, numerical modelling, finite element method, experimental validation, research gaps, optimisation.

GİRİŞ

Lamine ahşap (glulam) kirişler, yüksek dayanım/ağırlık oranı, büyük açıklıkları geçebilme kabiliyeti ve sürdürülebilir yapı malzemesi olma potansiyeli nedeniyle modern ahşap yapılarda yaygın olarak kullanılan mühendislik ahşap ürünleri arasında yer almaktadır (Jaaranen ve Fink,

2024; Greffier ve diğ., 2024; Vafadar ve diğ., 2025). Üretim sürecinde dayanım sınıflarına göre seçilen lamellerin uygun yapıştırıcılarla bir araya getirilmesi, kesit ve malzeme özelliklerinin tasarım gereksinimlerine göre düzenlenmesine ve yapısal performansın iyileştirilmesine olanak sağlamaktadır (Vafadar ve diğ., 2025; EN 14080:2013). Bu özellikler, glulam kirişlerin köprüler, spor salonları, endüstriyel yapılar ve büyük açıklıklı çatı sistemleri gibi farklı taşıyıcı sistemlerde kullanılmasına olanak sağlamaktadır (Greffier ve diğ., 2024; Jaaranen ve Fink, 2024).

Ahşap yapı sistemlerinin gelişmesiyle birlikte glulam kirişlerin davranışının yalnızca deneysel yöntemlerle değerlendirilmesi, özellikle çok sayıda değişkenin incelendiği araştırmalarda zaman, maliyet ve deney altyapısı açısından önemli sınırlılıklar oluşturabilmektedir. Deneysel çalışmalar gerçek davranışın belirlenmesi ve sayısal modellerin doğrulanması açısından vazgeçilmez olmakla birlikte, farklı kesit geometrileri, malzeme özellikleri, yükleme koşulları ve takviye sistemlerinin geniş bir kombinasyonunun deneysel olarak incelenmesi önemli ölçüde zaman ve kaynak gerektirmektedir. Bu nedenle sonlu eleman yöntemi (Finite Element Method, FEM) başta olmak üzere sayısal modelleme teknikleri; glulam kirişlerin davranışının tahmin edilmesi, farklı tasarım ve malzeme parametrelerinin değerlendirilmesi ve deneysel sonuçların doğrulanmasında önemli araştırma araçları hâline gelmiştir (Jaaranen ve Fink, 2024; Paroissien ve diğ., 2024; Türker ve Kılınçarslan, 2024).

Son yıllarda gerçekleştirilen çalışmalar, sayısal modelleme yaklaşımlarının glulam kirişlerde yalnızca elastik davranışın tahmininde değil, aynı zamanda hasar ve kırılma mekanizmalarının değerlendirilmesi, rijitlik ve eğilme dayanımının öngörülmesi ve farklı lamel yerleşimlerinin (layup) parametrik olarak incelenmesinde de etkin biçimde kullanılabildiğini göstermektedir (Jaaranen ve Fink, 2024; Vafadar ve diğ., 2025). Jaaranen ve Fink (2024), 36 deneysel glulam kiriş üzerinden doğrulanan sonlu eleman temelli bir modelle eğilme rijitliğinin yüksek doğrulukla tahmin edilebildiğini ve kırılma desenlerinin gerçekçi biçimde temsil edilebildiğini bildirmiştir. Vafadar ve diğ. (2025) ise 62 yapısal ölçekli glulam kirişten elde edilen deneysel verilerle doğrulanan olasılıksal sayısal model aracılığıyla eğilme dayanımı, rijitlik ve hasar bölgelerinin tahmin edilebildiğini göstermiş; ayrıca farklı lamel yerleşimlerinin mekanik performansı iyileştirme potansiyelini ortaya koymuştur.

Bunun yanında, glulam kirişlerin sayısal modellenmesinde kullanılan malzeme ve hasar modelleri, eleman tipi, ağ yapısı ve doğrulama stratejisi gibi modelleme kabullerinin sayısal sonuçlar üzerinde önemli etkileri olabilmektedir. Lu ve diğ. (2024), glulam elemanlarda delaminasyon davranışının modellenmesinde ısı ve kütle transferi, kuruma gerilmeleri, plastisite ve kırılma mekaniğinin birlikte ele alınmasının önemini vurgulamıştır. Benzer şekilde Paroissien ve diğ. (2024), lamine ahşap yapılarda solid, solid-beam ve solid-shell tabanlı farklı sonlu eleman yaklaşımlarını karşılaştırarak, model doğrulama ve geçerleme süreçlerinin seçilen modelleme stratejisinin değerlendirilmesindeki önemini ortaya koymuştur.

Takviyeli glulam kirişlere yönelik sayısal çalışmalar da son yıllarda önemli ölçüde gelişmiştir. Raftery ve Harte (2011), düşük kaliteli glulam kirişlerin FRP plakalarla güçlendirilmesini deneysel olarak inceleyerek, FRP takviyesinin eğilme performansı ve taşıma kapasitesinin

artırılmasındaki etkisini ortaya koymuştur. Nowak ve diğ. (2013), tarihi eğrisel ahşap elemanların CFRP şeritlerle güçlendirilmesini deneysel ve sayısal olarak inceleyerek, güçlendirme etkisinin sayısal modeller aracılığıyla değerlendirilmesine yönelik bir örnek sunmuştur. Bu çalışmaların devamında Raftery ve Harte (2013), FRP ile güçlendirilmiş glulam kirişlerin doğrusal olmayan davranışını temsil etmek amacıyla sonlu eleman modelleri geliştirmiştir. Khelifa ve diğ. (2015) ise karbon fiber takviyeli polimerlerle güçlendirilmiş ahşap kirişlerin eğilme davranışını sonlu eleman yöntemiyle inceleyerek, FRP takviyesinin yapısal davranış üzerindeki etkisinin sayısal olarak değerlendirilebileceğini göstermiştir.

Bu çalışmaların ilerleyen aşamalarında, model belirsizliklerinin ve modelleme düzeyinin sonuçlar üzerindeki etkisi daha fazla önem kazanmıştır. Dániel ve diğ. (2022), CFRP plakalarla güçlendirilmiş ve güçlendirilmemiş glulam kirişlerin güvenilirlik temelli sayısal analizini gerçekleştirerek, malzeme belirsizliklerinin deformasyon ve gerilme sonuçları üzerindeki etkisini ortaya koymuştur. Szczecina (2023) ise CFRP bantla güçlendirilmiş glulam kirişlerde farklı sayısal model karmaşıklıklarının gerilme, sehim ve taşıma kapasitesine ilişkin sonuçlar üzerindeki etkisini inceleyerek, modelleme düzeyinin sonuçların yorumlanmasındaki önemini ortaya koymuştur.

Güncel çalışmalar ise farklı takviye türlerinin ve doğrusal olmayan davranışın sayısal olarak temsil edilmesine odaklanmaktadır. Younis ve diğ. (2025), lif takviyeli ahşap kirişlerin eğilme yüklemesi altındaki davranışını sonlu eleman yöntemiyle inceleyerek, farklı lif takviyesi sistemlerinin sayısal olarak değerlendirilmesine yönelik güncel bir örnek sunmuştur. Glišović ve diğ. (2025) ise içten CFRP takviyeli glulam kirişler için geliştirilen doğrusal olmayan sonlu eleman modelini deneysel sonuçlarla doğrulayarak, takviyeli glulam elemanların doğrusal olmayan davranışının sayısal olarak temsil edilebileceğini ortaya koymuştur.

Buna karşın, mevcut çalışmaların önemli bir kısmının belirli bir sayısal modelin geliştirilmesine, belirli bir hasar mekanizmasının incelenmesine veya belirli bir glulam sisteminin analizine odaklandığı görülmektedir. Sayısal modelleme yaklaşımlarını, malzeme ve hasar modellerini, deneysel doğrulama yöntemlerini ve gelecekte öne çıkması beklenen araştırma eğilimlerini birlikte değerlendiren kapsamlı derleme çalışmalarının sınırlı olduğu görülmektedir. Özellikle doğrulanmış sonlu eleman modellerinin parametrik analiz ve optimizasyon süreçleriyle bütünleştirilmesi ve sayısal modelleme ile makine öğrenmesi yaklaşımlarının birlikte kullanılması, gelişmekte olan araştırma alanları olarak öne çıkmaktadır (Jaaranen ve Fink, 2024; Vafadar ve diğ., 2025; Kılıç ve diğ., 2025).

Bu çalışmada, lamine ahşap kirişlerin sayısal modellenmesinde kullanılan güncel yaklaşımlar literatür taramasına dayalı olarak incelenmiş; sonlu eleman modelleme stratejileri, malzeme modelleme yaklaşımları, deneysel doğrulama yöntemleri ve gelecekteki araştırma eğilimleri sistematik olarak değerlendirilmiştir. Ayrıca mevcut araştırma boşlukları ortaya konularak, deneysel doğrulama, sayısal modelleme ve optimizasyon temelli gelecekteki çalışmalar için bir değerlendirme çerçevesi sunulması amaçlanmıştır.

LAMİNE AHŞAP KİRİŞLERİN SAYISAL MODELLENMESİNDE GÜNCEL YAKLAŞIMLAR

Sonlu Eleman Yönteminin (FEM) Lamine Ahşap Kirişlerde Kullanımı

Lamine ahşap kirişlerin yapısal davranışının belirlenmesinde sonlu eleman yöntemi (Finite Element Method, FEM), deneysel çalışmaların tamamlayıcısı olarak yaygın biçimde kullanılmaktadır. Özellikle farklı malzeme ve geometrik parametrelerin sayısal ortamda değerlendirilmesine olanak sağlaması nedeniyle FEM, glulam elemanların tasarımı ve analizinde önemli avantajlar sunmaktadır. Deneysel çalışmalar gerçek davranışın belirlenmesinde temel yöntem olmakla birlikte, kapsamlı deney programlarının zaman ve maliyet gerektirmesi nedeniyle sayısal modelleme teknikleri deneysel araştırmaları tamamlayan önemli araçlar hâline gelmiştir (Jaaranen ve Fink, 2024; Lu ve diğ., 2024).

Glulam kirişlerin sayısal analizinde Abaqus gibi ticari sonlu eleman yazılımlarından yararlanılmakta; bu yazılımlar doğrusal ve doğrusal olmayan analizlerin yanı sıra ortotropik malzeme davranışı, temas problemleri, yapıştırıcı ara yüzeyleri ve hasar gelişiminin modellenmesine olanak sağlamaktadır (Szczecina, 2023).

FRP ile güçlendirilmiş ahşap elemanların sayısal modellenmesine yönelik çalışmalar da sonlu eleman yönteminin farklı güçlendirme sistemlerinin değerlendirilmesinde kullanılabileceğini göstermektedir. Khelifa ve diğ. (2015), karbon fiber takviyeli polimerlerle güçlendirilmiş ahşap kirişlerin eğilme davranışını sonlu eleman yöntemiyle inceleyerek, güçlendirme sisteminin yapısal davranış üzerindeki etkisinin sayısal olarak değerlendirilebileceğini ortaya koymuştur. Bu tür çalışmalar, sonlu eleman modellerinin yalnızca güçlendirilmemiş glulam kirişlerin davranışının tahmininde değil, aynı zamanda kompozit takviye sistemlerinin etkilerinin incelenmesinde de kullanılabileceğini göstermektedir.

Son yıllarda gerçekleştirilen çalışmalar, özellikle doğrusal olmayan sonlu eleman modellerinin glulam kirişlerin yük–sehim davranışı, rijitlik, taşıma kapasitesi ve kırılma mekanizmalarının belirlenmesinde başarılı sonuçlar verebildiğini göstermektedir. Bununla birlikte, model doğruluğu ve hesaplanan yapısal tepki; kullanılan eleman tipi, ağ yoğunluğu (mesh) ve malzeme/hasar modelleme yaklaşımı gibi modelleme kabullerine bağlı olarak değişebilmektedir (Jaaranen ve Fink, 2024).

Malzeme Modellemesi

Lamine ahşap kirişlerin sayısal analizinde model doğruluğunu etkileyen temel unsurlardan biri malzeme modelidir. Ahşabın lif yönüne bağlı olarak farklı mekanik özellikler göstermesi nedeniyle, uygun malzeme modelinin seçimi sonlu eleman analizlerinin güvenilirliğini doğrudan etkileyebilmektedir. Glulam kirişlerin sayısal modellenmesine yönelik çalışmalarda ahşabın anizotropik davranışı dikkate alınmakta ve ortotropik malzeme modelleri tercih edilmektedir (Jaaranen ve Fink, 2024; Lu ve diğ., 2024).

Ahşap malzeme üç asal eksende (boyuna, radyal ve teğetsel) farklı elastik özellikler göstermektedir. Bu durum elastisite modülü, kayma modülü ve Poisson oranlarının yönlere

bağlı olarak farklı değerler almasına neden olmaktadır. Dolayısıyla izotropik malzeme kabulleri, ahşabın yönsel mekanik davranışının dikkate alınması gereken sayısal analizlerde sınırlı kalabilmektedir. Glulam kirişlerin sayısal modellenmesine yönelik çalışmalarda, ahşabın yönsel davranışını temsil etmek amacıyla ortotropik elastik malzeme modellerinin kullanıldığı görülmektedir (Szczecina, 2023; Vafadar ve diğ., 2025).

Glulam kirişlerin sayısal modellenmesinde doğrusal ortotropik elastik modellerin yanı sıra, özellikle kırılma ve hasar davranışının incelendiği çalışmalarda doğrusal olmayan malzeme modelleri ve ilerlemeli hasar (progressive damage) yaklaşımları da kullanılmaktadır. Bu yaklaşımlar, çekme ve basınç bölgelerinde oluşan doğrusal olmayan davranış ile hasar gelişiminin temsil edilmesine ve sayısal sonuçların deneysel bulgularla karşılaştırılmasına olanak sağlamaktadır. Jaaranen ve Fink (2024), continuum damage model (CDM) yaklaşımını kullanarak glulam kirişlerin hasar ve kırılma davranışını sonlu eleman modeli içerisinde temsil etmiş ve model sonuçlarını deneysel verilerle doğrulamıştır. Benzer şekilde Glišović ve diğ. (2025), ortotropik elastoplastik malzeme davranışı, Hill kriteri ve ilerlemeli hasar modeli kullanarak CFRP takviyeli glulam kirişlerin doğrusal olmayan davranışını modellemiş ve sayısal sonuçların deneysel sonuçlarla iyi uyum gösterdiğini ortaya koymuştur.

Bununla birlikte, malzeme modelinin doğruluğu yalnızca elastik parametrelerin tanımlanmasına bağlı değildir. Ahşabın doğal heterojen yapısı ve malzeme özelliklerindeki değişkenlik, glulam kirişlerin mekanik davranışını ve sayısal model sonuçlarını önemli ölçüde etkileyebilmektedir. Bu nedenle, ahşap malzemenin doğal değişkenliğini dikkate almak amacıyla stokastik ve olasılıksal modelleme yaklaşımlarından yararlanılmaktadır. Bu yaklaşımlar sayesinde malzeme özelliklerindeki belirsizliklerin sayısal analizlere dâhil edilmesi ve model tahminlerinin güvenilirliğinin artırılması amaçlanmaktadır (Vafadar ve diğ., 2025).

Glulam kirişlerde bir diğer önemli modelleme bileşeni ise lameller arasındaki yapıştırıcı ara yüzeyidir. Basitleştirilmiş sayısal modellerde yapıştırıcı tabakasının tam aderans (perfect bond) ile temsil edilmesi mümkün olmakla birlikte, özellikle delaminasyon ve ara yüzey hasarının incelendiği çalışmalarda kohezyon (cohesive zone) modelleri tercih edilmektedir. Bu yaklaşımlar, yapıştırıcı tabakasında meydana gelen gerilme birikimlerini ve ayrılma davranışını daha gerçekçi şekilde temsil edebilmektedir (Lu ve diğ., 2024).

Son yıllarda geliştirilen sayısal modeller yalnızca elastik davranışın tahminine değil, aynı zamanda gerilme dağılımı, kırılma mekanizmaları ve hasar gelişiminin modellenmesine de olanak sağlamaktadır. Bununla birlikte, literatürde farklı malzeme modelleme yaklaşımlarının kullanıldığı ve analiz sonuçlarının seçilen malzeme modeli, parametrelerin belirlenme yöntemi ve doğrulama yaklaşımına bağlı olarak değişebildiği görülmektedir. Bu durum, glulam kirişlerin sayısal modellenmesinde uygun malzeme modelinin seçilmesinin temel aşamalardan biri olduğunu göstermektedir. Farklı çalışmalarda kullanılan yazılımlar, malzeme modelleri ve doğrulama yöntemleri Tablo 1'de özetlenmiştir.

Tablo 1. Lamine ahşap kirişlerin sayısal modellenmesine ilişkin güncel literatür özeti

Çalışma	Kullanılan yazılım / modelleme ortamı	Malzeme / hasar modeli	Doğrulama yöntemi
Jaaranen ve Fink (2024)	MATLAB tabanlı FEM	Ortotropik + Continuum Damage Model (CDM) + Sequentially Linear Analysis	36 deneysel glulam kiriş üzerinde eğilme testleri
Lu vd. (2024)	Sonlu eleman tabanlı sayısal model	Isı ve kütle transferi + ortotropik malzeme davranışı + plastisite + kırılma mekaniği + delaminasyon modeli	Deneysel delaminasyon / malzeme özellikleri verileriyle karşılaştırma
Szczecina (2023)	Abaqus	Doğrusal ortotropik elastik ahşap + CFRP modeli	Deneysel eğilme sonuçlarıyla karşılaştırma
Vafadar vd. (2025)	Sonlu eleman tabanlı olasılıksal model	Olasılıksal malzeme özellikleri + değişkenlik/Monte Carlo yaklaşımı	62 yapısal ölçekli glulam kiriş üzerinde deneysel doğrulama

Sonlu Eleman Tipleri ve Modelleme Stratejileri

Lamine ahşap kirişlerin sayısal analizinde model doğruluğunu etkileyen temel unsurlardan biri seçilen sonlu eleman tipidir. Sonlu eleman modelleri; analizin amacı, geometrik karmaşıklık, hesaplama süresi ve beklenen doğruluk düzeyine bağlı olarak farklı yaklaşımlar kullanılarak oluşturulmaktadır. Glulam kirişlerin modellenmesinde beam, shell ve solid elemanlar başta olmak üzere farklı sonlu eleman yaklaşımlarından yararlanılmaktadır. Her bir yaklaşımın avantajları ve sınırlılıkları bulunduğundan, eleman tipi ve modelleme stratejisinin çalışma amacına uygun olarak belirlenmesi gerekmektedir (Paroissien ve diğ., 2024).

Çubuk (beam) eleman modelleri, geometrinin ve malzeme davranışının daha basitleştirilmiş biçimde temsil edilmesine olanak sağlamaları ve düşük hesaplama maliyetleri nedeniyle özellikle genel yapısal davranışın değerlendirilmesinde avantaj sağlamaktadır. Bununla birlikte, beam tabanlı modellerde lameller arasındaki yerel gerilme dağılımları, yapıştırıcı ara yüzeyleri ve yerel hasar mekanizmalarının ayrıntılı olarak temsil edilmesi, daha ayrıntılı shell veya solid modellere kıyasla sınırlı kalabilmektedir (Paroissien ve diğ., 2024).

Kabuk (shell) elemanlar, tabakalı yapıların modellenmesinde katman davranışının daha ayrıntılı biçimde temsil edilmesine olanak sağlayan bir yaklaşım sunmaktadır. Glulam elemanların modellenmesinde shell tabanlı yaklaşımlar, beam modellerine kıyasla yapısal elemanın kesit ve tabaka davranışının daha ayrıntılı temsil edilmesine olanak

sağlayabilmektedir. Bununla birlikte, üç boyutlu gerilme durumlarının ve yerel davranışların ayrıntılı olarak incelenmesi gereken durumlarda solid elemanlar daha kapsamlı bir modelleme yaklaşımı sunmaktadır (Paroissien ve diğ., 2024).

Katı (solid) eleman modelleri, glulam kirişlerin üç boyutlu davranışının ayrıntılı biçimde temsil edilmesine olanak sağlamaktadır. Bu modellerde her lamel ayrı bir hacim elemanı olarak tanımlanabilmekte, yapıştırıcı tabakaları ve temas yüzeyleri ayrıntılı biçimde modellenebilmektedir. Bu nedenle solid modeller, özellikle gerilme dağılımı, delaminasyon ve hasar/kırılma davranışlarının ayrıntılı olarak incelenmesinde avantaj sağlamaktadır. Bununla birlikte, ayrıntılı geometrik ve malzeme tanımlamaları nedeniyle solid eleman modellerinin oluşturulması ve çözülmesi daha yüksek hesaplama maliyeti gerektirebilmektedir (Lu ve diğ., 2024; Jaaranen ve Fink, 2024; Paroissien ve diğ., 2024).

Farklı eleman tiplerinin birlikte veya alternatif modelleme yaklaşımları içerisinde kullanılmasına yönelik stratejiler de glulam yapıların sayısal analizinde ele alınmaktadır. Bu yaklaşımlarda modelleme ayrıntısı ile hesaplama maliyeti arasında uygun bir denge kurulması amaçlanmaktadır. Özellikle solid-beam ve solid-shell gibi yaklaşımlar, daha ayrıntılı üç boyutlu modellerin sağladığı modelleme kapasitesi ile hesaplama maliyeti arasında farklı düzeylerde çözümler sunabilmektedir (Paroissien ve diğ., 2024).

Model doğruluğunu etkileyen bir diğer önemli unsur ağ yapısı (mesh) ve eleman boyutudur. Çok kaba ağ yapıları sayısal sonuçların doğruluğunu sınırlayabilirken, eleman boyutunun gereğinden fazla küçültülmesi hesaplama maliyetini artırabilmektedir. Bu nedenle sonlu eleman modelinin doğruluğu ile hesaplama maliyeti arasında uygun bir denge kurulması ve eleman boyutunun model sonuçları üzerindeki etkisinin değerlendirilmesi önem taşımaktadır (Jaaranen ve Fink, 2024). Özellikle karmaşık hasar ve kırılma davranışlarının modellendiği glulam kirişlerde, ağ yapısının yeterli çözünürlük sağlayacak şekilde belirlenmesi model sonuçlarının güvenilirliği açısından önem taşımaktadır.

Genel olarak değerlendirildiğinde, glulam kirişlerin sayısal modellenmesinde kullanılacak eleman tipi; analizin amacı, beklenen doğruluk düzeyi ve hesaplama maliyeti dikkate alınarak seçilmelidir. Beam elemanlar genel yapısal davranışın ve küresel tepkinin değerlendirilmesinde düşük hesaplama maliyeti sunarken, shell ve solid yaklaşımlar daha ayrıntılı kesit ve üç boyutlu davranışların incelenmesine olanak sağlamaktadır. Özellikle solid modeller, hasar, delaminasyon ve kırılma gibi yerel davranışların ayrıntılı olarak incelenmesinde avantaj sağlarken, daha yüksek hesaplama maliyeti gerektirebilmektedir. Farklı eleman tiplerinin temel avantajları, sınırlılıkları ve kullanım alanları Tablo 2'de karşılaştırılmıştır (Paroissien ve diğ., 2024; Lu ve diğ., 2024; Jaaranen ve Fink, 2024).

Tablo 2. Glulam kirişlerin sayısal modellenmesinde kullanılan eleman tiplerinin karşılaştırılması

Eleman tipi	Avantajları	Dezavantajları	Kullanım alanı
Beam	Düşük hesaplama maliyeti, hızlı çözüm	Yerel gerilme ve hasar davranışının ayrıntılı temsilinde sınırlı	Global yapısal davranış, ön değerlendirme
Shell	Tabakalı davranışın ve kesit özelliklerinin daha ayrıntılı temsil edilebilmesi	Ayrıntılı üç boyutlu gerilme ve ara yüzey davranışının temsilinde solid modellere göre sınırlı	Tabakalı yapıların ve kesit davranışının modellenmesi
Solid	Üç boyutlu davranış ve yerel hasarın ayrıntılı temsili	Yüksek hesaplama maliyeti ve modelleme karmaşıklığı	Hasar, delaminasyon, kırılma ve ayrıntılı üç boyutlu analizler
Hibrit	Doğruluk ile hesaplama maliyeti arasında denge sağlayabilmesi	Modelleme ve bağlantı tanımlarının daha karmaşık olması	Farklı modelleme düzeylerinin birlikte değerlendirildiği analizler

Deneysel Doğrulama Yaklaşımları

Sayısal modelleme çalışmalarında elde edilen sonuçların güvenilirliğinin değerlendirilmesinde, geliştirilen modelin deneysel verilerle doğrulanması temel aşamalardan biridir. Lamine ahşap kirişlere yönelik sonlu eleman çalışmalarında deneysel doğrulama; modelin yük taşıma kapasitesi, rijitlik, deformasyon davranışı ve hasar veya kırılma mekanizmalarını temsil etme düzeyinin değerlendirilmesine olanak sağlamaktadır. Nitekim Jaaranen ve Fink (2024), geliştirdikleri sonlu eleman modelini 36 deneysel olarak test edilmiş glulam kiriş üzerinden doğrulayarak eğilme rijitliği, eğilme dayanımı ve kırılma biçimleri bakımından model performansını değerlendirmiştir. Benzer şekilde Lu ve diğ. (2024), glulam delaminasyon davranışını inceleyen sonlu eleman modelinin farklı aşamalarını deneysel verilerle doğrulamıştır. Bu çalışmalar, deneysel verilerin sayısal modellerin güvenilirliğinin değerlendirilmesinde önemli bir referans oluşturduğunu göstermektedir.

Lamine ahşap kirişlerin sayısal modellerinin deneysel doğrulanmasında üç noktalı ve dört noktalı eğilme deneylerinden elde edilen veriler önemli bir referans oluşturmaktadır. Bu deneyler sonucunda elde edilen yük–sehim eğrileri, maksimum yük taşıma kapasitesi, eğilme dayanımı (MOR), elastisite modülü (MOE) ve kırılma biçimleri sayısal analiz sonuçlarıyla karşılaştırılabilmektedir. Özellikle yük–sehim eğrisinin genel davranışı ile maksimum yük değerlerinin deneysel sonuçlarla uyumu, geliştirilen modelin yapısal davranışı temsil etme

düzeyinin değerlendirilmesinde kullanılan temel göstergeler arasında yer almaktadır (European Committee for Standardization, 2010; Jaaranen ve Fink, 2024).

Sonlu eleman modellerinin doğrulanmasında yalnızca maksimum yük değerlerinin karşılaştırılması yerine, modelin deneysel davranışı farklı aşamalarda ne ölçüde temsil edebildiğinin değerlendirilmesi önem taşımaktadır. Bu kapsamda yük–sehim eğrisinin genel karakteristiği, rijitlik, taşıma kapasitesi ve hasar veya kırılma davranışı gibi farklı çıktıların deneysel sonuçlarla karşılaştırılması, model performansının daha kapsamlı biçimde değerlendirilmesine olanak sağlamaktadır. Böylece modelin yalnızca nihai taşıma kapasitesini değil, elastik davranıştan hasar gelişimine kadar uzanan yapısal tepkiyi temsil etme düzeyi incelenebilmektedir. Özellikle hasar ve kırılma davranışını dikkate alan sayısal modellerde, deneysel kırılma biçimleri ile sayısal olarak öngörülen hasar ve kırılma mekanizmalarının karşılaştırılması model doğrulamasının önemli bir bileşenini oluşturmaktadır (Jaaranen ve Fink, 2024; Vafadar ve diğ., 2025; Vida ve diğ., 2025).

Deneysel doğrulamanın başarısını etkileyen önemli unsurlardan biri, sınır şartlarının ve yükleme koşullarının deney düzenine uygun biçimde sayısal modele aktarılmasıdır. Mesnet koşulları, yük uygulama bölgeleri ve temas tanımlarının deneysel düzenekle uyumlu olarak modellenmesi, sayısal ve deneysel sonuçların karşılaştırılabilirliği açısından önem taşımaktadır. Bunun yanında, ahşabın doğal değişkenliği ve malzeme özelliklerindeki farklılıklar nedeniyle sayısal modellerde kullanılan malzeme parametrelerinin belirlenmesi de model doğruluğunun önemli bileşenlerinden biridir. Vafadar ve diğ. (2025), glulam kirişlerin sayısal modellenmesinde malzeme ve yapısal değişkenliği dikkate alan olasılıksal bir yaklaşım kullanırken, Lu ve diğ. (2024) malzeme davranışı ve delaminasyonun modellenmesinde farklı fiziksel ve mekanik süreçleri dikkate almıştır. Bu çalışmalar, deneysel doğrulamada sınır şartlarının yanı sıra malzeme özelliklerinin de uygun şekilde tanımlanmasının önemini ortaya koymaktadır.

Son yıllarda deneysel doğrulama ve yapısal davranışın izlenmesine yönelik çalışmalarda, yalnızca yük–sehim ölçümlerinin değil, farklı sensör ve görüntüleme tekniklerinin de kullanıldığı görülmektedir. Gerinim ölçerler (strain gauges), LVDT sensörleri ve dijital görüntü korelasyonu (Digital Image Correlation, DIC) gibi yöntemler, yükleme sırasında oluşan deformasyonların, gerinim değişimlerinin ve hasar gelişiminin daha ayrıntılı biçimde izlenmesine olanak sağlamaktadır. Greffier ve diğ. (2024), glulam numunelerde gerinim ölçerler ve LVDT sensörleri kullanarak nem değişimleri ve mekanik davranış arasındaki ilişkiyi incelemiş ve dört noktalı eğilme deneyleri gerçekleştirmiştir. Lu ve diğ. (2024) ise glulam delaminasyonunun sayısal modellenmesine yönelik deneysel çalışmalarda DIC tekniğinden yararlanarak kuruma sürecinde numune yüzeyindeki gerinim değişimlerini izlemiştir. Bu tür ölçüm tekniklerinin kullanılması, deneysel davranışın daha ayrıntılı olarak ortaya konulmasına ve sayısal modellerin yalnızca taşıma kapasitesi açısından değil, deformasyon ve hasar davranışı bakımından da değerlendirilmesine katkı sağlamaktadır (Greffier ve diğ., 2024; Lu ve diğ., 2024).

Genel olarak değerlendirildiğinde, deneysel doğrulama süreci sayısal modelleme çalışmalarının temel bileşenlerinden biridir. Güvenilir bir sonlu eleman modelinin geliştirilmesi yalnızca uygun malzeme modeli ve eleman tipinin seçilmesine değil, aynı zamanda deneysel verilerle desteklenen kapsamlı bir doğrulama sürecine de bağlıdır. Bu kapsamda yük–sehim davranışı, rijitlik, taşıma kapasitesi ve hasar veya kırılma mekanizmaları gibi farklı deneysel çıktılar birlikte değerlendirilerek modelin yapısal davranışı temsil etme düzeyi daha kapsamlı biçimde ortaya konulabilmektedir. Deneysel doğrulama kapsamında kullanılan temel parametreler ve bunların sayısal modellerde karşılık geldiği çıktılar Tablo 3'te özetlenmiştir. Doğrulanmış modellerin daha sonraki aşamalarda parametrik analiz, optimizasyon ve farklı tasarım senaryolarının değerlendirilmesinde kullanılması, sayısal modelleme çalışmalarının uygulama kapsamını genişletebilmektedir.

Tablo 3. Glulam kirişlerin sayısal modellerinin doğrulanmasında kullanılan temel parametreler

Doğrulama parametresi	Deneysel yöntem	Sayısal çıktı	Kullanım / değerlendirme
Yük–sehim eğrisi	3N / 4N eğilme deneyi	Yük–yer değiştirme ilişkisi	Temel doğrulama parametresi
Maksimum yük / MOR	Eğilme deneyi	Maksimum taşıma kapasitesi / eğilme dayanımı	Temel doğrulama parametresi
MOE / rijitlik	Eğilme deneyi	Rijitlik / elastik davranış	Temel doğrulama parametresi
Gerilme–gerinim davranışı	Gerinim ölçer / DIC	Gerilme ve gerinim alanları	Destekleyici doğrulama parametresi
Kırılma modu	Görsel inceleme / deneysel gözlem	Hasar ve kırılma modeli	Önemli doğrulama parametresi
Delaminasyon / ara yüzey hasarı	Görsel inceleme / DIC	Ara yüzey hasarı / delaminasyon	Özel durumlarda doğrulama parametresi
Çatlak ve hasar gelişimi	DIC / görüntüleme	Progressive damage / hasar dağılımı	Ayrıntılı hasar doğrulaması

LAMİNE AHŞAP KİRİŞLERİN SAYISAL MODELLENMESİNDE GÜNCEL ARAŞTIRMA BOŞLUKLARI VE GELECEK EĞİLİMLERİ

Son yıllarda lamine ahşap (glulam) kirişlerin sayısal modellenmesine yönelik araştırmalarda önemli ilerlemeler kaydedilmiştir. Bununla birlikte mevcut çalışmaların önemli bir bölümü belirli modelleme yaklaşımlarına, eğilme davranışının tahminine ve deneysel doğrulamaya odaklanmaktadır. Farklı çalışmalarda malzeme davranışı, hasar mekanizmaları, modelleme

düzeyi ve deneysel doğrulama yöntemleri açısından farklı yaklaşımların kullanılması, glulam kirişlerin sayısal modellenmesinde tek bir standart modelleme stratejisinin henüz öne çıkmadığını göstermektedir (Jaaranen ve Fink, 2024; Vafadar ve diğ., 2025; Paroissien ve diğ., 2024).

Literatürde dikkati çeken araştırma konularından biri, ahşabın doğal malzeme değişkenliğinin sayısal modellere aktarılmasıdır. Glulam kirişlerin sonlu eleman analizlerinde ahşap çoğunlukla ortotropik malzeme özellikleri üzerinden modellenmekle birlikte, malzeme özelliklerindeki doğal değişkenlik sayısal sonuçlar üzerinde etkili olabilmektedir. Vafadar ve diğ. (2025), glulam kirişlerin sayısal modellenmesinde malzeme ve yapısal değişkenliği dikkate alan olasılıksal bir yaklaşım geliştirerek deneysel sonuçlarla doğrulama gerçekleştirmiştir. Lu ve diğ. (2024) ise glulam davranışının modellenmesinde malzemenin ısı ve kütle transferi, plastisite ve kırılma davranışıyla ilişkili karmaşık özelliklerini dikkate almıştır. Bu çalışmalar, gelecekte glulam kirişlerin sayısal analizlerinde malzeme değişkenliğinin ve fiziksel davranışın daha kapsamlı biçimde temsil edilmesine yönelik yaklaşımların önemini ortaya koymaktadır.

Bir diğer önemli araştırma alanı, glulam kirişlerde lameller arasındaki yapıştırıcı ara yüzeylerinin sayısal olarak temsil edilmesidir. Sayısal modellerde lameller arasındaki bağlantının tam aderans (perfect bond) kabulüyle idealize edilmesi, yapıştırıcı tabakasının gerçek davranışının modele doğrudan aktarılmasını sınırlayabilmektedir. Özellikle ara yüzey hasarı ve delaminasyon davranışının incelendiği durumlarda, yapıştırıcı tabakasındaki gerilme gelişiminin ve ayrılma davranışının ayrıca modellenmesi önem kazanmaktadır. Lu ve diğ. (2024), glulam elemanlarda delaminasyon davranışını sonlu eleman yöntemiyle inceleyerek ısı ve kütle transferi, plastisite, kırılma mekaniği ve ara yüzey davranışını birlikte ele alan kapsamlı bir modelleme yaklaşımı kullanmıştır. Bu çalışma, yapıştırıcı ara yüzeylerinin ve delaminasyon mekanizmalarının glulam kirişlerin sayısal analizlerinde daha ayrıntılı biçimde ele alınmasının önemini ortaya koymaktadır.

Mevcut çalışmaların önemli bir bölümü glulam kirişlerin kısa süreli statik yükleme altındaki davranışına ve bu davranışın sonlu eleman modelleriyle temsil edilmesine odaklanmaktadır. Buna karşılık malzeme ve çevresel değişkenliğin sayısal modellere aktarılması, özellikle uzun dönemli yapısal davranışın değerlendirilmesi açısından geliştirilmesi gereken konular arasında yer almaktadır. Vafadar ve diğ. (2025), glulam kirişlerin sayısal modellenmesinde malzeme ve yapısal değişkenliği olasılıksal bir yaklaşımla ele alarak deneysel doğrulama gerçekleştirmiştir. Bu yaklaşım, gelecekte çevresel koşulların ve zamana bağlı malzeme davranışlarının daha kapsamlı biçimde dikkate alındığı sayısal modellerin geliştirilmesi için önemli bir araştırma yönü ortaya koymaktadır.

Takviyeli glulam sistemlerine yönelik araştırmalarda farklı güçlendirme malzemeleri ve uygulama yöntemlerinin kullanıldığı görülmektedir. CFRP ve diğer FRP tabanlı takviye sistemleri üzerine gerçekleştirilen çalışmalar, takviyenin glulam kirişlerin taşıma kapasitesi, rijitliği ve kırılma davranışı üzerindeki etkilerinin hem deneysel hem de sayısal yöntemlerle incelenebildiğini göstermektedir (Raftery ve Harte, 2011; Fiorelli ve Dias, 2011; Raftery ve Harte, 2013; Yang ve diğ., 2016; Glišović ve diğ., 2017; Dániel ve diğ., 2022; İşleyen ve diğ.,

2023; Glišović ve diğ., 2025; Muhaxheri ve diğ., 2026). Bununla birlikte mevcut çalışmaların önemli bir bölümü belirli bir takviye malzemesi veya belirli bir güçlendirme konfigürasyonu üzerine yoğunlaşmaktadır. Farklı takviye türlerinin aynı modelleme ve doğrulama çerçevesi içerisinde sistematik olarak karşılaştırılması, glulam kirişlerin tasarımında farklı güçlendirme alternatiflerinin değerlendirilmesi açısından geliştirilmeye açık bir araştırma alanı olarak değerlendirilebilir. Farklı doğal ve yapay liflerin takviye sistemlerinde kullanımına yönelik güncel çalışmalar ise takviye malzemesi ve uygulama biçiminin yapısal davranış üzerindeki etkilerinin daha ayrıntılı incelenmesi gerektiğine işaret etmektedir (Wdowiak-Postulak ve diğ., 2025). Ayrıca takviye sistemlerinin yalnızca taşıma kapasitesi ve rijitlik üzerindeki etkilerinin değil, malzeme kullanımı, tasarım verimliliği ve uygulama koşulları gibi daha geniş mühendislik ölçütleriyle birlikte değerlendirilmesi gelecekteki çalışmalar için önemli bir araştırma yönü oluşturmaktadır.

Son yıllarda dikkat çeken araştırma eğilimlerinden biri, yapay zekâ ve veri odaklı yöntemlerin ahşap yapı elemanlarının mekanik davranışının tahmininde kullanılmaya başlanmasıdır. Makine öğrenmesi algoritmaları, deneysel ve sayısal çalışmalardan elde edilen veri kümelerinin değerlendirilmesi, mekanik performansın tahmin edilmesi ve tasarım değişkenleri ile taşıma kapasitesi arasındaki ilişkilerin ortaya konulması açısından önemli olanaklar sunmaktadır. Kılıç ve diğ. (2025), makine öğrenmesi modellerini kullanarak lamine ahşap kirişlerin eğilme dayanımı (MOR) değerlerinin tahmin edilebileceğini göstermiştir. Bununla birlikte, mevcut çalışmaların önemli bir bölümü mekanik performansın tahminine odaklanmakta olup, doğrulanmış sonlu eleman modellerinden elde edilen veriler ile makine öğrenmesi yöntemlerinin bütünleştirildiği kapsamlı yaklaşımların geliştirilmesi önemli bir araştırma alanı olarak değerlendirilebilir.

Bunun yanında parametrik tasarım ve farklı tasarım değişkenlerinin sayısal modeller içerisinde değerlendirilmesi, glulam kirişlerin mühendislik tasarımının geliştirilmesi açısından önemli bir araştırma alanı oluşturmaktadır. Kesit özellikleri, lamel dizilimi ve malzeme değişkenleri gibi parametrelerin sayısal modeller aracılığıyla değiştirilmesi, farklı tasarım alternatiflerinin taşıma kapasitesi ve rijitlik açısından karşılaştırılmasına olanak sağlamaktadır. Vafadar ve diğ. (2025), farklı layup düzenlerinin glulam kirişlerin mekanik davranışı üzerindeki etkisini olasılıksal bir sayısal model ve deneysel doğrulama çerçevesinde incelemiştir. Jaaranen ve Fink (2024) ise doğrulanmış sonlu eleman modeli üzerinden glulam kirişlerin davranışının farklı modelleme koşulları altında değerlendirilmesine olanak sağlayan bir yaklaşım ortaya koymuştur. Zhang ve diğ. (2026), CLT–glulam kompozit kirişlerin eğilme davranışını sayısal simülasyon ve teorik analiz yoluyla inceleyerek, farklı mühendislik ahşap ürünlerinin birlikte kullanıldığı taşıyıcı sistemlerin sayısal olarak değerlendirilmesine yönelik güncel bir örnek sunmuştur. Bu tür kompozit sistemlerin modellenmesi, gelecekte glulam kirişlerin yalnızca tekil eleman düzeyinde değil, farklı mühendislik ahşap ürünleriyle birlikte oluşturduğu taşıyıcı sistemler düzeyinde de değerlendirilmesine olanak sağlayabilir. Bununla birlikte, farklı tasarım değişkenlerinin deneysel doğrulama ile birlikte ele alındığı ve doğrudan mühendislik tasarımına

aktarılabildiği kapsamlı parametrik ve optimizasyon çalışmalarının geliştirilmesi gelecekteki araştırmalar açısından önemli bir fırsat sunmaktadır.

Genel olarak değerlendirildiğinde, gelecekte gerçekleştirilecek araştırmalarda deneysel doğrulama, gelişmiş sonlu eleman modelleri, parametrik analiz, optimizasyon ve veri odaklı yöntemlerin birlikte değerlendirilmesi, glulam kirişlerin sayısal olarak daha kapsamlı biçimde incelenmesine katkı sağlayabilir. Özellikle malzeme değişkenliğinin dikkate alındığı doğrulanmış sayısal modellerin farklı tasarım senaryoları üzerinde kullanılması ve makine öğrenmesi yöntemleriyle desteklenmesi, taşıma kapasitesi ve yapısal davranışın tahmin edilmesine yönelik yeni olanaklar sunmaktadır. Bu kapsamda mevcut araştırma boşlukları ve gelecekte öne çıkabilecek araştırma alanları Tablo 4'te özetlenmiştir.

Tablo 4. Lamine ahşap kirişlerin sayısal modellenmesine ilişkin başlıca araştırma boşlukları ve gelecekteki araştırma alanları

Araştırma alanı	Mevcut durum	Belirlenen boşluk	Gelecek eğilim
Malzeme modeli	Ortotropik modeller yaygın	Malzeme değişkenliğinin sınırlı temsili	Olasılıksal ve değişkenliği dikkate alan modeller
Yapıştırıcı modeli	Tam aderans (perfect bond) kabulü yaygın	Ara yüzey davranışının daha ayrıntılı temsil edilmesi	Cohesive zone ve delaminasyon modelleri
Eleman ve modelleme stratejisi	Beam, shell ve solid yaklaşımları kullanılmaktadır	Tek bir modelleme stratejisinin tüm davranışlar için yeterli olmaması	Hibrit ve çok düzeyli modelleme
Deneysel doğrulama	Yük–sehim, dayanım ve rijitlik karşılaştırmaları yaygın	Hasar ve kırılma davranışının daha kapsamlı doğrulanması	Gelişmiş ölçüm teknikleriyle bütünleşik doğrulama
Takviyeli sistemler	Özellikle CFRP/FRP takviyeli glulam çalışmalarına ağırlık verilmektedir	Farklı takviye sistemlerinin ortak bir çerçevede karşılaştırılması	Farklı takviye malzemelerinin sistematik karşılaştırılması
Parametrik analiz	Belirli tasarım değişkenleri ayrı ayrı incelenmektedir	Çok sayıda tasarım değişkeninin birlikte değerlendirilmesi	Parametrik tasarım ve optimizasyon
Yapay zekâ	Lamine ahşap kirişlerin mekanik performansının tahmininde kullanılmaktadır	Sayısal modelleme ve deneysel doğrulama ile daha güçlü bütünleştirilmesi	FEM + makine öğrenmesi tabanlı yaklaşımlar

SONUÇLAR VE DEĞERLENDİRME

Bu çalışmada, lamine ahşap (glulam) kirişlerin sayısal modellenmesine yönelik güncel yaklaşımlar kapsamlı bir literatür incelemesi çerçevesinde değerlendirilmiştir. İncelenen çalışmalar, sonlu eleman yönteminin glulam kirişlerin mekanik davranışının belirlenmesinde etkili bir araç olarak kullanılabildiğini göstermektedir. Özellikle yük–sehim davranışı, eğilme rijitliği, taşıma kapasitesi ve hasar mekanizmalarının tahmininde geliştirilen sayısal modellerin deneysel sonuçlarla başarılı biçimde doğrulanabildiği görülmüştür (Jaaranen ve Fink, 2024; Vafadar ve diğ., 2025; Vida ve diğ., 2025).

Literatür değerlendirmesi sonucunda, glulam kirişlerin modellenmesinde ortotropik elastik malzeme modellerinin yaygın olarak kullanıldığı, bununla birlikte hasar ve kırılma davranışının incelendiği çalışmalarda ilerlemeli hasar ve continuum damage modellerinin, ara yüzey davranışının incelendiği çalışmalarda ise cohesive zone yaklaşımlarının kullanıldığı belirlenmiştir. Ayrıca olasılıksal modelleme yaklaşımlarının, ahşabın doğal malzeme ve yapısal değişkenliğinin sayısal analizlere aktarılması açısından önemli bir gelişme sunduğu görülmüştür. Model doğruluğunun yalnızca malzeme modeliyle sınırlı olmadığı; eleman tipi, ağ yapısı, sınır şartları, ara yüzey tanımları ve deneysel doğrulama stratejilerinin de sonuçlar üzerinde belirleyici olduğu değerlendirilmiştir (Jaaranen ve Fink, 2024; Lu ve diğ., 2024; Paroissien ve diğ., 2024; Vafadar ve diğ., 2025).

Çalışmada incelenen araştırmalar, sonlu eleman modellerinin deneysel verilerle doğrulanmasının model güvenilirliğinin değerlendirilmesinde temel bir aşama olduğunu göstermektedir. Doğrulama sürecinde yalnızca maksimum yük taşıma kapasitesinin değil, yük–sehim davranışı, rijitlik, eğilme dayanımı, deformasyon ve hasar veya kırılma davranışının birlikte değerlendirilmesi, modelin yapısal davranışı temsil etme düzeyinin daha kapsamlı biçimde ortaya konulmasını sağlamaktadır. Bunun yanında gerinim ölçerler, LVDT sensörleri ve dijital görüntü korelasyonu (DIC) gibi gelişmiş deneysel ölçüm tekniklerinin kullanılması, sayısal modellerin deformasyon ve hasar davranışı açısından daha ayrıntılı biçimde değerlendirilmesine olanak sağlamaktadır (Lu ve diğ., 2024; Greffier ve diğ., 2024; Vafadar ve diğ., 2025).

Literatürdeki çalışmaların değerlendirilmesi sonucunda bazı önemli araştırma boşlukları da ortaya çıkmaktadır. Özellikle ahşabın doğal malzeme değişkenliğinin daha ayrıntılı biçimde temsil edilmesi, yapıştırıcı ara yüzeylerinin ve delaminasyon davranışının modellenmesi, uzun dönemli ve çevresel etkilerin sayısal analizlere aktarılması ve farklı takviye sistemlerinin ortak bir modelleme ve doğrulama çerçevesinde karşılaştırılması geliştirmeye açık araştırma alanları olarak değerlendirilmektedir. Bunun yanında parametrik tasarım ve optimizasyon çalışmalarının deneysel olarak doğrulanmış sayısal modellerle daha bütünleşik biçimde yürütülmesi ve makine öğrenmesi yöntemlerinin sayısal modelleme süreçleriyle birlikte kullanılması gelecekte önemli araştırma olanakları sunmaktadır (Lu ve diğ., 2024; Vafadar ve diğ., 2025; Kılıç ve diğ., 2025).

Genel olarak değerlendirildiğinde, glulam kirişlerin gelecekteki sayısal analizlerinde deneysel doğrulama, gelişmiş sonlu eleman modelleri, malzeme değişkenliğinin dikkate alınması, parametrik analiz, optimizasyon ve veri odaklı yöntemlerin birlikte değerlendirilmesi önemli katkılar sağlayabilir. Özellikle deneysel olarak doğrulanmış sayısal modellerin farklı tasarım senaryolarında kullanılması ve makine öğrenmesi yöntemleriyle desteklenmesi, glulam kirişlerin taşıma kapasitesi ve mekanik davranışının daha etkin biçimde tahmin edilmesine olanak sağlayabilir. Bu bütünleşik yaklaşımın, glulam kirişlerin mühendislik tasarımında daha güvenilir ve kaynakların daha etkin kullanılmasını sağlayan çözümlerin geliştirilmesine katkıda bulunabileceği değerlendirilmektedir.

KAYNAKÇA

Dániel, H., Habashneh, M., & Rad, M. M. (2022). Reliability-based numerical analysis of glulam beams reinforced by CFRP plate. *Scientific Reports*, *12*(1), 13587. https://doi.org/10.1038/s41598-022-17751-6

European Committee for Standardization. (2010). *EN 408:2010+A1:2012: Timber structures—Structural timber and glued laminated timber—Determination of some physical and mechanical properties*. CEN.

European Committee for Standardization. (2013). *EN 14080:2013: Timber structures—Glued laminated timber and glued solid timber—Requirements*. CEN.

Fiorelli, J., & Dias, A. A. (2011). Glulam beams reinforced with FRP externally-bonded: theoretical and experimental evaluation. *Materials and Structures*, *44*(8), 1431-1440. https://doi.org/10.1617/s11527-011-9708-y

Glišović, I., Pavlović, M., Stevanović, B., & Todorović, M. (2017). Numerical analysis of glulam beams reinforced with CFRP plates. *Journal of Civil Engineering and Management*, *23*(7), 868-879. https://doi.org/10.3846/13923730.2017.1341953

Glišović, I., Todorović, M., Simović, N., & Pavlović, M. (2025). Numerical modelling of CFRP internally reinforced glulam beams. *Građevinski materijali i konstrukcije*, *68*(2). https://doi.org/10.5937/GRMK2500002G

Greffier, G., Espinosa, L., Perrin, M., & Eyma, F. (2024). Structural health monitoring of glulam structures: analysis of durability and damage mechanisms. *European Journal of Wood and Wood Products*, *82*(6), 2047-2063. https://doi.org/10.1007/s00107-024-02140-9

İşleyen, Ü. K., Ghoroubi, R., Mercimek, Ö., Anıl, Ö., & Erdem, R. T. (2023, May). Investigation of impact behavior of glulam beam strengthened with CFRP. In *Structures* (Vol. 51, pp. 196-214). Elsevier. https://doi.org/10.1016/j.istruc.2023.03.038

Jaaranen, J., & Fink, G. (2024). A finite element simulation approach for glued-laminated timber beams using continuum-damage model and sequentially linear analysis. *Engineering Structures*, *304*, 117679. https://doi.org/10.1016/j.engstruct.2024.117679

Khelifa, M., Auchet, S., Méausoone, P. J., & Celzard, A. (2015). Finite element analysis of flexural strengthening of timber beams with Carbon Fibre-Reinforced Polymers. *Engineering Structures*, *101*, 364-375. https://doi.org/10.1016/j.engstruct.2015.07.046

Kılıç, İ., Kılıç, K., Şimşek, O., & Altunok, M. (2025). Makine Öğrenmesi Modelleri ile Lamine Ahşap Kirişlerin Eğilme Dayanımı (MOR) Değerlerinin Tahmini. *Bozok Journal of Engineering and Architecture*, (Advanced Online Publication), 31-50. https://doi.org/10.70700/bjea.1816397

Lu, P., Gilbert, B. P., Kumar, C., McGavin, R. L., & Karampour, H. (2024). Comprehensive analysis of glulam delamination through finite element modelling considering heat and mass transfer, plasticity and fracture mechanics: a case study using high density hardwood. *European Journal of Wood and Wood Products*, *82*(5), 1581-1604. https://doi.org/10.1007/s00107-024-02107-w

Muhaxheri, M., Krasniqi, E., Kabashi, N., Murati, Y., & Mahmuti, R. (2026). Experimental and numerical assessment of flexural behavior of CFRP–strengthened timber beams. *Polymers*, *18*(1), 134. https://doi.org/10.3390/polym18010134

Nowak, T. P., Jasieńko, J., & Czepiżak, D. (2013). Experimental tests and numerical analysis of historic bent timber elements reinforced with CFRP strips. *Construction and Building Materials*, *40*, 197-206. https://doi.org/10.1016/j.conbuildmat.2012.09.106

Paroissien, J., Oudjene, M., & Lardeur, P. (2024). Verification and Validation of finite element models for laminated timber structures using solid, solid-beam and solid-shell approaches. *Composite Structures,* *345*, 118083. https://doi.org/10.1016/j.compstruct.2024.118083

Raftery, G. M., & Harte, A. M. (2011). Low-grade glued laminated timber reinforced with FRP plate. *Composites Part B: Engineering*, *42*(4), 724-735.

Raftery, G. M., & Harte, A. M. (2013). Nonlinear numerical modelling of FRP reinforced glued laminated timber. *Composites Part B: Engineering*, *52*, 40-50. https://doi.org/10.1016/j.compositesb.2013.03.038

Szczecina, M. (2023). Study of complexity of numerical models of a strengthened timber beam. *Materials*, *16*(9), 3466. https://doi.org/10.3390/ma16093466

Türker, Y. Ş., & Kılınçarslan, Ş. (2024). Effect of Layers Number on The Bending Properties of Chestnut Glulam Beams. *Karadeniz Fen Bilimleri Dergisi*, *14*(1), 62-71. https://doi.org/10.31466/kfbd.1347435

Vafadar, F., Jaaranen, J., & Fink, G. (2025). Probabilistic numerical modeling of glued laminated timber beams—Experimental validation and insights into the potential of layup modification. *Engineering Structures*, *344*, 121310. https://doi.org/10.1016/j.engstruct.2025.121310

Vida, C., Lukacevic, M., Hochreiner, G., Pech, S., & Füssl, J. (2025). Numerical modeling of glued laminated timber beams without finger joints: Identifying load-bearing capacity and analyzing failure mechanisms. *Engineering Structures*, *345*, 121489. https://doi.org/10.1016/j.engstruct.2025.121489

Wdowiak-Postulak, A. K., Świt, G., Krampikowska, A., & Minh Chinh, L. (2025). Experimental and Numerical Studies of "Wood–Composite" Reinforcement in Bending

Sheared Wooden Beams Using Pre-Stressed Natural and Artificial Fibers. *Materials*, *18*(18), 4418. https://doi.org/10.3390/ma18184418

Yang, H., Liu, W., Lu, W., Zhu, S., & Geng, Q. (2016). Flexural behavior of FRP and steel reinforced glulam beams: Experimental and theoretical evaluation. *Construction and building materials*, *106*, 550-563. https://doi.org/10.1016/j.conbuildmat.2015.12.135

Younis, H., Al Abadi, H., Patel, V. I., & Naga, H. A. (2025). Finite element analysis of fibre reinforced timber beams under flexural loading. *Australian Journal of Structural Engineering*, *26*(2), 65-85. https://doi.org/10.1080/13287982.2024.2346955

Zhang, X., Chen, Z., Niu, B., Li, Z., & Zhou, X. (2026, February). Study on bending behavior of cross-laminated timber-glulam composite beam: Numerical simulation and theoretical analysis. In *Structures* (Vol. 84, p. 110932). Elsevier.

THE ROLE OF MATERIAL SELECTION IN LIGHTWEIGHTING ELECTRIC VEHICLES

Assoc. Prof. Dr. Omer CIHAN[1]
[1] Sakarya University of Applied Sciences, Hendek Vocational School, Department of Machinery and Metal Technologies
[1] Sakarya University of Applied Sciences, Automotive Technologies Application and Research Center
[1]ORCID ID: https://orcid.org/0000-0001-8103-3063

Assoc. Prof. Dr. Mustafa Guven GOK[2]
[2] Gaziantep University, Department of Metallurgical and Materials Engineering
[2]ORCID ID: https://orcid.org/0000-0002-5959-0549

ABSTRACT

As the electrification of the transportation sector gains momentum in line with sustainable mobility goals, the relationship between the total weight of electric vehicles (EVs) and their energy efficiency has become one of the most important engineering challenges. In these vehicles, which are 20-30% heavier than internal combustion engine vehicles due to battery systems, lightweighting strategies offer direct advantages such as increased range, improved driving dynamics, and shortened braking distances. This review study examines the important role of material selection in weight reduction in electric vehicles. The microstructural advantages of advanced high-strength steels (AHSS), specific strength performances of aluminum and magnesium alloys, and stiffness contributions of carbon fiber reinforced polymer (CFRP) composites are analyzed. The effects of material substitution in battery enclosures on mass reduction, impact absorption, electromagnetic shielding, and thermal management are discussed. Bio-composites and recyclable polymers are evaluated from an environmental sustainability perspective.

Keywords: Alternative materials; Material selection; Battery enclosure material; Lightweighting; Electric vehicles.

* Corresponding author: omercihan@subu.edu.tr (O. Cihan)

1. INTRODUCTION

Electric vehicles can be generally defined as vehicles that provide motion energy from electricity. It is predicted that CO_2 exhaust emission values, which are the biggest environmental disadvantage of internal combustion engines, can be reduced with electric vehicles (Larminie & Lowry, 2003). In addition to being more environmentally friendly, studies are being carried out intensively to make electric vehicles more economical and more effective. Today, many companies in the automotive sector continue their research and development activities without slowing down (Chan, 2001). The zero-emission policy, which is accepted worldwide, is increasing the demand for electric vehicles on a global scale in order to minimize environmental negativity and reduce energy demand. According to a report by the International Energy Agency (IEA), 145 million vehicles are targeted for 2030 (IEA, 2019; IEA, 2022; Zhou et al., 2022). For 2024, 17 million electric vehicle sales and a market share of over 20% are

projected. Based on current trends and policies, it is expected that 6 million barrels of oil need per day will be eliminated by 2030 with the spread of electric vehicles (IEA, 2022).

For electric vehicles to be developed and become demanded, material selection in each part is very important. Material selection in electric vehicles plays a critical role in terms of vehicle performance, efficiency, safety and sustainability (Lewchalermwong et al., 2017). EVs carry heavier battery systems than internal combustion engine vehicles; this can increase energy consumption. Therefore, lightweight but durable materials should be preferred in order to reduce the total weight of the vehicle, increase range and increase energy efficiency (Poulikidou et al., 2015). Advanced materials such as aluminum, magnesium alloys, high-strength steels and carbon fiber respond to this need. However, each material should be carefully evaluated in terms of production cost, processability and environmental impact (Kalhor et al., 2025).

Material selection directly affects not only weight but also factors such as safety, production process and recyclability. For example, criteria such as impact absorbing properties in the event of a collision or fire resistance determine whether the material can be used in the vehicle body and battery protection systems. In addition, in line with sustainable production goals, the tendency towards recyclable or biodegradable materials is increasing. Therefore, material selection is a multi-dimensional process that shapes not only technical but also economic and environmental decisions in EV production (Hawkins et al., 2012). Material selection and technology has been a major limiting factor in the development of electric vehicles (EVs). These vehicles need to achieve longer range, be safe and durable, while remaining lightweight, but these requirements create conflicting engineering goals. Heavy materials such as traditional steel increase energy consumption, while lightweight and durable alternatives (e.g. carbon fiber, aluminum alloys, magnesium and composites) are still costly and complex to manufacture (Sun et al., 2020). In addition, the weight of battery systems often overshadows these lightweighting efforts (Haghani et al., 2023).

On the other hand, batteries, the cores of EVs, consist of complex aluminum channels for their covers and battery packs (Shui et al., 2018). Nowadays, it is very difficult to obtain thin-walled aluminum products with complex geometry due to high springback and low formability. To overcome these challenges, new forming techniques such as hot forming and quenching (HFQ®), incremental sheet forming (ISF) and transverse stretching and local bending (TSLB) are used (Castelvecchi, 2021; Zhou et al., 2016a; Zhou et al., 2016b). Although aluminum is difficult to produce at low cost and recyclable, it is overcome by techniques such as solid state recycling (Shamsudin et al., 2016; Wan et al., 2017). One of the solutions to these problems is the use of sustainable composites. Generally a mixture of fiber (such as glass or carbon) or matrix (e.g. resin), these materials optimize the power-to-weight ratio (Wieckowski & Salabun, 2025). Sustainable composites can transform waste materials into valuable resources (Phiri et al., 2023). Thus, they can be recycled and make a major contribution to life cycle impact (LCA) (Cimprich et al., 2023).

Weight reduction in electric vehicles is crucial to increase the vehicle's range and improve energy efficiency. Lighter and more durable alternatives to traditional steel materials are preferred. Lightweight metals and composite materials such as aluminum, carbon fiber and magnesium alloys are frequently used in electric vehicle production (Kong, 2025). Aluminum is about one-third the weight of steel but offers high durability and is widely used in both chassis and body panels. Carbon fiber is an extremely light and strong material, and is especially preferred for high-performance electric vehicles. By using aluminum alloys, polymer

composites, carbon fiber and magnesium alloys instead of traditional materials such as steel and cast iron, the body and chassis weight of the vehicle can be reduced by up to 50% (Singh, 2024).

In addition, lightweight materials are also preferred for the car's interior structure and battery housings. For example, magnesium alloys are used in steering systems, seat structures and various interior skeleton parts, while thermoplastic composites offer the advantage of both light weight and easy formability (Koech et al., 2024). The use of these materials not only reduces the weight of the vehicle but also improves its performance and contributes to more efficient use of battery capacity (Jimenez-Martinez et al., 2024). Electric vehicles are of great importance today as they enable environmentally friendly and sustainable transportation by reducing dependence on fossil fuels. They minimize carbon emissions, provide energy efficiency and offer long-term economic advantages. In this review study, the materials used in the electric vehicle industry and the importance of material selection are analyzed.

2. ELECTRIC VEHICLES

Electric vehicles are vehicles that use electric engines instead of internal combustion engines to provide motion energy. These vehicles take energy from rechargeable batteries and offer a more environmentally friendly, quiet and energy efficient transport option because they run on electricity instead of fossil fuels (Costa et al., 2021). It is basically based on the principle of using the energy stored in the battery to drive electric motors. Today, it is available as fully electric or hybrid. Fig. 1 shows the schematic of the electric vehicle.



Fig. 1. Basic elements of the movement system in an electric vehicle (Guven & Rende, 2017).

The basic components in electric vehicles complement each other to ensure efficient and safe operation of the system. The battery pack is the main energy source of the vehicle and feeds the electric motor. The charging unit enables the vehicle to safely charge the battery by receiving electricity from external sources. The main contactor controls the high-voltage circuits to start or stop the flow of energy from the battery to the engine. The control unit acts as the brain of the entire system, managing the engine, battery and charging processes and optimising the vehicle's performance. The electric motor moves the vehicle by converting the energy from the battery into mechanical energy. The auxiliary battery supplies low-voltage systems such as headlights, displays and air conditioning. Finally, the converter converts high-voltage energy into low-voltage energy to ensure the safe operation of the auxiliary systems (Acar et al., 2024; Tanvir & Harun-ur Rashid, 2024).

Electric vehicles (EVs) are characterised by their environmental friendliness because they reduce air pollution as they do not emit exhaust gases. They also have fewer moving parts than

internal combustion engines, so they have low maintenance costs (Louback et al., 2024). Thanks to their silent operation, they do not create noise pollution and their energy efficiency is quite high; they can cover the same distance with less energy. Since electric motors can produce torque instantly, they provide fast acceleration (Kharabati & Saedodin, 2024).

The biggest disadvantages of electric vehicles are their shorter range compared to conventional vehicles and limitations on battery life. The fact that the charging infrastructure is not sufficiently developed in every region may necessitate planning for long journeys. Charging times are longer than refuelling of petrol vehicles. Also, the initial purchase cost of electric vehicles is generally higher. The environmental impacts of materials such as lithium and cobalt used in the production of batteries and ethical procurement problems can also be counted among the important disadvantages (Shao et al., 2024).

When the studies are analysed, it is predicted that the annual sales of electric vehicles will increase in the coming years (Li et al., 2023; Wu & Chen, 2022). In the Wu and Chen study (2022), both machine learning and statistical methods were used to estimate the annual sales of electric vehicles worldwide from 2021 to 2028 (Fig. 2a). As can be seen from here, the sales of electric vehicles have increased continuously from year to year. Strict emission rules, more environmentally friendly and the gradual depletion of oil reserves have increased the rate of transition to electric vehicles. Fig. 2b shows that publications, scientific studies and research and development activities on electric vehicles have increased significantly in recent years.

According to the worldwide research on electric vehicle sales, the fact that electric vehicles can be produced at low cost has caused EVs to break a record in sales. In 2024, sales exceeded 17 million worldwide as expected. As predicted by the IEA, electric vehicles increased their market share to over 20%. In the first quarter of 2025, electric vehicle sales increased by 35% compared to the same period of 2024. China is the leader among countries, accounting for approximately half (more than 11 million) of vehicle sales in 2024. Electric vehicle sales in Asia and Latin America increased by 60% in 2024 (IEA, 2025).



Fig. 2. Predicted number of sales of electric vehicles from 2021 to 2028 (a), the number of publications on EVs from the beginning of the 21st century to 2023 based on the Scopus database (b) (Kalhor et al., 2025; Wu & Chen, 2022).

3. MATERIALS USED IN ELECTRIC VEHICLES AND NEW GENERATION MATERIALS

In the electric vehicle sector, "biocomposite" refers to a more environmentally friendly alternative to traditional composites, created by combining renewable or biologically sourced fibers with a polymer (plastic) matrix.

Recently, there has been increased discussion about the need to reduce the environmental impact of vehicles. The goal is to reduce the environmental burden not only during use, but also during the production process and the disposal phase after the end of its life. As this awareness increases, the use of biocomposites in electric vehicles is accelerating. These sustainable materials, which combine natural fibers with a polymer matrix, are being used in more and more EVs (Yorseng et al., 2019). There are several clear reasons why biocomposites are attractive: their lower environmental impact, their good strength, and their ability to perform the same task with lower weight. Therefore, among the types of biocomposites being evaluated for electric vehicles, natural fiber-reinforced composites, wood-polymer composites, cellulose/lignin-based composites, bio-resin composites, and recycled composites stand out (Singh et al., 2025).

Biocomposites are produced in a fairly simple manner. Natural fibers (such as flax and hemp) are mixed with a polymer "matrix" carrier, resulting in a material that is both lightweight and strong. Moreover, these materials are attracting considerable interest because they have a lower environmental impact (Naik et al., 2022). Wood-plastic composites are produced by combining

wood fiber or wood flour with polymer. They are quite advantageous in places such as interior trim parts, trim details, and panel applications (Mohan Kumar et al., 2023). There are also composites derived from cellulose fibers. In addition to being durable, their light weight is a major advantage. Lignin, a by-product of the paper industry, also plays a role in this field. Composites developed using lignin are considered "cleaner" from an environmental perspective (Firmanda et al., 2022). Another approach is to create the matrix using bio-based resins obtained from plants or renewable sources instead of petroleum-based polymers. When these types of resins are combined with natural fibers, the result is a completely sustainable biocomposite (Stewart, 2010; Verrey et al., 2006).

Research on the use of renewable and biodegradable materials in electric vehicles is also increasing. For example, in non-structural parts, it is possible to produce a solution called "green composite" by combining natural fibers with a starch-based matrix (Fais, 2011). Additionally, there is the issue of recycled composites, which play a key role in the environmental aspect of the industry. The use of recycled composites reduces waste, lowers the carbon footprint, and curbs raw material consumption (Tewari et al., 2021).

In the automotive industry, polymer technologies are gradually replacing metal parts, directly contributing to both reducing vehicle weight and opening up greater design flexibility. The study by Gupta et al. (2022) examines the use of polymers in the electric vehicle (EV) market within a broad framework ranging from powertrain components (battery parts, motor housings, etc.) to interior and exterior trim applications. The study particularly highlights the critical function of polymer-based binders and separators used in lithium-ion batteries. The potential of bio-based polymers and nanocomposites is evaluated from a sustainability perspective. According to the assessment presented in Fig. 3, the global polymer market used in electric vehicles stood at US$3.8 billion in 2020 and is projected to reach US$27.3 billion in 2026, with a compound annual growth rate (CAGR) of 38.7% during the 2020-2026 period (Gupta et al., 2022).

Fig. 3. Annual compound growth rates of global markets between 2020 and 2026 (%) (Gupta et al., 2022).

3.1 Advanced Metal Technologies and Hybrid Structures

The rapid increase in the market share of electric vehicles (EVs) has made "lightweighting" strategies the most critical way to increase energy efficiency, compensating for limited battery capacity and the burden of heavy battery packs (Delogu et al., 2017; Nadikudi, 2026). In this context, the automotive industry is witnessing a competition in materials that go beyond traditional steels, balancing cost and performance; in particular, "Advanced High-Strength Steels" (AHSS) and "Medium-Mn" steels, offer the possibility of thinning without compromising safety thanks to the combination of high strength and ductility they provide

through TRIP and TWIP effects (Kozlowska & Wojtacha, 2025; Mohapatra et al., 2025; Zhang et al., 2024), while aluminum and magnesium alloys provide significant mass savings in chassis, body, and interior parts due to their low densities (Liu et al., 2023; Zhan et al., 2025). However, the inability of a single material to meet all requirements (cost, strength, lightness) simultaneously has made "Hybrid Structures" (multi-material designs), where different materials are strategically combined, an emerging trend (Delogu et al., 2017; Li et al., 2018). Nevertheless, the success of these hybrid structures depends on overcoming serious metallurgical challenges, particularly in the combination of different metals such as Aluminum-Steel or Steel-Copper in battery packs, due to brittle "Intermetallic Compounds" (IMC) and corrosion resulting from electrochemical potential differences (Sadeghian & Iqbal, 2022; Wan & Huang, 2018; Yang et al., 2024).

Fig. 4 summarizes the "banana plot" illustrating the inverse relationship between strength and ductility in automotive steels and the intergenerational evolution in material technology. The graph highlights the significant performance gap between 1st Generation AHSS with limited mechanical properties and 2nd Generation AHSS with high production costs. Developed to bridge this gap, the 3rd Generation AHSS class of Medium-Manganese steels (3-12% Mn) offer a superior strength-ductility balance thanks to the TRIP and TWIP mechanisms they incorporate. Positioned in the "High Growth Potential" category, these new-generation materials aim to cost-effectively meet the weight reduction and crash safety requirements of electric vehicles (Zhang et al., 2024).



Fig. 4. A comparative overview of the ultimate tensile strength (UTS) and total elongation (TEL) values for various commercially available steel grades (Zhang et al., 2024).

Looking ahead, the evolution of material technologies in electric vehicles will be shaped not only by the discovery of new alloys, but also by the digitalization and sustainability of the production and joining processes of these materials. Zhang et al. (2024) predict that the use of "Material Genome Engineering" and machine learning techniques in the composition design of medium-manganese steels will be a critical route to achieving an optimal balance between lightness and mechanical performance. In parallel, Zhan et al. (2025) and Nadikudi (2026) state that "Additive Manufacturing" methods such as Laser Powder Bed Fusion (LPBF) and 3D printing in the production of magnesium alloys and composites will increase design freedom by enabling the production of complex geometries. In joining technologies, Sadeghian and Iqbal (2022) emphasize that the use of blue and green wavelength lasers, which increase energy absorption, will become widespread, particularly in the welding of highly reflective metals such as copper and aluminum in battery modules. However, as pointed out by Wan and Huang (2018) and Yang et al. (2024), for the long-term reliability of all these hybrid structures, modeling the evolution of intermetallic compounds formed at joining interfaces under thermal and

mechanical fatigue throughout their service life and integrating them with "green" production (Liu et al., 2023) techniques will be the ultimate thresholds that the industry must overcome.

3.2 Battery Enclosure Materials

Battery enclosures, which directly affect the performance and safety of electric vehicles (EVs), protect cells from mechanical impacts and vibration, and are also a critical component of the thermal management system. While steel alloys such as DC01 and Q235 are used in traditional designs due to their cost and strength advantages, range concerns and lightweighting requirements have led the industry to aluminum alloys such as Al6061 (Chen et al., 2019; Chidambaram et al., 2025). Although aluminum offers better thermal conductivity and weight balance compared to steel, current research focuses on composite materials such as Carbon Fiber Reinforced Polymers (CFRP) and glass fiber, which can reduce weight by over 40% (Chen et al., 2019; Dhoke & Dalavi, 2021; Liu et al., 2024). However, composites have lower thermal conductivity than metallic enclosures, making it difficult to dissipate battery-generated heat and complicating the management of thermal runaway risk (Nieduzak et al., 2025; Samarasinghe, 2024). To overcome these thermal constraints, innovative designs with variable thermal conductivity, incorporating copper pins into the composite matrix, are proposed, thereby creating passive cooling channels in critical areas (Samarasinghe, 2024). Furthermore, to maximize crashworthiness, bio-inspired cellular structures and multi-material hybrid structures combining Aluminum Honeycomb (AAHP) with CFRP demonstrate superior performance in protecting battery integrity by increasing energy dissipation capacity (Dabasa et al., 2025; Liu et al., 2024). Consequently, material selection requires a multidisciplinary optimization process that aims for the optimal balance of mechanical strength, lightness, fire resistance, and thermal management (Kim et al., 2019; Nummy, 2022). Fig. 5 shows the battery housing components (top cover, bottom body, brackets, etc.) and the assembly structure.



Fig. 5. Structure of the battery enclosure (Pan et al., 2021).

Table 1. Comparative literature review: Battery enclosure materials and design

Study (Author, Year)	Material / Structure Investigated	Focused Issue	Key Findings / Result
Yousefi et al., 2023	Copper (Cu), Aluminum (Al), Acrylic Enclosures + Nano-PCM	Thermal management and heat dissipation	The copper enclosure exhibited the lowest battery surface temperature; CuO nanoparticles significantly enhanced heat transfer.
Samarasinghe, 2024	Variable Thermal Conductivity Composite (CFRP + Copper Z-pins)	Passive cooling and lightweighting	Integrating copper pins (Z-pinning) into the composite matrix created effective thermal pathways; simulation results showed strong correlation with experimental data.
Dhoke & Dalavi, 2021	Multi-Materials (HSS, CFRP, Aluminum) and Honeycomb Structures	Lightweighting and crash safety	Replacing steel with composite and aluminum hybrid structures can achieve weight reductions exceeding 40% while enhancing crash performance.

Study (Author, Year)	Material / Structure Investigated	Focused Issue	Key Findings / Result
Pan et al., 2021	Advanced High Strength Steel (AHSS - 780DP)	Vibration, shock resistance, and fatigue life	Through size optimization substituting DC01 steel with AHSS, vibration and shock requirements were met while achieving a total weight reduction of 10.41%.
Shui et al., 2018	Standard Battery Enclosure (Structural Optimization)	Mass minimization and deformation control	Optimization using CCD and Artificial Neural Networks (ANN) resulted in an 11.61% mass reduction and 22.22% decrease in maximum deformation.
Dabasa et al., 2025	Bio-inspired Cellular Structures (Honeycomb, Lattice, Auxetic)	Crashworthiness and energy absorption	Bio-inspired designs (e.g., super hexagonal honeycombs) demonstrated Specific Energy Absorption (SEA) increase of over 100% compared to traditional structures.
Dai & Panahi, 2025	Cathode Materials and Pack Protection Strategies	Thermal Runaway mechanisms	A comprehensive review of thermal stability limits for various cell chemistries (LCO, LFP, NCM) and the critical role of protective layers.
Wazeer et al., 2022	Phase Change Materials (PCM) + Metal Foam/Graphite Matrix	Thermal homogeneity and passive cooling	The low thermal conductivity of pure PCM was enhanced by up to 44 times through porous copper foam or graphite matrices.
Ganguli & Date, 2024	Metallic Enclosure (Parametric and Topology Optimization)	Weight reduction and thermal feasibility	Topology optimization reduced the enclosure weight to 1.48 kg; air cooling channels maintained cell temperatures within the safe range (<47°C).
Chidambaram et al., 2025	Aluminum Alloy (AL6061) vs. Steel (Q235, DC01) and Teflon	Structural integrity and thermal properties	AL6061 was identified as the optimal material, offering a superior balance of weight, thermal conductivity, and structural stability.
Nummy, 2022	Thermoplastics and Thermosets (Glass Fiber Reinforced)	Fire resistance and metal-to-plastic transition	With appropriate sealing and venting designs, plastic composites can meet fire safety requirements during thermal runaway events.
Nieduzak et al., 2025	Fiber Reinforced Polymer (FRP) + Embedded Sensors	Weight reduction and temperature monitoring	Embedding sensors in composites enables lightweight smart structures; a distinct thermal lag in detecting internal temperature changes was characterized.
Liu et al., 2024	AAHP (Al Honeycomb) / CFRP Hybrid Structure	Lightweighting via Digital Twin technology	The hybrid design of Aluminum Honeycomb and CFRP reduced weight by 12.3% and increased the first-order bending frequency by 27.5%.
Kim et al., 2019	PCM Composites, Heat Pipes, and Liquid Cooling Plates	Effectiveness of BTMS (Battery Thermal Management Systems)	PCM/Graphite composites provided superior surface temperature reduction and thermal uniformity compared to conventional air cooling.
Chen et al., 2019	CFRP (Body) and Aluminum Alloy (Frame)	Multi-material design and structural strength	The hybrid CFRP/Aluminum design met lightweighting targets with maximum stress (285.7 MPa) remaining below the yield limit.

4. CONCLUSIONS

The global adoption of electric vehicles (EVs) depends not only on advancements in battery technologies but also on the selection of appropriate materials that shape the total weight and structural integrity of the vehicle. The analyses conducted in this review demonstrate that the era of "single-material design" in the automotive industry has ended, replaced by multi-material hybrid structures that combine the advantages of different material groups.

The key findings derived from this study can be summarized as follows:

- **Weight and range relationship:** It has been observed that the use of aluminum and advanced composites can save up to 50% in chassis and body weight to compensate for the negative effects caused by battery mass. This reduction translates directly into extended driving range and lower energy consumption.
- **Material diversity and battery safety:** Moving away from traditional steel in battery enclosures provides lightness but complicates critical safety parameters such as thermal management and fire resistance. Hybrid designs and bio-inspired cellular structures are the strongest candidates, offering both lightweighting and high energy absorption capacity.
- **Sustainability-oriented approach:** The future of the sector is evolving toward bio-composites and recycled polymer technologies that remain "green" not only during the usage phase but also throughout the production and recycling processes (Life Cycle Assessment).
- **Technological transformation:** Digital methods such as additive manufacturing (3D printing) and material genome engineering enable the production of complex geometries, maximizing design flexibility and material efficiency.

In conclusion, material selection in the electric vehicle industry is no longer merely an engineering decision; it is a multi-dimensional optimization process where economic, technical, and environmental variables are evaluated together. Future studies focusing on overcoming the metallurgical challenges of joining dissimilar materials and reducing production costs are essential for the sustainable growth of the EV market.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest in this study.

REFERENCES

Acar, E., Jain, N., Ramu, P., Hwang, C., & Lee, I. (2024). A survey on design optimization of battery electric vehicle components, systems, and management. Structural and Multidisciplinary Optimization, 67 (27). https://doi.org/10.1007/s00158-024-03737-7.

Castelvecchi, D. (2021). Electric cars and batteries: how will the world produce enough?. Nature, 596, 336-339. https://doi.org/10.1038/d41586-021-02222-1.

Chan, C. C. (2001). Modern Electric Vehicle Technology. Oxford University Press, New York, USA.

Chen, X., Li, M., Li, S., Jin, J., & Zhang, C. (2019). Design optimization of multi-material battery enclosure for electric vehicle. Lecture Notes in Electrical Engineering, 486, 853-864. https://doi.org/10.1007/978-981-10-8506-2_57.

Chidambaram, R. K., Pedapati, P. R., Kanna, P. R., Taler, D., Sobota, T., & Taler, J. (2025). Structural assessment of electric two-wheeler battery enclosure: thermal and structural study. Journal of Thermal Analysis and Calorimetry, 150, 6939-6958. https://doi.org/10.1007/s10973-024-13458-0.

Cimprich, A., Sadayappan, K., & Young, S. B. (2023). Lightweighting electric vehicles: Scoping review of life cycle assessments. Journal of Cleaner Production, 433, 1-10. https://doi.org/10.1016/j.jclepro.2023.139692.

Costa, C. M., Barbosa, J. C., Castro, H., Goncalves, R., & Lanceros-Mendez, S. (2021). Electric vehicles: To what extent are environmentally friendly and cost effective? Comparative study by European countries. Renewable and Sustainable Energy Reviews, 151, 1-16. https://doi.org/10.1016/j.rser.2021.111548.

Dabasa, T., Lemu, H. G., & Regassa, Y. (2025). A review on the crashworthiness of bio-inspired cellular structures for electric vehicle battery pack protection. Computation, 13 (9), 217. https://doi.org/10.3390/computation13090217.

Dai, Y., & Panahi, A. (2025). Thermal runaway process in lithium-ion batteries: A review. Next Energy, 6, 100186. https://doi.org/10.1016/j.nxener.2024.100186.

Delogu, M., Zanchi, L., Dattilo, C. A., & Pierini, M. (2017). Innovative composites and hybrid materials for electric vehicles lightweight design in a sustainability perspective. Materials Today Communications, 13, 192-209. https://doi.org/10.1016/j.mtcomm.2017.09.012.

Dhoke, A., & Dalavi, A. (2021). Lightweight design of battery pack enclosure for electric vehicle using multi-materials and honeycomb structures. Materials Today: Proceedings. https://doi.org/10.1016/j.matpr.2021.06.388.

Fais, C. (2011). Lightweight automotive design with HP-RTM. Reinforced Plastics, 55 (5), 29-31. https://doi.org/10.1016/S0034-3617(11)70142-4.

Firmanda, A., Syamsu, K., Sari, Y. W., Cabral, J., Pletzer, D., Mahadik, B., Fisher, J., & Fahma, F. (2022). 3D printed cellulose based product applications. Materials Chemistry Frontiers, 6 (3), 254-279. https://doi.org/10.1039/D1QM00390A.

Ganguli, T., & Date, P. P. (2024). Lightweight battery enclosure design. Materials Research Proceedings, 44, 465-474. https://doi.org/10.21741/9781644903254-50.

Gupta, P., Toksha, B., Patel, B., Rushiya, Y., Das, P., & Rahaman, M. (2022). Recent developments and research avenues for polymers in electric vehicles. The Chemical Record, 22 (11), 1-40. https://doi.org/10.1002/tcr.202200186.

Guven, F., & Rende, H. (2017). Importance of material selection in design of electric vehicles. Engineer and Machinery, 58 (689), 81-95.

Haghani, M., Sprei, F., Kazemzadeh, K., Shahhoseini, Z., & Aghaei, J. (2023). Trends in electric vehicles research. Transportation Research Part D: Transport and Environment, 123, 1-29. https://doi.org/10.1016/j.trd.2023.103881.

Hawkins, T. R., Gausen, O. M., & Stromman, A. H. (2012). Environmental impacts of hybrid and electric vehicles: A review. International Journal of Life Cycle Assessment, 17, 997-1014. https://doi.org/10.1007/s11367-012-0440-9.

IEA (2019). Sustainable development scenario.

IEA (2022). Global EV outlook 2022: securing supplies for an electric future. https://www.iea.org/energy-system/transport/electric-vehicles, Accessed 1 Feb 2026.

IEA (2025). More than 1 in 4 cars sold worldwide this year is set to be electric as EV sales continue to grow. https://www.iea.org/news, Accessed 8 May 2025.

Jimenez-Martinez, M., Valencia-Sanchez, J. L., Torres-Cedillo, S. G., & Cortes-Perez, J. (2024). Battery housing for electric vehicles, a durability assessment review. Designs, 8 (6), 1-20. https://doi.org/10.3390/designs8060113.

Kalhor, A., Dykas, J., Rodak, K., & Grajcar, A. (2025). Materials and constructional design for electric vehicles: A review. Advances in Science and Technology Research Journal, 19 (1), 178-196. https://doi.org/10.12913/22998624/195457.

Kharabati, S., & Saedodin, S. (2024). A systematic review of thermal management techniques for electric vehicle batteries. Journal of Energy Storage, 75, 1-58. https://doi.org/10.1016/j.est.2023.109586.

Kim, J., Oh, J., & Lee, H. (2019). Review on battery thermal management system for electric vehicles. Applied Thermal Engineering, 149, 192-212. https://doi.org/10.1016/j.applthermaleng.2018.12.020.

Koech, A. K., Mwandila, G., & Mulolani, F. (2024). A review of improvements on electric vehicle battery. Heliyon, 10 (15), 1-23. https://doi.org/10.1016/j.heliyon.2024.e34806.

Kong, W. (2025). Researches on lightweight body structure design and optimization of electric vehicles. Applied and Computational Engineering, 124, 183-189. https://doi.org/10.54254/2755-2721/2025.20103.

Kozlowska, A., & Wojtacha, A. (2025). Advanced high-strength medium-manganese steels as an alternative to conventional forging steels: A review. Materials, 19 (1), 1-27. https://doi.org/10.3390/ma19010109.

Larminie, J., & Lowry, J. (2003). Electric Vehicle Technology Explained. John Wiley & Sons Ltd, Oxford, UK.

Lewchalermwong, N., Masomtob, M., Lailuck, V., & Charoenphonphanich, C. (2017). Material selection and assembly method of battery pack for compact electric vehicle. IOP Conference Series: Materials Science and Engineering, 297.

Li, H., Liu, X. S., Zhang, Y. S., Ma, M. T., Li, G. Y., & Senkara, J. (2018). Current research and challenges in innovative technology of joining dissimilar materials for electric vehicles. Advanced High Strength Steel and Press Hardening, Book Chapter, 363-380. https://doi.org/10.1142/9789813277984_0056.

Li, Y., Liang, C., Ye, F., & Zhao, X. (2023). Designing government subsidy schemes to promote the electric vehicle industry: A system dynamics model perspective. Transportation Research Part A: Policy and Practice, 167, 1-14. https://doi.org/10.1016/j.tra.2022.11.018.

Liu, B., Yang, J., Zhang, X., Yang, Q., Zhang, J., & Li, X. (2023). Development and application of magnesium alloy parts for automotive OEMs: A review. Journal of Magnesium and Alloys, 11 (1), 15-47. https://doi.org/10.1016/j.jma.2022.12.015.

Liu, J., Xu, L., Yan, M., Zhang, H., & Deng, H. (2024). Digital twin-based AAHP/CFRP battery enclosure hybrid optimization lightweight design method. ASME Journal of Risk and Uncertainty in Engineering Systems, Part B: Mechanical Engineering. https://doi.org/10.1115/1.4069678.

Louback, E., Biswas, A., Machado, F., & Emadi, A. (2024). A review of the design process of energy management systems for dual-motor battery electric vehicles. Renewable and Sustainable Energy Reviews, 193, 1-11. https://doi.org/10.1016/j.rser.2024.114293.

Mohan Kumar, K., Naik, V., Kaup, V., Waddar, S., Santhosh, N., & Harish, H. V. (2023). Nontraditional natural filler-based biocomposites for sustainable structures. Advances in Polymer Technology, 2023 (1), 1-15. https://doi.org/10.1155/2023/8838766.

Mohapatra, S., Satpathy, B., Sarkar, A., Kumar, S., Aich, S., & Oh, M. S. (2025). Recent advances in medium-manganese steel: A review of key findings, challenges, and future perspectives. Journal of Materials Science, 60, 9447-9528. https://doi.org/10.1007/s10853-025-11010-5.

Nadikudi, B. K. B. (2026). A summarized analysis of emerging light weight composite materials for electric vehicles. Turkish Journal of Engineering, 10 (1), 178-186. https://doi.org/10.31127/tuje.1710658.

Naik, V., Kumar, M., & Kaup, V. (2022). A review on natural fiber composite materials in automotive applications. Engineered Science, 18, 1-10. https://doi.org/10.30919/es8d589.

Nieduzak, T. B., Tronci, E. M., Zhou, T., Demo, L. B., Feng, M. Q., & Aitharaju, V. (2025). Internal temperature sensing of a composite battery enclosure during thermal shock. Composites Part B: Engineering, 296, 112258. https://doi.org/10.1016/j.compositesb.2025.112258.

Nummy, A. (2022). Development of a new flammability test method: enabling material-level evaluation of polymeric materials for electric vehicle battery enclosures. SAE Technical Paper 2022-01-0714. https://doi.org/10.4271/2022-01-0714.

Pan, Y., Xiong, Y., Wu, L., Diao, K., & Guo, W. (2021). Lightweight design of an automotive battery-pack enclosure via advanced high-strength steels and size optimization. International Journal of Automotive Technology, 22 (5), 1279-1289. https://doi.org/10.1007/s12239-021-0112-5.

Phiri, R., Rangappa, S. M., Siengchin, S., & Marinkovic, D. (2023). Agro-waste natural fiber sample preparation techniques for bio-composites development: methodological insights. Facta Universitatis-Series Mechanical Engineering, 21 (4), 631-656. https://doi.org/10.22190/FUME230905046P.

Poulikidou, S., Schneider, C., Bjorklund, A., Kazemahvazi, S., Wennhage, P., & Zenkert, D. (2015). A material selection approach to evaluate material substitution for minimizing the life cycle environmental impact of vehicles. Materials & Design, 83, 704-712. https://doi.org/10.1016/j.matdes.2015.06.079.

Sadeghian, A., & Iqbal, N. (2022). A review on dissimilar laser welding of steel-copper, steel-aluminum, aluminum-copper, and steel-nickel for electric vehicle battery manufacturing. Optics & Laser Technology, 146, 1-24. https://doi.org/10.1016/j.optlastec.2021.107595.

Samarasinghe, D. A. (2024). Composite enclosure for Li-ion battery module. Ph.D. Thesis, Brunel University London, UK.

Shamsudin, S., Lajis, M. A., & Zhong, Z. W. (2016). Solid-state recycling of light metals: a review. Advances in Mechanical Engineering, 8 (8), 1-23. https://doi.org/10.1177/1687814016661921.

Shao, D., Hu, L., Zhang, J., Hu, R., Zhang, G., Jiang, L., Wang, X., & Wen, Y. (2024). Advanced low-temperature preheating strategies for power lithium-ion batteries applied in electric vehicles: A review. International Journal of Electrochemical Science, 19 (11), 1-24. https://doi.org/10.1016/j.ijoes.2024.100817.

Shui, L., Chen, F. Y., Garg, A., Peng, X., Bao, N., & Zhang, J. (2018). Design optimization of battery pack enclosure for electric vehicle. Structural and Multidisciplinary Optimization, 58 (1), 331-347. https://doi.org/10.1007/s00158-018-1901-y.

Singh, R. (2024). Lightweight metals and alloys in electric vehicle manufacturing: enhancing performance and efficiency. International Journal of Science and Research Archive, 13 (2), 1735-1742. https://doi.org/10.30574/ijsra.2024.13.2.2343.

Singh, M. K., Palaniappan, S. K., Suyambulingam, I., Rangappa, S. M., Siengchin, S., & Marinkovic, D. (2025). Sustainable composite materials for electric vehicle applications: a comprehensive review. Facta Universitatis, Series: Mechanical Engineering, 1-28. https://doi.org/10.22190/FUME240828014S.

Stewart, R. (2010). Automotive composites offer lighter solutions. Reinforced Plastics, 54 (2), 22-28. https://doi.org/10.1016/S0034-3617(10)70061-8.

Sun, X., Li, Z., Wang, X., & Li, C. (2020). Technology development of electric vehicles: A review. Energies, 13 (90), 1-29. https://doi.org/10.3390/en13010090.

Tanvir, S. M., & Harun-ur Rashid, A. B. M. (2024). Battery charging technologies and standards for electric vehicles: A state-of-the-art review, challenges, and future research prospects. Energy Reports, 11, 5978-5998. https://doi.org/10.1016/j.egyr.2024.05.062.

Tewari, R., Singh, M. K., & Zafar, S. (2021). Utilization of forest and plastic wastes for composite manufacturing using microwave-assisted compression molding for low load applications. Journal of Polymer Research, 28 (11), 1-14. https://doi.org/10.1007/s10965-021-02778-6.

Verrey, J., Wakeman, M. D., Michaud, V., & Manson, J. A. E. (2006). Manufacturing cost comparison of thermoplastic and thermoset RTM for an automotive floor pan. Composites Part A: Applied Science and Manufacturing, 37 (1), 9-22. https://doi.org/10.1016/j.compositesa.2005.05.048.

Wan, B. B., Chen, W. P., Lu, T. W., Liu, F., Jiang, Z., & Mao, M. (2017). Review of solid state recycling of aluminum chips. Resources, Conservation and Recycling, 125, 37-47. https://doi.org/10.1016/j.resconrec.2017.06.004.

Wan, X., & Huang, Y. (2018). Friction stir welding of dissimilar aluminum alloy and steel: A review. The International Journal of Advanced Manufacturing Technology, 99, 1781-1811. https://doi.org/10.1007/s00170-018-2601-x.

Wazeer, A., Das, A., Abeykoon, C., Sinha, A., & Karmakar, A. (2022). Phase change materials for battery thermal management of electric and hybrid vehicles: A review. Energy Nexus, 7, 100131. https://doi.org/10.1016/j.nexus.2022.100131.

Wieckowski, J., & Salabun, W. (2025). Comparative sensitivity analysis in composite material selection: evaluating OAT and COMSAM methods in multi-criteria decision-making. Spectrum of Mechanical Engineering and Operational Research, 2 (1), 1-12. https://doi.org/10.31181/smeor21202524.

Wu, M., & Chen, W. (2022). Forecast of electric vehicle sales in the world and China based on PCA-GRNN. Sustainability, 14 (4), 1-14. https://doi.org/10.3390/su14042206.

Yang, Y., Luo, Z., Zhang, Y., & Su, J. (2024). Dissimilar welding of aluminium to steel: A review. Journal of Manufacturing Processes, 110, 376-397. https://doi.org/10.1016/j.jmapro.2023.12.060.

Yorseng, K., Sanjay, M. R., Tengsuthiwat, J., Pulikkalparambil, H., Parameswaranpillai, J., Siengchin, S., & Moure, M. M. (2019). Information in United States patents on works related to 'natural fibers': 2000-2018. Current Materials Science, 12 (1), 4-76.

Yousefi, E., Khaboshan, H. N., Jaliliantabar, F., & Abdullah, A. A. (2023). The effect of different enclosure materials and NePCMs on performance of battery thermal management system. Materials Today: Proceedings, 75 (1), 1-9. https://doi.org/10.1016/j.matpr.2022.09.261.

Zhan, L., Kong, C., Zhao, C., Cui, X., Zhang, L., Song, Y., & Jiang, Q. (2025). Recent advances on magnesium alloys for automotive cabin components: Materials, applications, and challenges. Journal of Materials Research and Technology, 36, 9924-9961. https://doi.org/10.1016/j.jmrt.2025.05.105.

Zhang, Y., Ye, Q., & Yan, Y. (2024). Processing, microstructure, mechanical properties, and hydrogen embrittlement of medium-Mn steels: A review. Journal of Materials Science & Technology, 201, 44-57. https://doi.org/10.1016/j.jmst.2024.03.014.

Zhou, X. Y., Hu, Z. L., Xiao, X., & Li, M. (2016a). Research on shredding process and characteristics of multi-material plates for recycled cars. Proceedings of the Institution of Mechanical Engineers, Part B: Journal of Engineering Manufacture, 230 (10), 1834-1844. https://doi.org/10.1177/0954405415598927.

Zhou, X. Y., Hu, Z. L., Qin, X., Tao, Y., & Hua, L. (2016b). Study on the stress characteristic and fatigue life of the shredder pin. Engineering Failure Analysis, 59, 444-455. https://doi.org/10.1016/j.engfailanal.2015.11.003.

Zhou, X., Jiang, J., Hu, Z., & Hua, L. (2022). Lightweight materials in electric vehicle. International Journal of Automotive Manufacturing and Materials, 1 (1), 3. https://doi.org/10.53941/ijamm0101003.

SPOR DİPLOMASİSİNDE SEMBOLİK İLETİŞİM VE FİLİSTİN MESELESİNİN ULUSLARARASI GÖRÜNÜRLÜĞÜ: LAMİNE YAMAL ÖRNEĞİ

SYMBOLIC COMMUNICATION IN SPORTS DIPLOMACY AND THE INTERNATIONAL VISIBILITY OF THE PALESTINIAN ISSUE: THE CASE OF LAMINE YAMAL

Assist. Prof. Dr. Dursun YILMAZ[1]
[1]Tokat Vocational School of Health Services, Tokat Gaziosmanpaşa University, Tokat, Türkiye
ORCID: 0000-0001-6161-3482

Assist. Prof. Dr. Erdal Taha AYDOĞDU[2]
[2] Tokat Vocational School, Tokat Gaziosmanpaşa University, Tokat, Türkiye
ORCID: 0000-0002-8587-2339

ÖZET

Günümüzde uluslararası spor organizasyonları, yalnızca sportif rekabetin gerçekleştiği etkinlikler olmanın ötesinde, küresel kamuoyunun şekillendiği ve siyasal ile insani meselelerin sembolik iletişim yoluyla görünürlük kazandığı önemli kamusal alanlara dönüşmüştür. Bu çalışma, Lamine Yamal'ın 2024 UEFA Avrupa Futbol Şampiyonası ile 2026 FIFA Dünya Kupası sürecinde Filistin meselesiyle ilişkilendirilen sembolik iletişim pratiklerini spor diplomasisi, sembolik sermaye ve dijital kamuoyu perspektiflerinden incelemektedir. Araştırmada nitel içerik analizi yöntemi kullanılmış; Yamal'ın kamuya açık sosyal medya paylaşımları, maçlardaki sembolik davranışları, röportajları ve uluslararası medya yansımaları bütüncül olarak analiz edilmiştir. Bulgular, Yamal'ın sportif başarısının uluslararası görünürlüğünü artıran bir sembolik sermayeye dönüştüğünü; çok kültürlü kimliğine yaptığı göndermeler ve Filistin meselesiyle ilişkilendirilen sembolik iletişim pratikleri aracılığıyla uluslararası kamuoyunda spor diplomasisi bağlamında değerlendirilen görünür bir spor figürü hâline geldiğini göstermektedir. Analiz, söz konusu sembollerin yalnızca bireysel kimlik temsili üretmekle kalmadığını, aynı zamanda Filistin meselesinin uluslararası görünürlüğünün yeniden üretilmesinde iletişimsel bir işlev üstlendiğini ortaya koymaktadır. Sonuç olarak çalışma, dijital çağda elit sporcuların sembolik iletişim pratikleri aracılığıyla uluslararası kamuoyunun oluşumuna ve insani meselelerin görünürlüğüne katkı sağlayabildiğini göstermekte; bireysel sporcuların spor diplomasisindeki rolüne ilişkin güncel literatüre özgün bir örnek sunmaktadır.

Anahtar Kelimeler: Spor diplomasisi, sembolik iletişim, Lamine Yamal, Filistin, dijital kamuoyu, sembolik sermaye, spor ve uluslararası ilişkiler.

ABSTRACT

International sporting events have evolved beyond arenas of athletic competition into significant public spaces where global public opinion is shaped and political and humanitarian issues gain visibility through symbolic communication. This study examines the symbolic communication practices associated with the Palestine issue in relation to Lamine Yamal during the period spanning UEFA EURO 2024 and the 2026 FIFA World Cup from the perspectives of sports diplomacy, symbolic capital, and the digital public sphere. Employing a qualitative content analysis approach, the study analyzes Yamal's publicly available social media posts, symbolic behaviors displayed during matches, interviews, and representations in the international media. The findings indicate that Yamal's sporting success evolved into a form of

symbolic capital that enhanced his international visibility. Through references to his multicultural identity and symbolic communication practices associated with the Palestine issue, he emerged as a highly visible sporting figure whose actions were interpreted within the context of sports diplomacy. The analysis further demonstrates that these symbolic practices not only represented his individual identity but also functioned as communicative mechanisms contributing to the reconstruction and international visibility of the Palestine issue. In conclusion, the study argues that, in the digital age, elite athletes can contribute to the formation of global public opinion and to the visibility of humanitarian issues through symbolic communication practices that extend beyond athletic performance. By examining Lamine Yamal as a contemporary case, the study contributes to the literature on sports diplomacy by highlighting the role of individual athletes in the production of symbolic capital and the shaping of international public discourse.

Keywords: Sports diplomacy, symbolic communication, Lamine Yamal, Palestine, digital public sphere, symbolic capital, sport and international relations.

GİRİŞ

Küreselleşme, dijital iletişim teknolojilerindeki gelişmeler ve sosyal medya platformlarının yaygınlaşması, uluslararası spor organizasyonlarının işlevini önemli ölçüde dönüştürmüştür. Günümüzde FIFA Dünya Kupası ve UEFA Avrupa Futbol Şampiyonası gibi organizasyonlar yalnızca sportif rekabetin gerçekleştiği etkinlikler değil, aynı zamanda devletlerin, kurumların ve bireysel aktörlerin uluslararası kamuoyuna mesaj iletebildiği stratejik iletişim alanları hâline gelmiştir. Bu dönüşüm, sporun yumuşak güç kapasitesini artırmış ve spor diplomasisini uluslararası ilişkiler literatürünün önemli çalışma alanlarından biri hâline getirmiştir (Nye, 2008; Murray, 2018).

Dijital iletişim ortamlarının gelişmesiyle birlikte sporcular da yalnızca sportif başarılarıyla değil, toplumsal, kültürel ve insani meselelerle ilişkilendirilen söylem ve sembolik iletişim pratikleri aracılığıyla küresel kamuoyunu etkileyebilen aktörlere dönüşmüştür (Rowe, 2019). Özellikle milyonlarca takipçiye ulaşabilen elit sporcular, sosyal medya paylaşımları, saha içi davranışları ve kamusal görünürlükleri sayesinde uluslararası medya gündeminin oluşumunda etkili olabilmektedir. Bu durum, sporcuların resmî diplomatik aktörlerden bağımsız biçimde uluslararası görünürlük üretebildiği yeni bir spor diplomasisi biçimini ortaya çıkarmıştır.

Bu dönüşümün güncel örneklerinden biri İspanya Millî Takımı oyuncusu Lamine Yamal'dır. Yamal, 2024 UEFA Avrupa Futbol Şampiyonası ile 2026 FIFA Dünya Kupası arasındaki süreçte sergilediği sportif performansın yanı sıra, çok kültürlü kimliğine ilişkin sembolik temsilleri ve özellikle 2026 yılında Filistin bayrağıyla verdiği görüntünün uluslararası medya ve dijital platformlarda geniş yankı uyandırmasıyla dikkat çekmiştir. Bu süreç, tekil bir sembolik eylemin spor medyası, haber kuruluşları ve dijital platformlar aracılığıyla nasıl yeniden dolaşıma sokulduğunu ve farklı anlam katmanları kazanabildiğini göstermesi bakımından dikkate değerdir.

Mevcut literatür incelendiğinde, spor diplomasisi çalışmalarının büyük ölçüde devlet politikaları, mega spor organizasyonları ve kurumsal diplomasi uygulamalarına odaklandığı; bireysel sporcuların sembolik iletişim pratiklerinin uluslararası medya ve dijital kamuoyu

tarafından nasıl çerçevelendiğini ele alan çalışmaların ise sınırlı olduğu görülmektedir. Özellikle Lamine Yamal örneği üzerinden Filistin meselesinin uluslararası görünürlüğünün spor diplomasisi ve sembolik iletişim perspektifinden incelendiği bütüncül akademik bir çalışmaya rastlanmamıştır. Bu boşluk, araştırmanın temel çıkış noktasını oluşturmaktadır.

Bu çalışmanın amacı, Lamine Yamal'ın 2024 UEFA Avrupa Futbol Şampiyonası ile 2026 FIFA Dünya Kupası sürecinde Filistin meselesiyle ilişkilendirilen sembolik iletişim pratiklerini spor diplomasisi perspektifinden analiz etmek ve bu pratiklerin uluslararası medya ile dijital kamuoyu tarafından nasıl anlamlandırıldığını ortaya koymaktır. Bu kapsamda aşağıdaki araştırma sorularına yanıt aranmaktadır:

1. Lamine Yamal'ın Filistin meselesiyle ilişkilendirilen sembolik iletişim pratikleri hangi görsel ve söylemsel göstergeler üzerinden ortaya çıkmaktadır?
2. Yamal'ın çok kültürlü kimliği ve uluslararası sportif görünürlüğü, bu sembolik iletişim pratiklerinin uluslararası medya ve dijital kamuoyunda çerçevelenmesini nasıl etkilemektedir?
3. Uluslararası medya, Yamal'ın sembolik iletişim pratiklerini spor diplomasisi bağlamında nasıl anlamlandırmaktadır?

KAVRAMSAL ÇERÇEVE

Bu çalışma, Lamine Yamal'ın Filistin meselesiyle ilişkilendirilen sembolik iletişim pratiklerini spor diplomasisi bağlamında açıklayabilmek amacıyla dört temel kuramsal eksen üzerine yapılandırılmıştır. Bu eksenler; yumuşak güç ve spor diplomasisi, sembolik iletişim ve sembolik sermaye, dijital kamuoyu ve görünürlük ile aktivist sporcu yaklaşımıdır.

1.1. Yumuşak Güç ve Spor Diplomasisi

Yumuşak güç, devletlerin uluslararası alanda askerî veya ekonomik yaptırımlardan ziyade kültürel çekicilik, değerler ve ikna kapasitesi aracılığıyla etki oluşturmasını ifade etmektedir (Nye, 2008). Spor ise bu etkinin görünür hâle geldiği en önemli diplomatik alanlardan biridir (Murray, 2018). Son yıllarda spor diplomasisi yalnızca devletlerin yürüttüğü resmî faaliyetlerle sınırlı kalmamış; yüksek görünürlüğe sahip sporcular da uluslararası kamuoyunu etkileyebilen diplomatik aktörler olarak değerlendirilmeye başlanmıştır (Manzenreiter, 2010). Bu çalışma, Lamine Yamal örneğinde bireysel sporcuların yüksek uluslararası görünürlüğe sahip sembolik eylemlerinin spor diplomasisi bağlamında nasıl anlamlandırıldığını ve bu eylemlerin uluslararası kamuoyunda nasıl dolaşıma girdiğini incelemektedir.

1.2.Sembolik İletişim, Kimlik ve Sembolik Sermaye

Sembolik iletişim, bireylerin kimliklerini yalnızca sözlü ifadelerle değil; görsel göstergeler, ritüeller, jestler ve diğer sembolik pratikler aracılığıyla da inşa ettiklerini kabul etmektedir. Bourdieu'nun (1991) sembolik sermaye yaklaşımına göre toplumsal tanınırlık ve meşruiyet, belirli semboller aracılığıyla güç ve etki üretebilmektedir. Hall'un (1996) kültürel kimlik yaklaşımı ise kimliğin sabit bir yapı değil, sürekli yeniden üretilen bir temsil süreci olduğunu vurgulamaktadır. Bu kuramsal çerçevede çalışma, Yamal'ın çok kültürlü kimliğine ilişkin sembolik temsilleri, uluslararası turnuvalarda görünür hâle gelen görsel göstergeleri ve Filistin

bayrağıyla ilişkilendirilen sembolik eylemini, sembolik sermaye üretimi ve kimlik temsili bağlamında değerlendirmektedir.

1.3.Dijital Kamuoyu, Görünürlük ve Ağ Toplumu

Dijital medya, sporcuların geleneksel medya kuruluşlarına bağımlı olmadan milyonlarca kişiye doğrudan ulaşabilmesini mümkün kılmıştır. Ağ toplumunda görünürlük, yalnızca iletişim kurmanın değil, aynı zamanda uluslararası gündem oluşturmanın da temel araçlarından biri hâline gelmiştir (Castells, 2015). Bu bağlamda sosyal medya platformları, sporcuların sembolik eylem ve temsillerinin küresel ölçekte dolaşıma girmesine ve uluslararası kamuoyu tarafından yeniden üretilmesine olanak sağlamaktadır (Manor, 2019). Çalışmada Yamal'ın dijital görünürlüğü, Filistin bayrağıyla ilişkilendirilen sembolik eylemin uluslararası medya ve dijital platformlar aracılığıyla nasıl dolaşıma sokulduğunu açıklayan bir iletişim mekanizması olarak ele alınmaktadır.

1.4. Aktivist Sporcu Yaklaşımı

Son yıllarda sporcular yalnızca sportif performanslarıyla değil, toplumsal ve insani meselelerde geliştirdikleri söylem ve eylemlerle de kamusal aktörler olarak öne çıkmaktadır. Aktivist sporcu yaklaşımı, sporcuların sosyal medya paylaşımları, kamusal açıklamaları ve sembolik davranışları aracılığıyla toplumsal tartışmalara katıldıklarını ve uluslararası kamuoyu üzerinde etki oluşturabildiklerini ileri sürmektedir (Coombs & Cassilo, 2017). Bu çalışma, Lamine Yamal'ın Filistin meselesiyle ilişkilendirilen sembolik eylemini doğrudan siyasal bir söylem olarak değil; uluslararası medya ve dijital kamuoyu tarafından insani dayanışma, kimlik temsili ve spor diplomasisi bağlamında yeniden anlamlandırılan bir iletişim pratiği olarak değerlendirmektedir.

2. YÖNTEM

2.1.Araştırmanın Deseni

Bu araştırma, Lamine Yamal'ın Filistin meselesiyle ilişkilendirilen sembolik iletişim pratiklerini spor diplomasisi bağlamında incelemek amacıyla nitel araştırma yaklaşımı çerçevesinde tasarlanmıştır. Araştırmada, yazılı, görsel ve dijital içeriklerde yer alan anlam örüntülerinin sistematik biçimde çözümlenmesine olanak sağlayan nitel içerik analizi yöntemi kullanılmıştır (Mayring, 2014). Bu yöntem, sembolik iletişim pratiklerinin yalnızca doğrudan söylemler üzerinden değil, görsel temsiller, kamusal davranışlar ve medya çerçevelemeleri üzerinden de analiz edilmesine imkân tanımaktadır.

Araştırmanın inceleme dönemini UEFA EURO 2024'ün başlangıç tarihi olan 14 Haziran 2024 ile 2026 FIFA Dünya Kupası'nın sona erdiği 19 Temmuz 2026 tarihleri arasındaki süreç oluşturmaktadır. Bu dönem, Lamine Yamal'ın uluslararası görünürlüğünün belirgin biçimde arttığı ve sembolik eylemlerinin uluslararası medya ile dijital platformlarda yoğun biçimde dolaşıma girdiği zaman aralığını kapsamaktadır.

2.2.Veri Toplama Süreci

Araştırmada ölçüt örnekleme (criterion sampling) yöntemi kullanılmıştır. Veri setine yalnızca araştırma sorularıyla doğrudan ilişkili, kamuya açık ve doğrulanabilir içerikler dâhil edilmiştir.

Analiz kapsamına alınan içeriklerin temel ölçütü, Lamine Yamal'ın Filistin meselesiyle ilişkilendirilen sembolik eylemlerini, çok kültürlü kimlik temsillerini veya bu eylemlerin uluslararası medya ve dijital kamuoyundaki yansımalarını içermesidir. Aynı içeriğin farklı platformlarda tekrar yayımlanması durumunda yalnızca ilk ve doğrulanabilir kaynak analiz birimi olarak değerlendirilmiştir.

Tablo 1. Araştırma Kapsamındaki Veri Kaynakları ve Ölçüt Örnekleme Kriterleri

Veri Türü	Veri Kaynağı	Seçim Ölçütü
Sosyal medya içerikleri	Lamine Yamal'ın doğrulanmış Instagram ve diğer resmî sosyal medya hesapları	Filistin bayrağı, çok kültürlü kimlik ve sembolik temsillere ilişkin kamuya açık paylaşımlar
Görsel veriler	UEFA ve FIFA resmî yayınları ile doğrulanmış görsel kayıtlar	Bayrak kullanımı, jestler ve sembolik davranışlar
Röportajlar ve açıklamalar	FIFA, UEFA ve doğrulanmış uluslararası medya röportajları	Kimlik, spor diplomasisi ve sembolik temsillere ilişkin açıklamalar
Medya içerikleri	BBC Sport, Reuters, Associated Press (AP), FIFA, UEFA ve diğer uluslararası haber kuruluşları	Filistin bayrağı olayı ve Yamal'ın sembolik iletişim pratiklerini konu alan haber ve analizler

Araştırmanın analiz birimini, Lamine Yamal'ın kamuoyunda Filistin meselesiyle ilişkilendirilen sembolik iletişim pratikleri ile bu pratiklerin uluslararası medya tarafından nasıl çerçevelendiğini gösteren her bir iletişim olayı oluşturmaktadır.

2.3. Veri Analizi

Veriler, Mayring'in (2014) nitel içerik analizi yaklaşımı doğrultusunda hem tümevarımsal hem de kuramsal kodlama teknikleri kullanılarak analiz edilmiştir. İlk aşamada veri setinde tekrar eden sembolik göstergeler açık kodlama yöntemiyle belirlenmiş; ikinci aşamada elde edilen kodlar araştırmanın kavramsal çerçevesi doğrultusunda tematik kategoriler altında birleştirilmiştir. Analiz sürecinde MAXQDA 2022 yazılımından yararlanılmıştır.

Tablo 2. Nitel İçerik Analizinde Kullanılan Tema ve Kodlar

Tema	Kodlar	Veri Kaynağı
Sembolik iletişim	Filistin bayrağı, görsel temsil, sembolik eylem	Sosyal medya, maç görüntüleri
Kimlik temsili	Çok kültürlü kimlik, ulusal aidiyet, kültürel temsil	Görseller, röportajlar
Spor diplomasisi	Uluslararası görünürlük, medya çerçevelemesi, sembolik sermaye	Uluslararası haberler, röportajlar
Dijital dolaşım	Viral içerik, yeniden paylaşım, dijital görünürlük	Sosyal medya ve haber içerikleri

2.4. Geçerlik ve Güvenirlik

Araştırmanın inandırıcılığını artırmak amacıyla veri çeşitlemesi (data triangulation) uygulanmıştır. Aynı iletişim olayı; sosyal medya içerikleri, doğrulanmış görsel kayıtlar ve uluslararası haber metinleri üzerinden karşılaştırmalı olarak incelenmiş; farklı veri kaynaklarından elde edilen bulguların tutarlılığı değerlendirilmiştir. Ayrıca analiz sürecinde yalnızca kamuya açık ve doğrulanabilir birincil veya güvenilir ikincil kaynaklardan elde edilen

veriler kullanılmış, doğrulanamayan sosyal medya iddiaları ve spekülatif içerikler analiz kapsamı dışında bırakılmıştır.

3. BULGULAR

3.1. Filistin Bayrağının Sembolik Bir İletişim Eylemine Dönüşmesi

Araştırma kapsamında incelenen veri setinde, Filistin meselesiyle ilişkilendirilen en belirgin sembolik iletişim pratiği; Lamine Yamal'ın 11 Mayıs 2026 tarihinde, FC Barcelona'nın La Liga şampiyonluk kutlamaları sırasında açık otobüs üzerinde Filistin bayrağını taşıması olarak tespit edilmiştir. Söz konusu eylem, herhangi bir yazılı açıklama veya doğrudan siyasi söylem eşliğinde gerçekleşmemiş; buna karşın görsel temsiliyet üzerinden kısa sürede uluslararası medya gündemine taşınmıştır. Associated Press'in (AP) aktarımına göre, yaklaşık 750 bin kişinin katıldığı bu kutlamalarda sergilenen jest, küresel ölçekte geniş yankı uyandırmıştır. Bu bulgu, ilgili dönemde Filistin meselesine dair en görünür sembolik iletişim pratiğinin, doğrudan siyasi bir beyandan ziyade görsel bir gösterge aracılığıyla üretildiğini ortaya koymaktadır.

Uluslararası haber içerikleri, bu sportif jestin siyasal ve insani anlamlar yüklenen sembolik bir pratik olarak çerçevelendiğini göstermektedir. Medya kuruluşlarının olaya yaklaşımındaki söylemsel farklılıklar dikkat çekicidir. Al Jazeera Sport, eylemi takdirle karşılayan bir üslup benimsemiş ve haberde şu doğrudan ifadeye yer vermiştir:

Özgün İngilizce metin: *"Spanish football star Lamine Yamal has been hailed as 'a very brave boy' for waving the Palestinian flag."* (Al Jazeera Sport, 2026)

Türkçe çevirisi: *"İspanyol futbol yıldızı Lamine Yamal, Filistin bayrağını dalgalandırdığı için 'çok cesur bir genç' olarak övgüyle karşılandı."*

Buna karşılık Associated Press (AP) aynı olayı daha nötr bir haber diliyle aktarmıştır. AP haberinde doğrudan bir alıntıya yer verilmemekle birlikte, Yamal'ın kutlamalar sırasında Filistin bayrağını taşıdığı belirtilmiş; teknik direktör Hansi Flick'in ise oyuncunun kendi kararlarını verebilecek olgunlukta olduğunu, ancak kutlamaların odağının sportif başarı olması gerektiğini ifade ettiği aktarılmıştır (Associated Press, 2026).

Bu bulgular, aynı sembolik eylemin farklı medya organları tarafından söylemsel olarak farklı çerçeveler içinde yeniden inşa edildiğini ve kamuoyuna sunulduğunu ortaya koymaktadır.

3.2. Uluslararası Medyanın Sembolik Eylemi Çerçevelemesi

Uluslararası medya kuruluşlarının haber metinleri incelendiğinde, Lamine Yamal'ın Filistin bayrağını taşıma eyleminin salt sportif bir kutlama görüntüsü olarak sınırlı kalmadığı; spor ile siyaset arasındaki ilişkinin yeniden tartışmaya açıldığı küresel bir iletişim olayı biçiminde sunulduğu görülmektedir. Associated Press (AP), olayı Barcelona şampiyonluk kutlamalarının öne çıkan görüntülerinden biri olarak aktarırken; Reuters ve Euronews ise haberlerinde bu jestin uluslararası diplomatik tartışmaları tetiklediğini vurgulamıştır. Euronews, olayın "futbolda siyaset tartışmalarını yeniden canlandırdığı" yönündeki değerlendirmesini doğrudan bir ifadeyle ortaya koymuştur:

Özgün İngilizce metin: *"Yamal waved a Palestinian flag, reviving debate over whether politics and sport should mix."* (Euronews, 2026)
Türkçe çevirisi: *"Yamal'ın Filistin bayrağını taşıması, spor ile siyasetin birbirine karıştırılıp karıştırılmaması yönündeki tartışmaları yeniden alevlendirdi."*
Reuters ise olayın ardından yöneltilen sorular üzerine teknik direktör Hansi Flick'in açıklamalarına yer vererek, eylemin sportif otoriteler nezdinde nasıl karşılandığını aktarmıştır.
İncelenen tüm haberlerde ortak olarak dikkat çeken husus, Yamal'ın Filistin'e ilişkin doğrudan politik bir açıklamasının bulunmamasına karşın, bayrak taşıma eyleminin medya tarafından uluslararası siyasi bağlam içinde anlamlandırılmasıdır. Bu çerçeveleme, Filistin bayrağının yalnızca bireysel bir sembol olmaktan çıkarak, spor diplomasisinin tartışma konusu hâline gelen bir iletişim göstergesine dönüştüğünü ve uluslararası kamuoyunda siyasal içerikli bir mesaj olarak konumlandırıldığını ortaya koymaktadır.

3.3.Kurumsal ve Siyasal Tepkilerin Farklılaşması

Veri setinde yer alan haberler, aynı sembolik eylemin farklı kurumsal aktörler tarafından farklı biçimlerde değerlendirildiğini göstermektedir.

3.3.1. FC Barcelona'nın Yaklaşımı

Uluslararası haber metinlerinde, FC Barcelona'nın söz konusu eylem karşısında kurumsal düzeyde temkinli bir tutum benimsediği gözlenmektedir. Euronews'ün aktardığına göre kulüp, İsrail Savunma Bakanı'nın eleştirileri üzerine konu hakkında herhangi bir resmî açıklama yapmamayı tercih etmiştir. Bu doğrultuda kulübün konumu, Euronews'te yer alan şu doğrudan ifadeyle özetlenmektedir:
Özgün İngilizce metin: *"The club said it had 'no comment' to make on the issue."* (Euronews, 2026)
Türkçe çevirisi: *"Kulüp, konu hakkında herhangi bir yorum yapmayacağını bildirdi."*
Teknik direktör Hansi Flick ise düzenlediği basın toplantısında konuya ilişkin bireysel sorumluluğu önceleyen bir açıklamada bulunmuştur:
Türkçe ifade: *"Normalde bu tür şeyleri tercih etmem. Kendisi artık yetişkin; karar kendisine aittir."* (Flick, basın toplantısı, 2026)
Bu açıklama, bireysel ifade özgürlüğünü tamamen reddetmeyen; ancak kulübün sportif kimliğini siyasal tartışmaların dışında tutmayı amaçlayan bir yaklaşım olarak dikkat çekmektedir.
Öte yandan, aynı sembolik eyleme devlet düzeyinde farklı bir anlam yüklendiği görülmektedir. İspanya Başbakanı Pedro Sánchez, Yamal'ın eylemini kamuoyu önünde desteklemiş ve Filistin bayrağının taşınmasının nefret söylemi olarak değerlendirilemeyeceğini ifade etmiştir (Anadolu Ajansı, 2026; Euronews Türkçe, 2026).
Bu bulgular, aynı sembolik göstergenin kulüp, devlet ve uluslararası medya gibi farklı kurumsal aktörler tarafından birbirinden farklı iletişim çerçeveleri içinde yeniden üretildiğini ve anlamlandırıldığını ortaya koymaktadır.

3.3.2. İspanya Hükümetinin Yaklaşımı

Kulübün temkinli yaklaşımına karşılık, İspanya Hükümeti'nin sembolik eyleme açık destek verdiği görülmektedir.

Uluslararası haberlerde, İsrail Savunma Bakanı Israel Katz'ın Yamal'ı "İsrail'e karşı nefreti körüklemekle" suçlamasının ardından Başbakan Pedro Sánchez'in futbolcuya destek verdiği aktarılmaktadır. Sánchez, Filistin bayrağını taşımanın nefret suçu olarak değerlendirilemeyeceğini belirtmiş ve Yamal'ın davranışının birçok İspanyolun Filistin halkıyla dayanışma duygusunu yansıttığını ifade etmiştir.

Bu bulgu, aynı sembolik eylemin devlet düzeyinde diplomatik söylemin bir parçası hâline geldiğini göstermektedir.

3.4. Uluslararası Medyanın Filistin Meselesinin Görünürlüğüne Katkısı

İncelenen haber içerikleri, Filistin bayrağına ilişkin görüntülerin kısa sürede uluslararası medya gündemine taşındığını göstermektedir.

Örneğin Filistin Futbol Federasyonunun Yamal'a teşekkür mesajı yayımladığı ve bu açıklamanın uluslararası haber ajansları tarafından geniş biçimde haberleştirildiği görülmektedir (Anadolu Ajansı, 2026).

Benzer şekilde The Guardian, 2026 FIFA Dünya Kupası sonrasında Gazze'de İspanya Millî Takımı'na yönelik sempatiyi aktarırken bunun nedenlerinden biri olarak Yamal'ın daha önce Filistin bayrağı taşımasını göstermektedir (The Guardian, 2026).

Bu bulgular, Filistin meselesinin uluslararası görünürlüğünün yalnızca siyasal aktörler aracılığıyla değil, yüksek görünürlüğe sahip sporcuların sembolik iletişim pratikleri ve bu pratiklerin medya tarafından yeniden çerçevelenmesi yoluyla da üretildiğini göstermektedir.

3.5. Bulguların Genel Değerlendirilmesi

Araştırmada incelenen tüm veri kaynakları birlikte değerlendirildiğinde dört temel bulgu ortaya çıkmaktadır:

Tablo 3. Filistin Meselesiyle İlişkilendirilen Sembolik İletişim Pratiklerinin Bulgulara Göre Dağılımı

Bulgular	**Kanıt**
Filistin meselesiyle ilişkilendirilen temel sembolik iletişim pratiği Filistin bayrağıdır.	AP, Reuters, UEFA görüntüleri
Yamal'ın doğrudan Filistin'e ilişkin siyasal bir açıklamasına ulaşılamamıştır.	Röportaj ve medya incelemesi
Uluslararası medya, sembolik eylemi spor-siyaset ilişkisi bağlamında çerçevelemiştir.	Reuters, Euronews, AP
Aynı sembolik eylem kulüp ve devlet tarafından farklı biçimlerde değerlendirilmiştir.	Reuters, Euronews, AP

4. TARTIŞMA VE SONUÇ

Bu araştırma, Lamine Yamal'ın 2024 UEFA Avrupa Futbol Şampiyonası ile 2026 FIFA Dünya Kupası arasındaki dönemde Filistin meselesiyle ilişkilendirilen sembolik iletişim pratiğinin spor diplomasisi bağlamında nasıl anlamlandırıldığını incelemiştir. Bulgular, araştırmanın başlangıcındaki varsayımlardan farklı olarak, Yamal'ın Filistin konusunda doğrudan siyasal söylemler geliştiren bir "aktivist sporcu" profilinden ziyade, tek bir sembolik eylemin uluslararası medya, dijital platformlar ve siyasal aktörler tarafından farklı anlamlarla yeniden üretilmesi üzerinden küresel görünürlük kazanan bir spor figürü olduğunu ortaya koymuştur.

Araştırmanın en önemli bulgusu, Filistin bayrağının taşınmasının tek başına uluslararası bir diplomatik tartışmayı tetikleyebilmiş olmasıdır. Yamal'ın bu eylemine ilişkin doğrudan politik bir açıklaması tespit edilememesine rağmen, uluslararası medya kuruluşları, siyasal aktörler ve dijital kullanıcı toplulukları söz konusu görüntüyü farklı bağlamlarda yeniden yorumlamıştır. Bu durum, sembolik iletişim süreçlerinde anlamın yalnızca mesajı üreten aktör tarafından değil, mesajın dolaşıma girdiği medya ekosistemi tarafından da üretildiğini göstermektedir. Hall'un (1996) temsil yaklaşımı ile Castells'in (2015) ağ toplumu kuramı birlikte değerlendirildiğinde, sembollerin dijital dolaşım sürecinde yeni anlam katmanları kazanması bu araştırmanın bulgularını açıklayan temel mekanizmalardan biri olarak öne çıkmaktadır.

Bulgular ayrıca spor diplomasisinin yalnızca devletlerin yürüttüğü resmî diplomatik faaliyetlerle sınırlı olmadığını göstermektedir. Yumuşak güç yaklaşımının ortaya koyduğu üzere (Nye, 2008), uluslararası görünürlüğe sahip bireyler de kültürel ve sembolik etkileri aracılığıyla kamuoyu oluşturabilmektedir. Ancak bu araştırma, bireysel sporcuların diplomatik etkisinin çoğu zaman doğrudan siyasal söylemlerden değil, uluslararası medya tarafından yeniden çerçevelenen sembolik eylemlerden beslendiğini ortaya koymaktadır. Bu yönüyle çalışma, spor diplomasisinin bireysel aktörler üzerinden nasıl işlediğine ilişkin literatüre güncel bir örnek sunmaktadır (Murray, 2018; Dubinsky, 2019).

Araştırmada dikkat çeken bir diğer bulgu, aynı sembolik eylemin farklı kurumsal aktörler tarafından farklı biçimlerde yorumlanmasıdır. FC Barcelona'nın kurumsal tarafsızlığı korumaya yönelik temkinli yaklaşımı ile İspanya Hükümeti'nin Yamal'a açık destek vermesi, spor alanında ortaya çıkan sembolik eylemlerin yalnızca sportif bağlamda kalmadığını; devlet, kulüp ve medya gibi farklı aktörlerin kendi kurumsal öncelikleri doğrultusunda yeniden anlamlandırıldığını göstermektedir. Böylece tek bir görsel temsil, spor iletişiminin sınırlarını aşarak diplomatik ve siyasal tartışmaların parçası hâline gelebilmektedir.

Bourdieu'nun (1991) sembolik sermaye yaklaşımı açısından değerlendirildiğinde ise Yamal'ın sportif başarısı, tek başına diplomatik etki üretmemektedir. Asıl belirleyici unsur, sahip olduğu sportif görünürlüğün medya tarafından yüksek dolaşıma sahip sembolik bir iletişim olayına dönüştürülmesidir. Başka bir ifadeyle, sportif başarı uluslararası görünürlüğü mümkün kılan başlangıç noktası olurken, sembolik sermaye bu görünürlüğün medya ve dijital kamuoyu tarafından yeniden üretilmesiyle oluşmaktadır.

Araştırmanın bir diğer önemli sonucu, dijital çağda sporcuların uluslararası kamuoyu üzerindeki etkisinin yalnızca sosyal medya paylaşımlarından ibaret olmadığını göstermesidir.

Çalışmada incelenen olayda, doğrudan siyasal içerikli açıklamalar bulunmamasına rağmen görsel bir sembolün uluslararası haber kuruluşları, sosyal medya kullanıcıları ve siyasal aktörler tarafından yeniden dolaşıma sokulması, spor iletişiminde "ikincil anlam üretimi"nin önemini ortaya koymaktadır. Böylece sembolik eylemin etkisi, onu gerçekleştiren kişinin niyetinden bağımsız olarak medya dolaşımı içinde yeniden inşa edilmektedir.

Bu araştırmanın çeşitli sınırlılıkları bulunmaktadır. Çalışma tek bir sporcu ve tek bir sembolik olay üzerinden yürütülmüş olup, yalnızca kamuya açık ve doğrulanabilir veri kaynakları analiz edilmiştir. Ayrıca araştırmada doğrulanamayan sosyal medya içerikleri ile yapay zekâ tarafından üretilmiş görseller analiz kapsamı dışında bırakılmıştır. Dolayısıyla ulaşılan bulgular, Lamine Yamal örneğiyle sınırlı olup diğer sporculara doğrudan genellenemez.

Gelecek araştırmalarda farklı sporcuların benzer sembolik iletişim pratiklerinin karşılaştırmalı olarak incelenmesi, farklı uluslararası krizler bağlamında spor diplomasisinin nasıl işlediğinin araştırılması ve dijital medya platformlarında sembolik eylemlerin kullanıcı etkileşimleri üzerindeki etkisinin nicel yöntemlerle analiz edilmesi literatüre önemli katkılar sağlayacaktır.

Sonuç olarak bu çalışma, Lamine Yamal'ın Filistin meselesiyle ilişkilendirilen sembolik eyleminin uluslararası medya ve dijital kamuoyu tarafından yeniden çerçevelenerek küresel görünürlük kazandığını göstermektedir. Bulgular, spor diplomasisinin yalnızca devletlerin yürüttüğü resmî diplomatik faaliyetlerden ibaret olmadığını; uluslararası görünürlüğe sahip sporcuların sembolik eylemlerinin de medya ve dijital iletişim süreçleri aracılığıyla diplomatik anlam üretebildiğini ortaya koymaktadır. Bu yönüyle çalışma, spor diplomasisi, sembolik iletişim ve dijital kamuoyu literatürünün kesişim noktasında yer alan güncel ve özgün bir örnek sunmaktadır.

KAYNAKÇA

Associated Press. (2026, May 12). *Barcelona's Lamine Yamal waves Palestinian flag during La Liga title celebrations*. https://apnews.com/article/d60e697991db60d9a9ce21b19099d32c

Bourdieu, P. (1991). *Language and symbolic power*. Harvard University Press.

Castells, M. (2015). *Networks of outrage and hope: Social movements in the Internet age* (2nd ed.). Polity Press.

Coombs, D. S., & Cassilo, D. (2017). Athletes and/or activists: LeBron James and Black Lives Matter. *Journal of Sport and Social Issues, 41*(5), 425–444. https://doi.org/10.1177/0193723517719662

Euronews. (2026, May 15). *FC Barcelona tells Euronews "no comment" after Israel accuses Lamine Yamal of fuelling hatred*. https://www.euronews.com/my-europe/2026/05/15/fc-barcelona-tells-euronews-no-comment-after-israel-accuses-yamal-of-fuelling-hatred

Euronews. (2026, May 15). *Spain's PM Pedro Sánchez defends Lamine Yamal over Palestinian flag row*. https://www.euronews.com/my-europe/2026/05/15/spains-pm-pedro-sanchez-proud-of-barcelonas-yamal-in-palestinian-flag-row

Hall, S. (1996). Introduction: Who needs identity? In S. Hall & P. du Gay (Eds.), *Questions of cultural identity* (pp. 1–17). SAGE.

Lincoln, Y. S., & Guba, E. G. (1985). *Naturalistic inquiry*. SAGE.

Manor, I. (2019). *The digitalization of public diplomacy*. Palgrave Macmillan.

Manzenreiter, W. (2010). The Beijing Games in the western imagination of China: The weak power of soft power. *Journal of Sport and Social Issues, 34*(1), 29–48. https://doi.org/10.1177/0193723509358968

Mayring, P. (2014). *Qualitative content analysis: Theoretical foundation, basic procedures and software solution*. Klagenfurt. https://nbn-resolving.org/urn:nbn:de:0168-ssoar-395173

Murray, S. (2018). *Sports diplomacy: Origins, theory and practice*. Routledge.

Nye, J. S., Jr. (2008). *The powers to lead*. Oxford University Press.

Rowe, D. (2019). *The media sports cultural complex: Local and global dimensions* (2nd ed.). Routledge.

UEFA. (2024, July 15). *Lamine Yamal named UEFA EURO 2024 Young Player of the Tournament*. https://www.uefa.com/euro2024/news/

FIFA. (2026). *FIFA World Cup 2026: News, interviews and press conferences*. https://www.fifa.com/

Reuters. (2026, May 20). *Flick says he does not like political gestures after Yamal waved Palestinian flag*. Reuters.

AFP Fact Check. (2026, July 20). *Image of Yamal with Palestinian flag after World Cup is AI-generated*. https://factcheck.afp.com/doc.afp.com.C2NG4G3

Al Jazeera. (2026, May 12). *Lamine Yamal's Palestine flag wave hailed by fans, activists and athletes*. https://www.aljazeera.com/sports/2026/5/12/lamine-yamals-palestine-flag-wave-hailed-by-fans-activists-and-athletes

Anadolu Ajansı. (2026, May 15). *Sanchez, Filistin bayrağı sallayan Lamine Yamal'ı eleştiren İsrailli bakana sert cevap verdi*. https://www.aa.com.tr/tr/dunya/sanchez-filistin-bayragi-sallayan-lamine-yamali-elestiren-israilli-bakana-sert-cevap-verdi/3937958

Euronews Türkçe. (2026, May 15). *İspanya Başbakanı Sanchez'den Filistin bayrağı açan Lamine Yamal'a destek: "Gurur duyuyoruz"*. https://tr.euronews.com/my-europe/2026/05/15/ispanya-basbakani-sanchezden-filistin-bayragi-acan-lamine-yamala-destek-gurur-duyuyoruz

Lead Stories. (2026, July 21). *Fact check: Fake photo shows Lamine Yamal draped in the Palestine flag holding World Cup trophy*. https://leadstories.com/hoax-alert/2026/07/fact-check-fake-photo-shows-lamine-yamal-draped-in-the-palestine-flag-holding-world-cup-trophy.html

T24. (2026, May 15). *Şampiyonluk kutlamalarında Filistin bayrağı taşıyan Lamine'ye İsrail'den tepki, Sánchez'den destek: "Onunla gurur duymak için bir neden daha"*. https://t24.com.tr/dunya/sampiyonluk-kutlamalarinda-filistin-bayragi-tasiyan-lamineye-israilden-tepki-snchezden-destek-onunla-gurur-duymak-icin-bir-neden-daha,1321984

The Guardian. (2026, July 21). *Gaza celebrates World Cup victory of Spanish team they took into their hearts*. https://www.theguardian.com/football/2026/jul/21/gaza-celebrates-world-cup-victory-of-spanish-team-they-took-into-their-hearts

NATO ANKARA ZİRVESİ BAĞLAMINDA LİDERLİK PROFİLİNİN İLETİŞİM YOLUYLA İNŞASI: CUMHURBAŞKANI RECEP TAYYİP ERDOĞAN ÖRNEĞİ

THE CONSTRUCTION OF A LEADERSHIP PROFILE THROUGH COMMUNICATION IN THE CONTEXT OF THE NATO ANKARA SUMMIT: THE CASE OF PRESIDENT RECEP TAYYIP ERDOĞAN

Assist. Prof. Dr. Erdal Taha AYDOĞDU[1]
[1] Tokat Vocational School, Tokat Gaziosmanpaşa University,
Tokat, Türkiye
ORCID: 0000-0001-6830-8763

Assist. Prof. Dr. Dursun YILMAZ[2]
[2]Tokat Vocational School of Health Services, Tokat Gaziosmanpaşa University,
Tokat, Türkiye
ORCID: 0000-0001-6161-3482

ÖZET

Uluslararası zirveler, devlet liderlerinin yalnızca diplomatik müzakereler yürüttükleri platformlar değil, aynı zamanda liderlik profillerini ulusal ve uluslararası kamuoyuna yönelik stratejik iletişim yoluyla inşa ettikleri çok katmanlı alanlardır. Buna karşın NATO zirvelerine ilişkin literatür büyük ölçüde güvenlik politikaları, ittifak ilişkileri ve dış politika kararlarına odaklanmakta; liderlik profilinin iletişimsel süreçlerle nasıl inşa edildiğini iletişim bilimi perspektifinden inceleyen çalışmalar ise sınırlı kalmaktadır. Bu araştırma, söz konusu boşluktan hareketle 2026 NATO Ankara Zirvesi bağlamında Cumhurbaşkanı Recep Tayyip Erdoğan'ın liderlik profilinin hangi iletişim stratejileriyle inşa edildiğini ortaya koymayı amaçlamaktadır. Çalışmada liderlik profili, liderin kişisel özelliklerini ifade eden bir kavram olarak değil; söylem, sembolik temsil, diplomatik etkileşim ve dijital iletişim uygulamaları aracılığıyla kamuoyu nezdinde üretilen ve görünür kılınan liderlik kimliği olarak ele alınmaktadır. Araştırma, nitel araştırma yaklaşımı çerçevesinde durum çalışması desenine dayanmaktadır. Veri setini, NATO Ankara Zirvesi sürecinde Cumhurbaşkanı Recep Tayyip Erdoğan'ın resmî konuşmaları, basın toplantıları, Cumhurbaşkanlığı açıklamaları, ikili görüşmelere ilişkin resmî bilgilendirmeler, kurumsal dijital platformlarda paylaşılan içerikler ile zirve sürecine ait görsel ve sembolik iletişim materyalleri oluşturmaktadır. Verilerin analizinde söylem analizi ve tematik analiz tekniklerinden yararlanılacak; bulgular stratejik liderlik iletişimi, çerçeveleme kuramı, stratejik anlatı yaklaşımı ve kamu diplomasisi perspektifinde değerlendirilecektir. Araştırmanın, NATO zirvelerini iletişim bilimi perspektifinden ele alarak liderlik profilinin iletişim yoluyla inşasını açıklayan bütüncül bir analitik çerçeve sunması beklenmektedir. Bu doğrultuda çalışma, liderliğin uluslararası zirvelerde yalnızca politik kararlar ve diplomatik müzakerelerle değil; söylemsel, sembolik, diplomatik ve dijital iletişim pratiklerinin etkileşimiyle şekillenen dinamik bir süreç olduğunu ortaya koymaktadır. Bu yönüyle araştırmanın siyasal iletişim, uluslararası iletişim, liderlik iletişimi ve kamu diplomasisi literatürüne kuramsal ve yöntemsel katkı sağlaması, ayrıca uluslararası zirvelerde liderlik iletişiminin analizine yönelik uygulanabilir bir kavramsal çerçeve geliştirmesi beklenmektedir.

Anahtar Kelimeler: NATO Ankara Zirvesi, liderlik iletişimi, liderlik profilinin inşası, stratejik liderlik iletişimi, siyasal iletişim, kamu diplomasisi, söylem analizi.

ABSTRACT

International summits are not merely platforms where state leaders conduct diplomatic negotiations; they are also multidimensional communication arenas in which leadership profiles are constructed through strategic communication directed toward both national and international audiences. Despite this, the existing literature on NATO summits has primarily focused on security policies, alliance relations, and foreign policy decisions, while studies examining how leadership profiles are communicatively constructed from a communication studies perspective remain limited. Addressing this gap, the present study aims to examine the communication strategies through which President Recep Tayyip Erdoğan's leadership profile was constructed during the 2026 Ankara NATO Summit. In this study, the concept of *leadership profile* is not approached as a reflection of an individual leader's personal or psychological characteristics; rather, it is conceptualized as a communicatively constructed leadership identity that is produced and made visible through discourse, symbolic representation, diplomatic interaction, and digital communication practices in the eyes of national and international publics. The study adopts a qualitative case study design. The dataset consists of President Erdoğan's official speeches, press conferences, statements released by the Presidency, official briefings concerning bilateral meetings, content shared through the official digital communication channels of the Presidency and related institutions, and visual and symbolic communication materials produced throughout the summit. The data will be analyzed using discourse analysis and thematic analysis, and the findings will be interpreted within the theoretical frameworks of strategic leadership communication, framing theory, strategic narrative, and public diplomacy. The study is expected to offer a holistic analytical framework for explaining the communicative construction of leadership profiles in the context of NATO summits from a communication studies perspective. Accordingly, it seeks to demonstrate that leadership at international summits is not constituted solely through political decision-making and diplomatic negotiations but is dynamically constructed through the interaction of discursive, symbolic, diplomatic, and digital communication practices. In doing so, the study is expected to make both theoretical and methodological contributions to the literature on political communication, international communication, leadership communication, and public diplomacy, while also proposing a conceptual framework applicable to the analysis of leadership communication in international summit diplomacy.

Keywords: Ankara NATO Summit, leadership communication, communicative construction of leadership profile, strategic leadership communication, political communication, public diplomacy, discourse analysis.

İHRACATTA ÜRÜN VE ÜLKE BAZINDA UYULMASI GEREKEN KURALLARIN TÜRKİYE DIŞ TİCARETİNE ETKİSİ

THE IMPACT OF PRODUCT- AND COUNTRY-SPECIFIC RULES TO BE COMPLIED WITH IN EXPORTS ON TURKEY'S FOREIGN TRADE

Lecturer Yücel NAKTİYOK
Department of Foreign Trade, Vocational School of Social Sciences, Atatürk University, Erzurum, TÜRKİYE
ORCID ID: https://orcid.org/0000-0002-8429-1946

Lecturer Fatih UZUNBOY
Department of Foreign Trade, Vocational School of Social Sciences, Atatürk University, Erzurum, TÜRKİYE
ORCID ID: https://orcid.org/0000-0002-7045-1072

ÖZET

Bu çalışma, uluslararası ticarete uyulması gereken kurallar arasında yer alan, ülke ve ürün bazlı kuralların Türkiye'nin dış ticaret performansı üzerindeki etkilerini incelemeyi amaçlamaktadır. Uluslararası ticarette tarife dışı engeller olarak adlandırılan teknik ve idari kurallar öncelikli olarak, üründe kalite ve can güvenliğini hedeflerken, ülke bazlı kurallar ikili veya çok taraflı yapılan anlaşmalarla ülkelerin birbirlerine sağladıkları imtiyazlar yanında, çeşitli tarife dışı engelleri de kapsamaktadır. Ürün bazlı kurallar; gıda, kimyasal, orman ve hayvancılık, radyasyon ve ahşap ambalaj malzemesi gibi sınıflara ayrılmış olup, temelde kalite ve can güvenliğini esas almaktadır. AB'ye ihracatta Avrupa'ya Uygunluk (CE) işareti ve Avrupa Normu (EN) standartları, İslam ülkelerinde helal sertifikası, ABD'de FDA (Amerika Birleşik Devletleri Gıda ve İlaç Dairesi) onayı, Rusya GOST-R sertifikası, ahşap ambalaj malzemelerinde ISPM-15, bitki sağlığı standartları (fitosaniter) gibi belgeler talep edilmektedir. Bu belgelerle birlikte teknik ve idari yükümlülükler, ürün bazlı kuralların uluslararası pazarlara mal satışında ön koşul olma niteliğini göstermektedir. Ülke bazlı kurallar ise ikili veya çok taraflı yapılan anlaşmalarla ülkelerin birbirlerine sağladıkları imtiyazlar yanında, çeşitli tarife dışı engelleri de kapsamaktadır. Türkiye'nin yapmış olduğu, AB ile Gümrük Birliği Anlaşması, EFTA ve diğer ülkelerle Serbest Ticaret Anlaşmaları, Genelleştirilmiş Tercihler Sistemi (GTS) gibi tek taraflı ayrıcalıkları içeren anlaşmalar, ülke ihracatına gümrük muafiyeti, ihracat kolaylığı ve tercihli ticaret gibi olumlu katkılar sağlamaktadır. Diğer yandan çeşitli ülkelerin uyguladığı görünmez engeller, menşe kuralları, anti-damping/korunma önlemleri ve AB'nin Sınırda Karbon Düzenleme Mekanizması (CBAM) çerçevesinde uyguladığı ek vergiler gibi tarife dışı engeller ihracatı olumsuz yönde etkilemektedir. Nitel içerik analizi yöntemiyle Ticaret Bakanlığı, Türkiye İhracatçılar Meclisi, TÜİK ve Avrupa Komisyonu kaynaklarından derlenen verilere dayanan çalışma, Türkiye'nin başlıca ihracat pazarları olan AB, ABD ve komşu ülkeler özelinde bu kuralların ortaya çıkardığı fırsat ve tehditleri değerlendirmektedir. Bu çalışma; büyük firmalar ile sermaye ve üretim kapasitesi sınırlı KOBİ'ler arasındaki rekabet farkının derinleşme riskine dikkat çekerek, Türkiye'nin öncelikli olarak AB ve ABD

standartlarına uyum kapasitesini güçlendirmesi, sertifikasyon çalışmalarına ağırlık vermesi ve pazar çeşitlendirmesine yönelmesi gerektiği sonucuna varmaktadır.
Anahtar Kelimeler: İhracat Kuralları, Tarife Dışı Engeller, CE İşareti, CBAM, Menşe Kuralları, Türkiye Dış Ticareti.

ABSTRACT

This study aims to examine the effects of country- and product-based rules, which are among the regulations that must be complied with in international trade, on Türkiye's foreign trade performance. Technical and administrative regulations, referred to as non-tariff barriers in international trade, primarily aim to ensure product quality and consumer safety, while country-based rules encompass not only the privileges granted to countries through bilateral or multilateral agreements but also various non-tariff barriers. Product-based rules are classified into categories such as food, chemicals, forestry and livestock, radiation, and used goods, and are fundamentally based on quality and consumer safety. For exports to the European Union, documents such as the European Conformity (CE) marking and European Norm (EN) standards are required; in Islamic countries, halal certification; in the United States, FDA (Food and Drug Administration) approval; in Russia, the GOST-R certificate; ISPM-15 compliance for wooden packaging materials; and phytosanitary certificates for plant health standards. Together with these documents, the associated technical and administrative obligations demonstrate that product-based rules constitute a prerequisite for selling goods in international markets. Country-based rules, on the other hand, include not only the privileges granted through bilateral or multilateral agreements but also various non-tariff barriers. Agreements concluded by Türkiye, such as the Customs Union Agreement with the European Union, Free Trade Agreements with EFTA and other countries, and unilateral preferential arrangements such as the Generalized System of Preferences (GSP), provide positive contributions to the country's exports through customs exemptions, export facilitation, and preferential trade. Conversely, non-tariff barriers such as invisible trade barriers imposed by various countries, rules of origin, anti-dumping and safeguard measures, and additional taxes implemented under the European Union's Carbon Border Adjustment Mechanism (CBAM) negatively affect exports. Based on data compiled from the Ministry of Trade, the Turkish Exporters Assembly (TİM), the Turkish Statistical Institute (TurkStat), and the European Commission using the qualitative content analysis method, this study evaluates the opportunities and threats arising from these rules, with a particular focus on Türkiye's main export markets, namely the European Union, the United States, and neighboring countries. The study concludes that Türkiye should primarily strengthen its capacity to comply with European Union and United States standards, place greater emphasis on certification activities, and pursue market diversification, while also drawing attention to the risk of a widening competitive gap between large firms and small and medium-sized enterprises (SMEs) with limited capital and production capacity.
Keywords: Export Regulations, Non-Tariff Barriers, CE Marking, CBAM, Rules of Origin, Türkiye's Foreign Trade.

1. GİRİŞ

1990'lı yıllarda Sovyetler Birliği ve Doğu Bloğu ülkelerindeki rejim değişikliği, Dünya Ticaret Örgütü'nün ticaretin önündeki engelleri azaltması, AB benzeri yeni ticari ortaklıkların kurulması ve ülkeler arası serbest ticaret anlaşmaları gibi gelişmeler, uluslararası ticarette olumlu katkılar sunmuştur. Ancak bu olumlu gelişmeler sonucu ülkeler arası soğuk savaş faaliyetleri hız keserken, küresel alanda ticaret savaşlarını da beraberinde getirmiştir. Dünya Ticaret Örgütü'nün çalışmaları ve çeşitli anlaşmalar yoluyla dünya genelinde gümrük tarifeleri azaltılırken, tarife dışı engeller olarak adlandırılan ürün ve ülkeye özgü kurallar, günümüz uluslararası ticaretinin belirleyici unsurları arasında öne çıkmaktadır. Ürün kalite ve güvenliğine yönelik standartlar, kontrol ve uygunluk belgeleri, sağlık sertifikaları, menşe kuralları, anti-damping ve korunma önlemleri ile son dönemde ön plana çıkan çevreyi korumaya yönelik düzenlemeler, ihracatçı firmaların hedef pazara erişimini doğrudan etkilemektedir (Dünya Ticaret Örgütü [DTÖ], 2024).

Türkiye'nin halihazırda yürürlükte olan, AB ile Gümrük Birliği Anlaşması, EFTA ülkeleri ve 23 ülke ile Serbest Ticaret Anlaşması (STA) ve 6 ülke ile devam eden STA müzakereleri (T.C. Ticaret Bakanlığı, 2026), firmalarımıza pazara erişim ve maliyet avantajı sağlarken, diğer yandan bu ülkelerin uyguladığı çok sayıda düzenlemeye de uyum zorunluluğu getirmektedir. AB, ABD, Rusya ve Uzak Doğu ülkelerinde uygulamaya konulan ticari normlar dünya ticaretinde belirleyici rol oynamakla birlikte, Dünya Ticaret Örgütü de bu ülkelerin normlarına uygun hareket sahası belirlemektedir (DTÖ, 2024). Bu durum, ihracatçı firmalar için uluslararası ticarette: Avrupa'ya Uygunluk (CE) işareti, Avrupa Normu (EN) standartları, Amerika Birleşik Devletleri Gıda ve İlaç Dairesi (FDA) onayı, Rusya GOST-R sertifikası ve Türkiye'de Dış Ticarette Risk Esaslı Kontrol Sistemi (TAREKS) denetimleri gibi kapsamlı bir uygunluk yükümlülüğü getirmektedir (T.C. Ticaret Bakanlığı, 2018-2026). Benzer şekilde, Körfez ve Orta Doğu ülkeleri ile Afrika ülkeleri gibi diğer önemli pazarlarda da kendine özgü tarife ve tarife dışı uygulamalar söz konusudur.

Bu çalışmanın amacı, ihracatta ürün ve ülke bazlı kuralları sistemli ve anlaşılır bir biçimde ele alarak, bu kuralların Türkiye'nin dış ticaret performansı üzerindeki etkilerini güncel veriler ışığında değerlendirmektir.

2. KAVRAMSAL ÇERÇEVE: TİCARETTE TEKNİK ENGELLER VE ÜLKE BAZLI DÜZENLEMELER

Türkiye'de Ticaret Bakanlığı bünyesinde yer alan Ürün Güvenliği ve Denetimi Genel Müdürlüğü; Dünya Ticaret Örgütü'nün Ticarette Teknik Engeller Anlaşması (TBT) çerçevesinde (DTÖ, 2024), ülkelerin insan sağlığı, çevre ve tüketici güvenliğini korumak amacıyla getirdiği teknik düzenlemelerin uyum sürecini yönetmektedir. Bakanlık oldukça farklı olan bu düzenlemelerin firmalar açısından maliyetini düşürmek ve dış ticaret engeli işlevi olmasını ortadan kaldırmak için gerekli tedbirlerin alınması ve diğer ülkelerle uyum sağlamak amacıyla sürekli güncelleme faaliyetleri içerisindedir. Literatürde bu düzenlemeler iki temel başlıkta ele alınmaktadır.

2.1. Ürün Bazlı Kurallar

Ürün bazlı kurallar, malın fiziksel ve kimyasal özelliklerine göre kalite ve can güvenliğini esas almakta olup, malın yurt dışı pazara girişi esnasında taşıması gereken teknik, sağlık, dinî veya kültürel nitelikteki koşulları içerir. Kurallar gıda, kimyasal, orman ve hayvancılık, radyasyon vb. olmak üzere neredeyse tüm ürünleri kapsayacak biçimde sınıflandırılmıştır. Günümüzde tüketici sağlığını ve ürün kalitesini güvence altına almak isteğe bağlı bir tercih değil, zorunlu bir ön koşul haline gelmiştir. Kurallara uymamanın maliyeti yalnızca finansal değil, itibar ve pazar kaybı gibi hususlara ek olarak sınırda geri çevrilme veya imha riskleri de firmaları olumsuz yönde etkilemektedir.

Dış ticarette uyulması gereken ürün bazlı kuralları ana ürün gruplarına göre özetleyecek olursak:

Gıda ürünleri: sağlık, hijyen, izlenebilirlik ve katkı maddesi sınırlamaları gibi denetimlere tabi tutulan en hassas ihracat türüdür.

Kimyasal ürünler: insan sağlığı ve çevresel etkileri nedeniyle en sıkı denetime tabi tutulan ürün grupları arasında yer almaktadır. Avrupa Birliği, ABD ve diğer ülkeler dış ticarette kimyasal mevzuatına uyumu zorunlu kılmaktadır.

Orman ve Hayvancılık Ürünleri: hem biyogüvenlik hem de sürdürülebilirlik endişeleri nedeniyle özel düzenlemelere tabidir. Hayvansal ürünlerin yasa dışı kesim veya hastalık taşıma riskinden arındırıldığına dair belge (veteriner sağlık sertifikaları), dış pazara girişin ön koşuludur.

Radyasyon Kontrolü: gıda ve tarım ürünlerinde nükleer kirlenme riskine karşı özellikle belirli bölgelerden gelen ürünler için radyasyondan ari belgesi talep edilir. Belge Türkiye Atom Enerjisi Kurumu tarafından verilmektedir.

CE ve EN: CE (Avrupa'ya Uygunluk) işareti, ürünün Avrupa Birliği'nin sağlık, güvenlik ve çevre standartlarına uygunluğunu gösteren belgedir. EN (Avrupa Normu) standartları, ürünlerin teknik performans ve güvenlik kriterlerini tanımlayan hususlardan oluşur. CE ve EN olmadan birçok ürün grubunun AB pazarına girişi mümkün değildir. Avrupa Birliği'ne ihraç edilecek makine, elektrikli cihaz, oyuncak ve tıbbi cihaz gibi ürünlerin CE işareti taşınması zorunludur; bu işareti taşımayan ürünün AB üyesi ülkelere ihracatı mümkün değildir. CE işaretinin teknik altyapısını EN standartları oluşturmakta, Türkiye'de bu standartlar Türk Standartları Enstitüsü (TSE) tarafından ulusal standart olarak yayımlanmaktadır (Türk Standartları Enstitüsü [TSE], 2026.).

FDA onayı: Amerika Birleşik Devletleri Gıda ve İlaç Dairesi (FDA) onayı, gıda, ilaç ve kozmetik ürünlerin ABD pazarına girişinde kullanılır.

GOST-R sertifikası, Rusya pazarına giren ürünlerin teknik güvenlik standartlarına uygunluğunu onaylayan bir belgedir.

Helal sertifikası ve Koşer belgesi: Körfez ülkeleri ve Malezya gibi İslam ülkelerine yapılan ürünlerin ihracatında dini ve kültürel uygunluğu gösteren belge olup, müftülükler tarafından

düzenlenir. Koşer belgesi ise aynı tür ürünler için İsrail tarafından istenilen bir belge olup hahambaşılar tarafından düzenlenir.
ISPM-15 standardı: ahşap ambalaj malzemeleri ve diğer orman ürünlerinde hammaddeden ambalajlamaya kadar üretim zincirinin bütün aşamaları için kullanılan belgedir.
Fitosaniter sertifika uygulamaları: bitkisel ürünlerin zararlı organizma ve hastalıklardan ari olduğunu belgelemektedir.
Yukarıda başlıcaları açıklanan ürün bazlı kurallar, neredeyse tüm pazarlarda ortak bir gereklilik olarak ihracatçı firmaların karşısına çıkmaktadır.

2.2. Ülke Bazlı Kurallar

Ülke bazlı kurallar ise ikili veya çok taraflı anlaşmalarla ülkelerin birbirlerine tanıdığı ayrıcalık veya kısıtlamaları kapsayan geniş bir düzenleme alanıdır. Bu kurallar, ürün bazlı düzenlemelerden farklı olarak malın niteliğinden çok ihracatçı ülkenin alıcı ülke ile ilişki ve anlaşmalarına göre düzenlenmektedir. Ülke bazlı kurallar, ihracatçılara önemli ayrıcalık ve maliyet avantajları sağlar veya ticareti sınırlayıcı rol oynayabilir. Ayrıcalıkların menşe kurallarına bağlı olması, uygulamada ciddi bir bilgi birikimi gerektirmektedir.
Özetle ülke bazlı kurallar ticaret ortaklarının veya bir bölgenin kendine özgü uyguladığı tarife, tarife dışı önlem ve menşe rejimlerinden oluşur. Bu kapsamda menşe kuralları, ürünün hangi ülkede 'yeterli işçilik ve işlem' gördüğünü belirleyerek tercihli tarife avantajlarından yararlanıp yararlanamayacağını belirlemektedir. İthal ürünlerin yerli üreticiye zarar verdiği tespit edildiğinde ise anti-damping ve diğer koruma önlemleri uygulanarak ilave gümrük vergileri tahsil edilir. Bu durum hem Türkiye'nin ithalatını hem de üçüncü ülkelerin Türk ürünlerine yönelik uygulamalarını kapsayan çift yönlü bir araçtır.
Avrupa Birliği'nin Sınırda Karbon Düzenleme Mekanizması (CBAM) son dönemde işlerlik kazanan bir diğer ülke bazlı düzenlemedir. Bu mekanizma, karbon yoğun sektörlerde AB dışından yapılan ithalata gömülü emisyona dayalı ek bir maliyet ortaya çıkarmaktadır.

3. ARAŞTIRMANIN AMACI VE YÖNTEMİ

Bu çalışmada, resmi kurum raporları ve diğer veri kaynakları taranarak nitel içerik analizi yöntemi uygulanmıştır. Ticaret Bakanlığı, TİM, TÜİK ve Avrupa Komisyonu verileri, birbirini doğrulayacak şekilde çapraz karşılaştırmaya tabi tutulmuş olup, bu yöntem, sayısal verilerin ötesinde kuralların yapısal ve stratejik anlamını da görmemizi sağlamaktadır.
Veri kaynakları arasında Dünya Ticaret Örgütü'nün ticarette teknik engellere ilişkin düzenlemeleri (DTÖ, 2024), T.C. Ticaret Bakanlığı mevzuat duyuruları ve serbest ticaret anlaşması verileri (T.C. Ticaret Bakanlığı, 2018-2026, 2026), Resmî Gazete tebliğleri (Resmî Gazete, 2001, 2025), Türkiye İstatistik Kurumu (TÜİK) dış ticaret istatistikleri (Türkiye İstatistik Kurumu [TÜİK], 2018-2026), Türkiye İhracatçılar Meclisi'nin sürdürülebilirlik raporları (Türkiye İhracatçılar Meclisi [TİM], 2025), Avrupa Komisyonu'nun CBAM'a ilişkin resmî açıklamaları (Avrupa Komisyonu, 2023-2026) yer almaktadır. Çalışma kapsamında, Türkiye'nin başlıca ihracat pazarları olan Avrupa Birliği, Amerika Birleşik Devletleri ve komşu

bölge ülkeleri (Rusya/BDT, Orta Doğu) özelinde uygulanan ürün ve ülke bazlı kurallar sistematik olarak derlenmiş; bu kuralların dış ticarete olası etkileri değerlendirilmiştir. Araştırma, dış ticareti koordine eden kurum ve ihracatçı firmalara yol haritası sunmayı amaçlamaktadır.

4. AVRUPA BİRLİĞİ PAZARINA YÖNELİK KURALLAR VE ETKİLERİ

Türkiye ile AB arasındaki Gümrük Birliği Anlaşması, işlenmiş tarım ve sanayi ürünlerinde karşılıklı gümrük muafiyeti yoluyla ihracata önemli bir artış sağlamıştır. Bu anlaşma, Türk sanayisinin AB standartlarına uyum sağlamasına aracı olurken diğer yandan, AB'nin teknik mevzuatının Türkiye tarafından da uygulanmasını öngörmektedir (Türkiye-Avrupa Topluluğu Ortaklık Konseyi, 1995). Bu çerçevede 4703 sayılı Kanun ile CE işareti uygulaması Türkiye'de de zorunlu hale getirilmiş (Resmî Gazete, 2001); ithalat ve ihracatta ürün güvenliği denetimleri, Ticaret Bakanlığı tarafından yürütülen TAREKS (Dış Ticarette Risk Esaslı Kontrol Sistemi) üzerinden risk esaslı olarak gerçekleştirilmektedir. EN standartlarına uygun olmayan ve CE işareti taşıması gerektiği halde taşımayan bir ürünün AB üyesi ülkelere ihracatı fiilen mümkün olmadığından, bu uygulama özellikle KOBİ statüsündeki firmalar için önemli bir uyum maliyeti oluşturmaktadır.

AB pazarına yönelik en son düzenleme, Avrupa Yeşil Mutabakatı'nın bir parçası olan Sınırda Karbon Düzenleme Mekanizmasıdır (CBAM) (Avrupa Komisyonu, 2023-2026). CBAM, hâlihazırda demir-çelik, alüminyum, çimento, gübre, elektrik ve hidrojen gibi karbon yoğun ürünleri kapsamakta olup, 1 Ocak 2026 itibarıyla ise kesin uygulama dönemine geçilerek gömülü emisyonlara dayalı bir karbon bedeli alınması zorunlu hale getirilmiştir. Karbon maliyetindeki artış, karbon yoğunluğu yüksek ürünler ihraç eden Türk ihracatçıları AB pazarında fiyat rekabeti açısından dezavantajlı konuma düşürecektir.

5. AMERİKA BİRLEŞİK DEVLETLERİ PAZARINA YÖNELİK KURALLAR VE ETKİLERİ

Amerika Birleşik Devletleri'nin 2 Nisan 2025 tarihinde yürürlüğe koyduğu yeni gümrük tarife rejimi, dünya ticaretinde önemli bir kırılma yaratmıştır. Bu düzenleme kapsamında otomotiv, elektronik ve tekstil gibi sektörlerde gümrük tarifeleri yükseltilmiş, ihracatçı firmalar maliyet ve fiyatlandırma stratejilerini yeniden gözden geçirmek zorunda kalmıştır.

2026 yılı itibarıyla çelik, alüminyum ve otomobil sektörlerinde uygulanan Section 232 tarifelerinin yürürlüğü devam ederken, Türkiye, diğer birçok ticaret ortağıyla eşit düzeyde uygulanan genel Section 122 tarifesine tabi tutulmaktadır. T.C. Ticaret Bakanlığı bu durumu, Türkiye'nin ABD nezdinde dengeli bir ticaret ortağı olarak değerlendirildiğinin göstergesi şeklinde yorumlamaktadır (T.C. Ticaret Bakanlığı, 2018-2026); bununla birlikte, önceki dönemde yüksek oranlı IEEPA'ya (Uluslararası Ekonomik Acil Durum Yetkileri Yasası) tabi olan bazı ülkelerin genel oran seviyesine çekilmesiyle, Türkiye'nin tarife avantajı bir ölçüde azalmıştır. Çelik ve alüminyum sektörlerinde ise Section 232 tarifeleri, ABD pazarına yönelik Türk ihracatını doğrudan etkileyen maliyet unsuru olma özelliğini korumaktadır.

6. DİĞER PAZARLARDA ÜLKEYE ÖZGÜ DÜZENLEMELER

Rusya ve Bağımsız Devletler Topluluğu (BDT) ülkelerine yönelik ticarette GOST-R uygunluk sertifikası zorunluluğu devam etmektedir (T.C. Ticaret Bakanlığı, 2018-2026). Ayrıca Rusya'nın zaman zaman uyguladığı ihracat yasakları (2026 yılı içinde kükürt ihracatına getirilen yasak) Türk firmaların hammadde temininde belirsizlik yaratmaktadır.

Orta Doğu ve Körfez ülkelerine yönelik ihracatta ise gıda, kozmetik ve ilaç sektörlerinde helal sertifikası pazara girişin ön koşuludur. Benzer ürünler için İsrail pazarında koşer belgesi aranmaktadır.

Bunların yanı sıra Türkiye, hem ithalatta yerli üreticisini korumak amacıyla çeşitli ülkelere yönelik anti-damping ve korunma önlemi soruşturmaları açmakta (paslanmaz çelik levha, kalaylı çelik levha, PET reçine, karton gibi ürünlerde), hem de kendi ihracatçıları üçüncü ülkelerde benzer soruşturmalara konu olabilmektedir (T.C. Ticaret Bakanlığı, 2018-2026).

Öte yandan Pan-Avrupa-Akdeniz Menşe Kümülasyonu, Türkiye-Birleşik Krallık ve Türkiye-Malezya gibi serbest ticaret anlaşmaları kapsamında uygulanan tercihli menşe kuralları, Türk ihracatçılarına gümrük vergisi muafiyeti yoluyla önemli bir rekabet avantajı sunmaktadır (T.C. Ticaret Bakanlığı, 2026). Nitekim Türkiye'nin halihazırda EFTA ülkeleri ile birlikte Makedonya, Bosna-Hersek, Filistin, Tunus, Fas, Mısır, Arnavutluk, Gürcistan, Karadağ, Sırbistan, Şili, Morityus, Güney Kore, Malezya, Moldova, Faroe Adaları, Singapur, Kosova, Venezuela, Birleşik Krallık, BAE ve Katar olmak üzere toplam 23 ülkeyle STA'sı yürürlükte olup, Lübnan, Sudan ve Ukrayna ile imzalanan STA'lar ise iç onay süreçlerinin tamamlanmasını müteakip yürürlüğe girecektir. Ayrıca Japonya, Endonezya ve Körfez Arap Ülkeleri İşbirliği Konseyi (KİK) ile STA müzakereleri halen sürdürülmektedir (T.C. Ticaret Bakanlığı, 2026).

Genelleştirilmiş Tercihler Sistemi (GTS), ABD, Kanada, Japonya, Rusya vb. gelişmiş ülkelerin gelişmekte olan ülkelere tek taraflı olarak tanıdığı gümrük indirimlerini ifade etmektedir. Bu sistem, karşılıklılık ilkesi aranmaksızın uygulanması nedeniyle alıcı ülkelerin inisiyatifine bağlı bir durum taşımaktadır. Türkiye bazı ülkelerin sağlamış olduğu tercihli düzenlemelerden yararlanmakla birlikte, bazı ülkelere de tek taraflı vergi indirimleri uygulamaktadır.

7. BULGULAR: KURALLARA UYUMUN TÜRKİYE DIŞ TİCARETİNE ETKİLERİ

Aşağıdaki Tablo 1, başlıca ürün bazlı kuralları ve dış ticarete etkilerini özetlemektedir.

Kural / Belge	Uygulandığı Pazar	Kapsadığı Ürün Grupları	Dış Ticarete Etkisi
CE İşareti / EN Standartları	AB	Makine, elektrikli cihaz, oyuncak, tıbbi cihaz vb.	Zorunlu; taşımayan ürün AB pazarına giremez
FDA Onayı	ABD	Gıda, ilaç, kozmetik, tıbbi cihaz	Zorunlu kayıt ve uygunluk denetimi
GOST-R Sertifikası	Rusya/BDT	Elektrikli cihaz, makine, gıda	Zorunlu uygunluk belgesi

Kural / Belge	Uygulandığı Pazar	Kapsadığı Ürün Grupları	Dış Ticarete Etkisi
Helal Sertifikası	Körfez/Orta Doğu, Malezya	Gıda, kozmetik, ilaç	Pazara giriş için fiilen zorunlu
ISPM-15	Çoğu ülke	Ahşap ambalaj malzemesi	Isıl işlem/damgalama zorunlu
Fitosaniter Sertifika	Tüm ülkeler	Bitkisel ürün, taze meyve-sebze	Bitki sağlığı denetimi zorunlu

Kaynak: T.C. Ticaret Bakanlığı (2018-2026) ve Türk Standartları Enstitüsü (2026).

Tablo 2 ise ülke bazlı kuralları ve bunların Türkiye dış ticaretine yansımalarını göstermektedir.

Ülke / Bölge	Uygulanan Kural / Mekanizma	Kapsadığı Ürünler	Türkiye Dış Ticaretine Etkisi
Avrupa Birliği	CBAM (Sınırda Karbon Düzenleme Mekanizması)	Demir-çelik, alüminyum, çimento, gübre, elektrik, hidrojen	2026 itibarıyla karbon sertifikası maliyeti; rekabet gücünde aşınma riski
Avrupa Birliği	Gümrük Birliği menşe/uygunluk kuralları, TAREKS denetimi	Sanayi ürünlerinin tamamı	Menşe ispatı ve teknik denetim yükümlülüğü
ABD	Section 232 (çelik-alüminyum-otomotiv) ve Section 122 genel tarifeleri	Çelik, alüminyum, otomotiv ve genel sanayi ürünleri	İlave gümrük vergisi; fiyat rekabetinde dezavantaj
Rusya / BDT	İhracat yasakları, GOST-R uygunluk zorunluluğu	Kükürt, tahıl, çeşitli sanayi ürünleri (karşı yönde)	Girdi tedarikinde belirsizlik, uygunluk maliyeti
Çin ve üçüncü ülkeler	Anti-damping / korunma önlemi soruşturmaları (iki yönlü)	Çelik levha, sentetik iplik, PET reçine, karton vb.	İthalatta ilave vergi; yerli üretici korunurken girdi maliyeti artabilir
Serbest Ticaret Anlaşması Ortakları	Tercihli menşe kuralları (Pan-Avrupa-Akdeniz, İngiltere, Malezya STA'ları)	STA kapsamındaki tüm ürünler	Menşe kümülasyonu sağlanırsa gümrük vergisi muafiyeti

Kaynak: T.C. Ticaret Bakanlığı (2018-2026)

Tablolardan hareketle, ürün ve ülke bazlı kuralların Türkiye dış ticaretine etkisinin iki yönlü olduğu söylenebilir. Bir yandan CE işareti, EN standartları ve tercihli menşe kuralları gibi düzenlemelere uyum sağlayan firmalar, özellikle AB pazarı, Gümrük Birliği'nin sağladığı avantajlar sayesinde Türk ihracatçılar için en önemli pazar konumundadır. Ancak CE, EN ve CBAM gibi düzenlemeler, bu avantajı etkileyen uyum ve maliyet yükü oluşturmaktadır. Standartlara erken ve kapsamlı uyum sağlayan firmalar, rakiplerine karşı sürdürülebilir bir

rekabet üstünlüğü elde etmektedir. AB pazarındaki başarı, yalnızca fiyat değil, teknik ve çevresel uyum kapasitesiyle doğrudan ilişkilendirilmektedir.

ABD tarafından son yıllarda uygulanan FDA ve Section 232/122 tarifeleri gibi düzenlemeler, stratejik sektörlerde ilave maliyet ve rekabet engelleyici etkiye sahiptir. ABD pazarının büyüklüğü ve satın alma gücü, uyum maliyetlerini telafi edebilecek önemli bir fırsat sunmaktadır. ABD'ye yönelik ihracatta sertifikasyon süreçlerine erken yatırım yapan firmalar, uzun vadede istikrarlı bir pazar payı yakalayabilir. Türkiye'nin ABD pazarındaki payının artırılması, sertifikasyon altyapısının geliştirilmesine yönelik alınacak tedbirlere bağlı görünmektedir. Sertifikasyon ve uyum maliyetlerinin yüksekliği, sermaye ve üretim kapasitesi sınırlı KOBİ'leri büyük firmalara göre dezavantajlı konuma sokmaktadır. KOBİ'lere yönelik destek mekanizmaları geliştirilmediği takdirde, rekabet farkının derinleşmesi kaçınılmaz görünmektedir.

Komşu ülkeler, coğrafi yakınlık ve kültürel benzerlik sayesinde Türk ihracatçılar için uyum maliyetinin nispeten az olduğu pazarlardır. Bu pazarlardaki siyasi ve ekonomik istikrarsızlık, ticari öngörülebilirliği zaman zaman ciddi biçimde etkilemektedir. GOST-R gibi ülkeye özgü sertifikasyon gereklilikleri, bölgesel ticaretin de bir uyum bilgisi gerektirdiğini göstermektedir. Bölgesel pazarlara ağırlık verilmesi, AB ve ABD pazarlarındaki risklere karşı önemli bir denge unsuru olarak ortaya çıkmaktadır.

Ayrıca, anti-damping ve korunma önlemleri gibi araçların karşılıklı işletilmesi, Türkiye'nin hem yerli üreticisini üçüncü ülke kaynaklı ucuz ithalata karşı koruma imkanı bulduğunu hem de kendi ihracatçılarının benzer önlemlere maruz kalabileceğini göstermektedir. Bu düzenlemeler, yeterince açık ve ayrıntılı olmadığı durumlarda özellikle küçük ölçekli ihracatçılar için önemli zorluklara dönüşmektedir. Dolayısıyla tarife dışı engellerin doğru sınıflandırılması, etkili bir ihracat planlamasının ön koşulu haline gelmektedir.

8. SONUÇ VE ÖNERİLER

Bu çalışmada, ihracatta ürün ve ülkeye göre uyulması gereken kuralların Türkiye dış ticaretine etkileri, güncel mevzuat ve raporlar ışığında değerlendirilmiştir. Bulgular, söz konusu kuralların mali yükümlülük içeren gümrük tarifelerinden daha etkin bir rol üstlendiğini; CE işareti, EN standartları, helal sertifikası ve fitosaniter belgeler gibi ürün bazlı kuralların pazara giriş için ön koşul niteliği taşıdığını; CBAM, ABD tarife uygulamaları ve GOST-R gibi ülke bazlı düzenlemelerin ise özellikle karbon yoğun ve stratejik sektörlerde maliyet artışı ile birlikte sertifikasyon altyapısının da güçlenmesi zorunluluğunu ortaya koymaktadır. Özellikle Türkiye, AB'ye önemli miktarda çelik ve alüminyum ihraç eden ülkeler arasında yer aldığından, CBAM uygulamalarından en fazla etkilenecek ülkelerden biri olarak değerlendirilmektedir.

Diğer yandan kuralların öngördüğü yüksek maliyetli laboratuvar testleri ve belgelendirme süreçleri KOBİ'ler için ek maliyet ve pazara giriş tereddütleri oluşturmakta, CE ve EN uyumluluk süreci, firmaların bütün tedarik zincirinde dönüşüm gerektirmektedir. Gümrük Birliği'nin tarım ve hizmetler gibi alanları kapsam dışında bırakması, AB ile Türkiye arasında tam bir ticari birlik sağlanmasını engellemiştir. Türkiye'nin tarımsal ihracat potansiyelinin,

fitosaniter altyapısı ile yakından ilişkisi olup, belge süreçlerindeki gecikmeler, özellikle çabuk bozulabilir ürünlerde ciddi ekonomik kayıplara yol açabilmektedir. Bu durum ihracatçı firmaların mevzuat takibini kurumsal bir sürece dönüştürmesini zorunlu kılar. Belge veya bilgi eksikliği sonucu ihracatın gümrükte alıkonulma veya iadesi Türk ihracatçılar için önemli pazar kayıplarına yol açabilir. Türkiye'nin tercihli tarife avantajlarını artıracak STA'ların ürün yelpazesinin dar olması pazar kayıplarına neden olabilmektedir.

Bu çerçevede, Türkiye'nin dış ticaret rekabet gücünün korunması ve artırılması için: (1) Türkiye Emisyon Ticaret Sistemi'nin AB nezdinde tanınırlığının sağlanarak CBAM yükünün mahsup edilebilir hale getirilmesi, (2) Türkiye'nin öncelikli olarak AB ve ABD standartlarına uyum kapasitesini güçlendirerek, test ve belgelendirme altyapısını, sadece firma düzeyinde değil, ulusal ölçekte geliştirilmesi, (3) yeni serbest ticaret anlaşmaları yapılarak, tercihli menşe kurallarından daha geniş ölçekte yararlanılması, (4) tek bir pazarın (özellikle AB) düzenleyici riskleri veya kaos ortamlarında ülke ihracatının sekteye uğramaması için, Türk ihracatçılarının geleneksel pazarlara olan bağımlılığının azaltılması, (5) Gümrük Birliği Anlaşmasının Türkiye lehine güncellenerek genişletilmesi, (6) belgelendirme işlemlerinin kolaylaştırılması, özellikle KOBİ'lere uluslararası pazarlara erişim kolaylığı kamu-özel sektör iş birliği sonucu, KOBİ ölçeğindeki ihracatçılara sertifikasyon ve emisyon raporlama süreçlerinde teknik ve finansal destek sağlanması, (7) ihracatçıların menşe kurallarına tam anlamıyla hakim olmasını sağlayacak çalışmalar yapılması, (8) standartlara uygun üretim yapan ihracatçı firmaların, pazara giriş süresini kısaltıcı tedbirlerin alınması, (9) AB, ABD ve Rusya tarafından yeni uygulamaya konulan sertifikasyon ve çeşitlendirmeden oluşan kurallara, uyum çalışmaları Türkiye dış ticaretinin vizyonu haline getirilmeli, (10) KOBİ'ler ile büyük firmalar arasındaki rekabet farkının derinleşme riski iyi analiz edilerek STK ve kamu otoritelerinin planlı müdahalesi zorunlu hale getirilmelidir.

Yukarıda sayılan öneriler; ihracatta uyulması gereken ülke ve ürün bazlı kuralları, Türkiye açısından risk faktörü olmaktan çıkarıp, rekabet üstünlüğü haline dönüştürecektir.

9. TARTIŞMA

Ürün ve ülke bazlı kurallar Türkiye dış ticareti açısından fırsat mı, tehdit mi?

Kurallar, uluslararası pazarlara girişin tali unsuru olmaktan çıkıp, doğrudan belirleyicisi haline gelmiştir. Bu kurallar, ürünün özelliğine bağlı teknik standartlardan, ülkeler arası siyasi ve ekonomik ilişkilere bağlı ayrıcalık ve yasaklar gibi geniş bir yelpaze oluşturmaktadır. Türkiye gibi ihracata dayalı büyümeyi hedefleyen ekonomi için bu kuralların doğru okunması, rekabet gücünün artarak devam etmesi açısından hayati önem taşımaktadır. Türkiye'nin bu kuralları fırsata dönüştürerek sürdürülebilir ihracat büyümesi sağlaması; uyum kapasitesi, belgelendirme çalışmaları, katma değeri yüksek teknolojik ürünlere öncelik verme ve pazar çeşitlendirmesine bağlıdır.

Türkiye'nin AB ve diğer ülkelerle yapmış olduğu Gümrük Birliği ve STA'lar çerçevesinde uyguladığı uyum süreci ile gümrük vergisi niteliği taşımayan ancak ticareti fiilen kısıtlayan bu

kurallardan, rakip ülkelere göre daha az etkilenecek, teknik ve idari düzenlemeler fırsata dönüşecektir.

Türkiye Emisyon Ticaret Sistemi'nin (TR-ETS) AB tarafından tanınması halinde, ihracatçı firmalarımızın karbon bedelinden muaf tutulması, dünyanın en büyük pazarlarından bir olan AB'ye ihracatta maliyet avantajı kazandıracaktır.

Gümrük tarifelerinin ticaret anlaşmalarıyla azaltılması, tarife dışı engellere uyum kabiliyetini kazanmış Türkiye için fırsat olacaktır. Ayrıca helal belgesi kullanan İslam ülkelerindeki pazarların büyüklüğü, dini, kültürel ve coğrafi yakınlık nedeniyle Türkiye için önemli bir fırsat teşkil etmektedir.

Sonuç olarak uluslararası ticaret, ülkelerin çıkarlarını dengelemeyi amaçlayan kurallar üzerine kurulmuştur. Bu nedenle ürün ve ülke bazlı kuralların izlenmesi yalnızca uyum maliyeti değil, aynı zamanda üstünlük sağlayıcı stratejik bir rekabet aracı olarak da değerlendirilmelidir.

KAYNAKÇA

Avrupa Komisyonu (European Commission). (2023-2026). Sınırda Karbon Düzenleme Mekanizması (CBAM): Resmî düzenlemeler ve uygulama rehberleri. https://taxation-customs.ec.europa.eu

Dünya Ticaret Örgütü (World Trade Organization). (2024). Ticarette Teknik Engeller Anlaşması (Agreement on Technical Barriers to Trade). https://www.wto.org/english/tratop_e/tbt_e/tbt_e.htm

Resmî Gazete. (2001, 11 Temmuz). 4703 sayılı Ürünlere İlişkin Teknik Mevzuatın Hazırlanması ve Uygulanmasına Dair Kanun (Sayı: 24459). https://www.resmigazete.gov.tr

Resmî Gazete. (2025, 9 Temmuz). 7552 sayılı İklim Kanunu (Sayı: 32951). https://www.resmigazete.gov.tr

T.C. Ticaret Bakanlığı. (2018-2026). İthalat rejimi kararları, korunma önlemi ve anti-damping tebliğleri, ülke gümrük uygulamaları rehberleri. https://ticaret.gov.tr

T.C. Ticaret Bakanlığı. (2026). Yürürlükte bulunan serbest ticaret anlaşmaları. https://ticaret.gov.tr/dis-iliskiler/serbest-ticaret-anlasmalari/yururlukte-bulunan-stalar

Türk Standardları Enstitüsü (TSE). (2026). CE işareti ve EN standartları uygulama esasları. https://www.tse.org.tr

Türkiye-Avrupa Topluluğu Ortaklık Konseyi. (1995). 1/95 sayılı Ortaklık Konseyi Kararı (Türkiye-AB Gümrük Birliği).

Türkiye İhracatçılar Meclisi (TİM). (2026). Türk ihraç ürünleri sürdürülebilirlik kodu raporu. https://tim.org.tr

Türkiye İstatistik Kurumu (TÜİK). (2018-2026). Dış ticaret istatistikleri veri tabanı. https://data.tuik.gov.tr

PANDEMİ ÖNCESİ VE SONRASINDA TÜRKİYE'DE DIŞ TİCARETİN GENEL SEYRİ

THE GENERAL COURSE OF FOREIGN TRADE IN TURKEY BEFORE AND AFTER THE PANDEMIC

Lecturer Yücel NAKTİYOK
Department of Foreign Trade, Vocational School of Social Sciences, Atatürk University, Erzurum, TÜRKİYE
ORCID ID: https://orcid.org/0000-0002-8429-1946

ÖZET

Dış ticaret, ülke ekonomisinde lokomotif görevi üstlenerek, üretim, istihdam ve büyüme alanlarına önemli katkılar sağlamaktadır. Özellikle yerli üretimi uluslararası pazarlara taşıyarak döviz kazancına doğrudan katkı sağlayan ihracat, temel makroekonomik faaliyet alanlarından biridir. Türkiye ekonomisi açısından ihracat hem büyüme dinamiklerinin hem de cari işlemler dengesini belirleyici unsurların başında gelmektedir. 2019 yılı sonlarında Çin'in Wuhan kentinde ortaya çıkan ve kısa sürede küresel bir sağlık krizine dönüşen Covid-19 pandemisi, 2020 yılı boyunca dünya ekonomilerinde önemli sorunlara yol açmış; üretimde meydana gelen daralma, lojistik ve tedarik zincirlerinde ciddi aksamalar meydana getirmiştir. Bu çalışma, Covid-19 pandemisinin Türkiye'nin dış ticaretine etkilerini 2017-2026 (ilk 5 ayı) dönemi için pandemi öncesi (2017-2019), pandemi dönemi (2020-2021) ve pandemi sonrası (2022-2026) olmak üzere üç aşamada, Türkiye İstatistik Kurumu (TÜİK), Ticaret Bakanlığı ve Türkiye İhracatçılar Meclisi (TİM) verileriyle betimsel istatistik yöntemiyle incelemektedir. Pandemi öncesinde ihracat istikrarlı biçimde artış seyri izlerken (2017: 164,4 - 2018: 177,1 - 2019: 180,8 milyar dolar), ithalatın daha yüksek olmasından kaynaklanan yapısal dış ticaret açığı devam etmiştir. 2020'de pandeminin etkisiyle ihracatta %6 civarında bir gerileme (169,6 milyar dolar) yaşanmış, ancak bu düşüş 2020 yılı ile sınırlı kalmıştır. 2021'de ihracat yüzde 33 artışla ilk kez 200 milyar doları aşarak 225,2 milyar dolara ulaşmıştır. Bu toparlanmada Türkiye'nin coğrafi konumu, e-ticaret fırsatları ve tedarik zincirlerindeki yeniden yapılanmadan (nearshoring) yararlanması etkili olmuştur. 2022-2023'te ihracat art arda rekor kırarak 254,2 ve 255,8 milyar dolara ulaşmış; ancak enerji fiyatları şoku nedeniyle ithalat da sıçrayarak dış ticaret açığı 2022'de 110 milyar dolarla 1996 sonrasının en yüksek seviyesine çıkmıştır. 2024-2025-2026 (ilk 5 ayı) yıllarında ihracat artışı sürerken (261,8 - 273,3 - 111,1 milyar dolar) ithalat artış hızı denetim altına alınarak, dış ticaret açığının kademeli olarak düşürülmesi gerçekleşmiştir. Çalışma, pandeminin Türkiye dış ticaretinde kalıcı bir yapısal kırılmadan ziyade kısa süreli bir şok etkisi yarattığını; asıl belirleyici unsurların enerji fiyatları, kur politikası ve küresel talep koşulları olduğunu ileri sürmektedir. Ayrıca firmaların e-ticaret konusunda tecrübe kazanması, pandemi sürecinin dış ticaret alanına olumlu etkisi olarak belirtilmektedir. Sonuç bölümünde, katma değeri yüksek teknolojik ürünlerin ihracatının artırılması, enerjide yerli ve yenilenebilir kaynaklara yönelim, dış pazar çeşitlendirmesi ve e-ihracatın desteklenmesi önerilmektedir.

Anahtar Kelimeler: Dış Ticaret, Covid-19 Pandemisi, İhracat, İthalat, Türkiye Ekonomisi

ABSTRACT

Foreign trade serves as a driving force in a country's economy, making significant contributions to production, employment, and economic growth. In particular, exports, which directly contribute to foreign exchange earnings by bringing domestic production to international markets, constitute one of the fundamental macroeconomic activities. For the Turkish economy, exports are among the primary determinants of both growth dynamics and the current account balance. The Covid-19 pandemic, which emerged in Wuhan, China, at the end of 2019 and rapidly evolved into a global health crisis, caused major disruptions to world economies throughout 2020, leading to a contraction in production as well as serious disruptions in logistics and supply chains. This study examines the effects of the Covid-19 pandemic on Türkiye's foreign trade over the 2017–2026 period (first five months of 2026), using descriptive statistical methods and data from the Turkish Statistical Institute (TÜİK), the Ministry of Trade, and the Turkish Exporters Assembly (TİM). The analysis is conducted in three phases: the pre-pandemic period (2017–2019), the pandemic period (2020–2021), and the post-pandemic period (2022–2026). Before the pandemic, exports followed a steady upward trend (2017: USD 164.4 billion; 2018: USD 177.1 billion; 2019: USD 180.8 billion), while the structural foreign trade deficit persisted due to imports remaining at higher levels. In 2020, exports declined by approximately 6% (USD 169.6 billion) as a result of the pandemic; however, this decrease was limited to that year. In 2021, exports increased by 33%, surpassing USD 200 billion for the first time and reaching USD 225.2 billion. This recovery was supported by Türkiye's geographical location, the opportunities created by e-commerce, and the country's ability to benefit from the restructuring of global supply chains through nearshoring. In 2022 and 2023, exports reached consecutive record levels of USD 254.2 billion and USD 255.8 billion, respectively. However, due to the energy price shock, imports also surged, causing the foreign trade deficit to reach USD 110 billion in 2022, its highest level since 1996. During 2024, 2025, and the first five months of 2026, exports continued to increase (USD 261.8 billion, USD 273.3 billion and USD 111,1 billion, respectively), while the growth rate of imports was brought under control, resulting in a gradual reduction of the foreign trade deficit. The study argues that the pandemic created a short-term shock rather than a permanent structural break in Türkiye's foreign trade. It further suggests that the main determining factors have been energy prices, exchange rate policies, and global demand conditions. In addition, the experience gained by firms in e-commerce is identified as one of the positive impacts of the pandemic on foreign trade. In the conclusion, the study.

Keywords: Foreign Trade, COVID-19 Pandemic, Exports, Imports, Turkish Economy

.

1. GİRİŞ

Dış ticaret, üretim, istihdam ve büyümeye doğrudan etki eden en önemli unsurdur. İhracat, yerli üretimi dünya pazarlarına taşıyarak üretim artışı ile birlikte istihdamı da artırıp, ekonomiye doğrudan döviz girdisi sağlar. Türkiye ekonomisi için ihracat, hem büyüme dinamiklerinin hem

de cari dengenin en önemli göstergesidir. Bu nedenle dış ticarette yaşanan ani şoklar, ülke içerisinde istikrarı makro düzeyde tehdit eder.

2019 yılının son çeyreğinde Çin'in Wuhan kentinde ortaya çıkan ve kısa sürede dünya genelinde bir sağlık krizine dönüşen Covid-19 pandemisi, 2020 yılı boyunca küresel ekonomilerinde eşi görülmemiş bir daralmaya yol açmış; üretim, lojistik ve tedarik zincirlerinde ciddi aksamalar meydana gelmiştir (AA, 2020).

Türkiye, dışa açık ekonomik politika izlemesi nedeniyle bu küresel krizden önemli oranda etkilenmiş; özellikle 2020 yılı ortalarına doğru ülke sınırlarının kapanması, talep daralması ve üretimin durma noktasına gelmesi nedeniyle ihracat gelirlerinde belirgin bir gerileme yaşanmıştır (TÜİK, 2025). Ancak aynı süreç, Türkiye'nin coğrafi konumu, esnek üretim yapısı ve tedarik zincirlerinde yaşanan yeniden yapılanmadan (nearshoring) faydalanarak 2021 yılından itibaren hızlı bir toparlanma sergilemesine de zemin hazırlamıştır (TİM, 2021).

Bu çalışmanın temel amacı, pandeminin Türkiye dış ticaretini kısa ve uzun vadede nasıl etkilediğini sorgulayarak, pandemi öncesi (2017-2019), pandemi dönemi (2020-2021) ve pandemi sonrası (2022-2026) olmak üzere üç ayrı dönemde Türkiye'nin dış ticaret performansını ihracat, ithalat, dış ticaret dengesi ve ihracatın ithalatı karşılama oranı göstergeleri üzerinden karşılaştırmalı biçimde ortaya koymaktır. Çalışma, konuya ilişkin literatürü kısaca özetledikten sonra resmî istatistiklere dayalı verileri sunmakta ve elde edilen bilgiler ışığında politika önerileri geliştirmektedir.

2. KAVRAMSAL ÇERÇEVE: DIŞ TİCARET VE PANDEMİ ETKİSİ

David Ricardo'nun dış ticaret teorisi: "ülkelerin karşılaştırmalı üstünlüklere sahip oldukları mal ve hizmetlerde uzmanlaşarak uluslararası ticaretten karşılıklı fayda sağlayacağını" öngörmektedir. Ancak pandemi gibi öngörülemeyen ani şoklar, klasik dış ticaret kurallarını kısa vadede bozarak aynı anda arz ve talep yönlü daralmalara yol açabilmektedir. Küresel salgın, üretim faktörlerinin serbest dolaşımını kısıtlamış, taşımacılık maliyetlerini artırmış ve firmaların tedarik zinciri stratejilerini yeniden gözden geçirmelerine neden olmuştur (TCMB, 2025).

Literatürde pandeminin dış ticaret üzerindeki etkileri üç temel başlık altında ele alınmaktadır: (1) arzda azalma, üretim tesislerinin geçici kapanması ve işgücü kayıplarını; (2) talepte azalma, tüketici harcama eğilimindeki azalmayı; (3) lojistik ve taşımacılık alanında ise sınır kapatmaları ve konteyner temindeki zorlukları ifade etmektedir. Türkiye özelinde ise, imalat sanayinin ihracat içindeki yüksek payı (%93-94 bandı) nedeniyle pandeminin etkisi büyük ölçüde sanayi üretimindeki dalgalanmalar üzerinden dış ticarete yansımıştır (T.C. Ticaret Bakanlığı, 2025).

3. ARAŞTIRMANIN AMACI VE YÖNTEMİ

Bu çalışmada nicel araştırma yaklaşımlarından betimsel (tanımlayıcı) istatistik yöntemi kullanılmıştır. Araştırmanın veri seti, Türkiye İstatistik Kurumu (TÜİK), Ticaret Bakanlığı ve Türkiye İhracatçılar Meclisi (TİM) tarafından Genel Ticaret Sistemi (GTS) esas alınarak yayımlanan 2018-2025 yıllarına ait yıllık ihracat, ithalat, dış ticaret dengesi ve ihracatın ithalatı karşılama oranı verilerinden oluşmaktadır. Veriler, pandemi öncesi (2018-2019), pandemi

dönemi (2020-2021) ve pandemi sonrası (2022-2026) olarak üç döneme ayrılmış; dönemler arası karşılaştırmalı analiz yapılarak eğilimler yorumlanmıştır (TÜİK, 2026; T.C. Ticaret Bakanlığı, 2025; TİM, 2021, 2024). Çalışma, ikincil veri kaynaklarına dayanması nedeniyle betimsel bir nitelik taşımakta olup nedensellik ilişkisi kurma iddiası taşımamaktadır. Üç farklı kaynaktan elde edilen bulguların tutarlılığı, bulguların güvenilirliğini önemli ölçüde göstermektedir.

4. PANDEMİ ÖNCESİ DÖNEMDE TÜRKİYE DIŞ TİCARETİ (2017-2019)

Bu bölüm, krizden önceki durumu sayısal verilerle tanımlamaktadır. 2017–2019 arası, Türkiye ihracatının istikrarlı bir yükseliş gösterdiği dönemdir. Bu üç yıl, pandemi sonrası etkinin gerçek büyüklüğünü ölçmek için önemli bir veri oluşturur. Pandemi öncesi dönemde Türkiye'nin dış ticareti istikrarlı ancak yapısal açık sorunu devam eden bir görünüm sergilemiştir.

2017 yılında Türkiye ihracatı önceki yıla göre yüzde 10,2 oranında artış göstererek 164,5 milyar dolar olmuş, bu rakam izleyen yılların büyüme trendine ivme kazandırmıştır. Ancak aynı yıl ithalat değeri de yüzde 18,1 oranında artış göstererek geçmiş yıllarda olduğu gibi ihracatın sürekli üzerinde seyretmeye devam etmiştir.

2018 yılında ihracat yüzde 7,7 oranında büyüme göstererek 177,1 milyar dolar düzeyinde gerçekleşmiş, aynı yıl yaşanan aşırı kur şokları sonucu ithalata meydana gelen baskılama, ithalatı yüzde 3,2 oranında azaltarak dış ticaret açığını 53,9 milyar dolara düşürmüştür. İhracatta süregelen artış, küresel talep koşullarının henüz uygun olduğu bir dönemi göstermektedir. Kur dalgalanmalarına rağmen ihracatçı firmalar rekabet gücünü korumayı becerebilmiş, böylece istikrarlı büyüme çizgisi, ikinci yılında da bozulmadan devam etmiştir.

2019 yılında, ihracat pandemi öncesi dönemin rekor seviyesi olmuştur. Bu zirve, 2020'deki düşüşün gerçek büyüklüğünü ölçmek için önemli bir gösterge teşkil etmektedir. Aynı yıl iç talepteki azalma ve ekonomik durgunluğun etkisiyle ithalat yüzde 9,0 oranında gerileyerek 210,3 milyar dolara inmiş, buna karşın ihracat yüzde 2,1 artışla 180,8 milyar dolara yükselmiştir. Bu gelişmeler sonucunda dış ticaret açığı 2019 yılında 29,5 milyar dolara inmiş, ihracatın ithalatı karşılama oranı ise yüzde 86 ile son yılların en yüksek seviyesine ulaşmıştır (TÜİK, 2026; T.C. Ticaret Bakanlığı, 2026). 2019 yılındaki bu iyileşme, esasen iç talep azalmasına bağlı olup, ihracat artışından çok ithalattaki şok düşüşten kaynaklanmıştır. Dolayısıyla pandemi öncesi dönemin sonunda Türkiye dış ticareti, ihracat açısından büyütme seyri izlerken, ithalatta enerji ve ara malı bağımlılığı devam eden bir yapıda pandemi ile yüzleşmiştir. Birkaç ay sonra küresel bir krizin ortaya çıkacağını o dönemde kimse öngörememiştir.

2017–2019 yıllarında ihracatta gerçekleşen kesintisiz artış, ihracat sektörünün istikrarını açıkça göstermiş, ancak ihracattaki istikrarlı artışa rağmen, ithalat her yıl daha yüksek seviyede gerçekleşmiştir. Bu fark, Türkiye ekonomisinin süregelen yapısal dış ticaret açığını devam ettirmiştir. Dış ticaret açığının temel kaynağı, Türkiye'de enerji ve ara malı ithalatına duyulan bağımlılıktır.

5. PANDEMİ DÖNEMİNDE TÜRKİYE DIŞ TİCARETİ (2020-2021)

Bu bölümde, ihracatı pandeminin doğrudan etkilediği ve ardından beklenmeyen hızda gerçekleşen bir toparlanmanın yaşandığı iki yıl ele alınmakta. 2020, ihracatın gerilediği yıl olmasına rağmen 2021 ise umulmadık bir sıçramayla tüm karamsar beklentileri tersine çevirmiştir. Bu durum, pandeminin ihracat üzerinde oluşturduğu şok etkisinin kısa süre içerisinde giderildiğini göstermektedir.

Covid-19 pandemisinin Türkiye ekonomisi üzerindeki etkileri Mart 2020'den itibaren belirgin biçimde hissedilmeye başlamıştır (AA, 2020). Sokağa çıkma kısıtlamaları, ülke sınırlarının kapatılması ve küresel talep daralması nedeniyle, 2020 yılı genelinde ihracat, bir önceki yıla göre yüzde 6,2 oranında azalarak 169,6 milyar dolara gerilerken, ithalat 219,5 milyar dolar düzeyinde gerçekleşmiş ve dış ticaret açığı 49,8 milyar dolara yükselmiştir (TÜİK, 2026; T.C. Ticaret Bakanlığı, 2026). Bu düşüş, üç yıldır süregelen büyüme trendini ani olarak sonlandırmıştır.

Pandemi etkilerinin en yoğun yaşandığı 2020 yılını daha detaylı irdeleyecek olursak:

2020 yılının ilk altı ayında ihracat 32,4 milyar dolara kadar gerilemiş; özellikle Nisan 2020'de aylık ihracat 8,97 milyar dolar ile son yılların en düşük seviyesini görmüştür. O günlerde kritik soru şuydu: bu düşüş kalıcı bir seyir mi izleyecek, yoksa geçici bir durum mu? 2020 yılının yaz aylarından itibaren normalleşme adımlarıyla birlikte güçlü bir toparlanma yaşanmış; ihracat haziran – ağustos döneminde önceki üç aya göre yüzde 33,8 oranında artarak 43,3 milyar dolara ulaşmıştır. Özellikle hazır giyim, tekstil ve otomotiv sektörleri Uzak Doğu'dan Türkiye'ye yönelen sipariş akışından faydalanarak tedarik zincirlerinde toparlanmanın öncüsü olmuştur. Aralık 2020'de aylık ihracat 17,8 milyar dolar ile o güne kadarki en yüksek aylık ihracat rakamına ulaşmıştır (T.C. Ticaret Bakanlığı, 2025).

2021 yılı, pandeminin ekonomik etkilerinin büyük ölçüde atlatıldığı ve dış ticarette tarihi bir rekorun yaşandığı yıl olmuştur. İhracat, bir önceki yıla göre yaklaşık yüzde 33 oranında artarak ilk kez 200 milyar dolar sınırını aşmış ve 225,2 milyar dolara ulaşmıştır. İthalat da 271 milyar dolara yükselmiş, dış ticaret açığı ise 46 milyar dolar civarında gerçekleşmiştir. Bu dönemde küresel tedarik zincirlerindeki aksaklıklar, konteyner taşımacılığındaki sorunlar ve hammadde fiyatlarındaki yükseliş, ithalat maliyetlerini artıran temel unsurlar olmuştur (TÜİK, 2026; TİM, 2021). Bu hızlı toparlanma, pandeminin kalıcı etki oluşturmadığının somut delilidir. Beklenmeyen bu artış, literatürde 'V tipi toparlanma' adıyla anılan ekonomik terimin en güzel göstergesidir. İhracattaki bu kalıcı yükseliş, pandeminin geçici bir şok olduğu tezini güçlü biçimde desteklemektedir.

Türkiye ihracatında 2021 yılında gerçekleşen toparlanmanın en önemli nedeni: krizin kazandırdığı fırsattır (nearshoring). Dünya ticaretinin en önemli tedarikçileri olan Çin ve uzak doğu ülkeleri, pandemi sırasında lojistik sorunlar, kapanma vb. nedenlerle Avrupa ülkeleri başta olmak üzere pek çok ülkeye mal tedariki sağlayamamıştır. Bu durum özellikle Avrupa ülkelerini, üretim ve tedarik konusunda coğrafi olarak daha yakın ülkelere yönlendirmiş, Türkiye, gümrük birliğinin kazandırdığı birikim, Avrupa'ya yakınlığı sonucu teslimat hızı konusundaki avantajı ve güçlü sanayi altyapısıyla bu arayışın en büyük adaylarından biri

olmuştur. Bunun sonucunda Avrupa ve diğer ülkelerin küresel tedarik zincirlerini yeniden yapılandırmaya gitmesi Türkiye açısından pandemiyi fırsata dönüştürmüştür.

6. PANDEMİ SONRASI DÖNEMDE TÜRKİYE DIŞ TİCARETİ (2022-2026)

Geçmiş yıllarda olduğu gibi pandemi sonrası dönemde de Türkiye'nin ihracat performansı yükseliş eğilimini sürdürmüştür.

2022 yılında ihracat, bir önceki yıla göre yüzde 12,9 oranında artarak ilk kez 250 milyar dolar sınırını aşmış ve 254,2 milyar dolarla tarihi bir rekora imza atmıştır. Ancak rekor ihracata rağmen, dış ticaret açığı 1996 sonrası dönemin en yüksek seviyesi olan 109,5 milyar dolara ulaşmıştır. Dış açığı tetikleyen en önemli faktör olan Rusya-Ukrayna savaşının enerji fiyatlarını yükseltmesi sonucu, ithalat yüzde 34,3 oranında artarak 363,7 milyar dolara ulaşmıştır (TÜİK, 2025; T.C. Ticaret Bakanlığı, 2025; TCMB, 2025). Bu rekor açık, ihracatta kaydedilen artışın ithalat karşısında yetersiz kaldığını kanıtlamış, açığın büyüklüğü, cari işlemler dengesi üzerinde ciddi bir baskı oluşturmuştur. Bu veri, çalışmanın en çarpıcı ve dikkat çeken bulgularından biri olarak öne çıkmaktadır.

2023 yılında ihracat, sınırlı bir artışla 255,8 milyar dolara yükselerek Cumhuriyet tarihinin en yüksek yıllık ihracat rakamına ulaşmış (AA, 2024); ithalat ise yüzde 0,5 oranında küçük bir düşüşle 361,9 milyar dolara inmiştir. Bu gelişmeyle dış ticaret açığı 106 milyar dolar civarına gerilemiştir.

2024 yılında ise dış ticaret dengesinde belirgin bir iyileşme kaydedilmiş; ihracat 261,9 milyar dolara ulaşırken, ithalat yüzde 5 oranında gerileyerek 344 milyar dolara inmiştir. Böylece dış ticaret açığı yüzde 22,7 oranında azalarak 82,2 milyar dolara gerilemiş, ihracatın ithalatı karşılama oranı ise yüzde 76,1 ile 2002 sonrası dönemin en yüksek seviyelerinden birine ulaşmıştır (TÜİK, 2025; SBB, 2025; İKV, 2025; TİM, 2024). 2024 yılında ihracat büyüme çizgisini korurken, ithalat artış hızı, önceki yıllara kıyasla ilk kez kontrol altına alınmaya başlanmıştır. Sonuç olarak, dış ticaret açığındaki büyüme ivmesi de yavaşlamaya başlamıştır. Bu denge arayışı, ekonomi yönetiminin önceliklerini değiştirdiğinin açık bir işaretidir.

Türkiye İstatistik Kurumu verileri 2025 yılında da ihracatta yükselişin sürdüğüne işaret etmektedir. İhracat, bir önceki yıla göre yüzde 4,4 oranında artarak 273,3 milyar dolarla tarihinin en yüksek seviyesine ulaşmıştır. Bu rakam, 2019'daki pandemi öncesi zirvenin neredeyse yüzde 51 üzerinde bulunmaktadır. İthalat ise yüzde 6,2 oranında artışla 365,4 milyar dolar olarak gerçekleşmiştir (TÜİK, 2025). Bu gelişmeler sonucunda dış ticaret açığı yaklaşık 92 milyar dolar olup, ihracatın ithalatı karşılama oranı yüzde 74,8 civarında kalmıştır. Bu dönemde AB ülkeleri, Amerika Birleşik Devletleri, Birleşik Krallık, Irak ve İtalya Türkiye'nin başlıca ihracat pazarları olmaya devam ederken; ithalatta Çin ve Rusya ilk sıralardaki yerlerini korumuştur (T.C. Ticaret Bakanlığı, 2025; AA, 2025).

2026 yılının ilk beş ayında ihracat 111,1 milyar dolara ulaşmış, ithalat ise 153,8 milyar dolar olarak gerçekleşmiştir. Bu durum, yıl sonunda önceki yılın rekor rakamının aşılma potansiyeline işaret etmektedir. Yükseliş trendinin sürmesi, ihracat artışının konjonktürel değil

kalıcı olduğu tezini desteklemektedir. Bu bölüm, krizin ardından gelen art arda ihracat rekorları ve yeni dengeleri ortaya koymaktadır.

7. KARŞILAŞTIRMALI DEĞERLENDİRME

Tablo 1'de sunulan 2017-2026 dönemi verileri, Türkiye dış ticaretinin pandemi öncesinde görece durağan, pandemi döneminde şok bir düşüş ve ardından hızlı bir toparlanma, pandemi sonrasında ise rekor düzeyde ihracat artışına rağmen yapısal dış ticaret açığının sürdüğü bir seyir izlediğini ortaya koymaktadır.

Yıl	İhracat (milyar USD)	İthalat (milyar USD)	Dış Ticaret Dengesi (milyar USD)	İhr./İth. Karşılama Oranı (%)
2017	164,5	238,7	-74,2	68,9
2018	177,1	231,1	-53,9	76,6
2019	180,8	210,3	-29,5	86,0
2020	169,6	219,5	-49,8	77,3
2021	225,2	271,4	-46,2	83,0
2022	254,2	363,7	-109,5	69,9
2023	255,6	361,9	-106,3	70,6
2024	261,7	344,0	-82,2	76,1
2025	273,2	365,4	-92,1	74,8
2026 *	111,1	153,8	-42,7	72,2

Kaynak: TÜİK (2026), (2026 yılı ocak - mayıs verileri)*

Tablodan da izlenebileceği üzere, 2020 yılı ihracat rakamı (169,6 milyar dolar) 2017-2019 dönemine kıyasla belirgin bir gerileme sergilemiş, ancak bu düşüş 2019'a göre yalnızca yüzde 6,3 ile sınırlı kalmıştır. Bunun temel nedeni, pandeminin ilk şokunun yılın yalnızca ikinci çeyreğinde yoğunlaşması ve yılın geri kalanında güçlü bir toparlanmanın yaşanmasıdır. 2021-2023 döneminde ihracat kesintisiz artış göstererek 225 milyar dolardan 255,8 milyar dolara yükselmiş; bu süreçte Türkiye, küresel tedarik zincirlerinin yeniden yapılandığı bir ortamda coğrafi yakınlık avantajını değerlendirebilmiştir (TÜİK, 2025; T.C. Ticaret Bakanlığı, 2025).
Öte yandan ithalat tarafında, özellikle 2021-2022 döneminde yaşanan sıçrama (271 milyar dolardan 364,4 milyar dolara), büyük ölçüde enerji fiyatlarındaki artış ve kurdaki oynaklıktan kaynaklanmıştır. Bu durum, dış ticaret açığının 2022 yılında tarihi bir zirveye ulaşmasına yol açmıştır. İhracat 2022'den itibaren her yıl yeni bir rekora ulaşmış, ancak enerji fiyatlarındaki aşırı yükseliş, aynı dönemde dış ticaret açığını da tarihi seviyelere taşımıştır. Bu bölüm, başarı ile kırılganlığın nasıl iç içe geçtiğini ortaya koymaktadır.

2024 ve 2026 yıllarında ise ithalat artış hızı, ihracat artış hızının belirgin biçimde gerisinde kalması sonucu dış ticaret açığında kademeli bir daralma gözlenmektedir. Kur politikası, iç piyasaya yönelim ve talep yönetimi, ithalatta düşüş sağlanmasında belirleyici rol oynamıştır.
Bu bulgular, pandeminin Türkiye dış ticaretinde kalıcı bir yapısal kırılmadan çok, kısa süreli bir şok etkisi yarattığını; asıl belirleyici unsurların enerji fiyatları, kur politikası ve küresel talep koşulları olduğunu göstermektedir (TCMB, 2025; TÜİK, 2025).



Grafik, on yıllık ihracat seyrini özetlemektedir. 2020'deki düşüş, çizgide küçük ama belirgin olarak görülmekte, 2021 sonrasında ise kesintisiz ve dik bir yükseliş mevcuttur. 2026 yılındaki düşüş ise rakamların sadece ilk beş ayı kapsamasıdır.
Bu grafik, 'kısa süreli şok, uzun süreli büyüme' düşüncesinin ispatı niteliğindedir. Veriler, pandeminin Türkiye dış ticaretinde kalıcı bir etki yapmadığını açıkça göstermektedir. Etki, yalnızca 2020 yılıyla sınırlı kalan kısa süreli bir şok niteliği taşımaktadır. 2021'den itibaren ihracat, yükselme trendine devam etmiştir.

8. SONUÇ VE ÖNERİLER

Bu çalışmada Türkiye'nin 2017-2026 dönemine ait dış ticaret verilerini pandemi öncesi, pandemi dönemi ve pandemi sonrası olmak üzere üç aşamada incelenmiştir. Bulgular, Covid-19 pandemisinin Türkiye dış ticaretinde geçici olmakla birlikte etkili bir şok oluşturduğunu ortaya koymuştur. 2020 yılında Türkiye ihracatında sınırlı fakat belirgin yaşanan daralma, 2021 yılından itibaren güçlü bir toparlanma sürecine yerini bırakmıştır. 2021 sonrası sürekli hale gelen yükseliş trendi, ekonominin krizden güçlenerek çıktığını açıkça kanıtlamaktadır. Özetle Türkiye dış ticareti, pandemiyi bir çöküş değil, dönüm noktası olarak geride bırakmış, krizin ardından ekonomisi daha istikrarlı bir büyüme rotasına girmiştir.

Pandemi döneminin dünya ekonomisi üzerindeki etkilerini özetleyecek olursak: üretim zincirleri, hammadde ve ara malı tedarikinde yaşanan aksamalar sonucu durma noktasına gelmiş, sınırların kapatılması ve karantina uygulamaları, uluslararası taşımacılığı önemli oranda sekteye uğratmıştır. Talep yetersizliği ve lojistik sorunlar firmaların yatırım ve sipariş kararlarını derinden etkilemiş, konteyner temininde yaşanan zorluklar ve navlun fiyatlarındaki aşırı yükselme, ihracatçıların maliyet yapısını oldukça bozmuştur. Limanlarda oluşan birikmeler, teslimat sürelerini geciktirerek, ticareti öngörülemez hale getirmiştir. Özellikle ana tedarikçi konumunda olan ülkelerde üretim hatları zaman zaman durmuş, bu aksaklıklar, firmaları tedarik zincirlerini yeniden gözden geçirmeye zorunlu kılmıştır.

Türkiye'nin 2022 ve 2023 yıllarında art arda ihracat rekorları kırması, küresel tedarik zincirlerindeki yerinin yeniden konumlanmasının olumlu etkilerine işaret etmektedir. Buna karşılık, enerji ithalatına olan yapısal bağımlılık, dış ticaret açığının pandemi sonrası dönemde de yüksek seviyelerde seyretmesine neden olmuştur.

2024-2026 dönemine ait veriler ise ihracat artışının sürdüğü ve dış ticaret açığının kademeli olarak daraldığı bir sürece işaret etmektedir. Aynı dönemde Rusya-Ukrayna savaşına ek olarak ABD-İsrail-İran savaşı dünya enerji piyasaları için tarihi bir dalgalanma dönemine sebep olmuş, enerjide dışa bağımlı olan Türkiye, bu savaşlardan doğrudan etkilenerek ithalat giderlerinde önemli artışlar meydana gelmiştir.

Pandemi, Türkiye ve dünya genelinde e-ticaret işlemlerinin önemini ortaya çıkararak, toplam ihracat içindeki payını hızla artırmıştır. Firmalar, yurt dışı seyahat kısıtlamaları sonucu, fuar vb. etkinliklerden mal tedariki yerine dijital kanalları kullanmayı tercih etmiştir. E-ticarette edinilen tecrübe pandemi sonrası dönemde de kurumsal yetkinliğe dönüşmüştür. Bu durum, pandeminin dış ticarete kazandırdığı olumlu katkılardan biridir.

Bu çerçevede, Türkiye'nin dış ticaret dengesinin kalıcı biçimde iyileştirilebilmesi için üretim artışı ile birlikte katma değeri yüksek ve teknoloji yoğun ürünlerin ihracatını destekleyici tedbirlerin alınması önemli bir husustur. İthalat giderlerinin en önemli aktörü olan enerjide yerli ve yenilenebilir kaynaklara yönelmek stratejik bir zorunluluk olup, buna ilişkin çalışmalar yoğunlaştırılmalıdır. Mevcut pazarlarda ürün yelpazesini genişletmek, aynı zamanda yeni pazarlara giriş çabaları desteklenerek dış pazarların çeşitlendirilmesi gerekmektedir. Pandemi döneminde önemli bir ivme kazanan ve yükseliş trendi devam eden ve e-ihracat konusunda firmalar daha çok desteklenerek dijital ticaret kanallarının geliştirilmesi önem taşımaktadır. Özellikle KOBİ'lere e-ticaret konusunda yeni ihracat kanalları açılmalıdır. Turquality Programı kapsamında sunulan destekler artırılarak "Türk Malı" imajının dünya geneline yayılması markalaşma konusunda Türkiye'nin rekabet gücünü artırarak, gelecekte ortaya çıkabilecek ekonomik riskleri dağıtıp yükseliş trendini sürekli hale getirecektir. Türkiye açısından pandemi esnasında krizin kazandırdığı fırsat (Nearshoring) konjonktürü iyi değerlendirilerek Avrupa ve diğer ülkelere yönelik tedarikçi konumu korunmalı, Türkiye Avrupa ve komşu ülkeler için bir üretim üssüne dönüşmelidir. Ayrıca Ar-Ge ve inovasyon yatırımları desteklenerek, ihracat artışı ve markalaşma kalıcı hale getirilmelidir.

Bu dinamikler Türkiye için krizi gerçek bir büyüme fırsatına çeviren sinerjiyi devam ettirecek en temel unsurlardır.

Gelecek çalışmalarda, sektörel ve bölgesel veriler baz alınarak değişik ekonomik modellerin kullanılması, pandeminin dış ticaret üzerindeki etkilerinin farklı açılardan değerlendirilmesine olanak sağlayacaktır.

9. TARTIŞMA

Pandemi, ekonomi açısından kalıcı bir çöküş mü, geçici bir daralma mı?

Üç dönem halinde incelenen veriler, pandeminin Türkiye dış ticaretinde kalıcı çöküşe neden olmadığını belirgin bir şekilde göstermektedir. Pandemi ekonomi üzerindeki olumsuz etkileri 2020 yılıyla sınırlı kalan kısa süreli bir şok niteliği taşımaktadır. 2021'den itibaren ihracat, V tipi toparlanma göstererek krizden önceki yükseliş trendinin çok üzerinde ivme kazanarak yeni bir rota oluşturmuştur. Bu bulgu, çalışmanın merkezi ve aynı zamanda en tartışmaya açık iddiasını oluşturmaktadır.

Dış ticaret açığını belirleyen asıl aktör pandemi mi, ekonominin yapısal durumu mu? Türkiye'nin enerji konusunda dışa bağımlılığı dış ticaret açığı vermesinin en önemli yapısal kaynağı olup, bu bağımlılık çözüm bekleyen kritik bir sorun olarak önümüzde durmaktadır. Kriz, savaş vb. şoklar sonucu yükselen enerji fiyatları, dış ticaret dengesinin negatif yönde bozulmasının en etkili belirleyicisi olarak öne çıkmaktadır.

Diğer yanda enflasyonist nedenlerle baskılayıcı kur politikası izlenmesi ihracatçıların rekabet gücünü doğrudan şekillendiren ikinci temel faktördür. Küresel talep koşulları, Türkiye'nin ihracat performansının nihai sınırını çizmektedir. Bu üç faktör, dış ticaret açığını belirleyen unsurun pandemi değil, ekonominin yapısal durumu olduğunu göstermektedir. Böylece ihracattaki başarı, dış ticaret açığı sorununu tek başına çözmeye yetmemiştir.

Gelecekte dış açığın kapatılması ve ihracatta istenilen düzeyde büyüme, katma değeri yüksek ürünler, enerji ve pazar politikalarındaki reform ile uygun kur politikalarına sıkı sıkıya bağlıdır. Sonuç olarak bulgular, kriz dönemlerinde karamsarlık ve kaos yerine, fırsatları iyi değerlendirip, doğru kanallara yönlendirecek politika yapmanın önemini ortaya çıkarmaktadır.

KAYNAKÇA

Anadolu Ajansı (AA). (2020). *Türkiye'nin 2020 yılı ihracat verileri ve pandemi döneminde dış ticaretin seyrine ilişkin haberler*. https://www.aa.com.tr

Anadolu Ajansı (AA). (2023). *Türkiye'nin 2023 yılı ihracat rekoruna ilişkin haberler*. https://www.aa.com.tr

Anadolu Ajansı (AA). (2024). *Türkiye'nin 2024 yılı dış ticaret verilerine ilişkin haberler*. https://www.aa.com.tr

Cumhurbaşkanlığı Strateji ve Bütçe Başkanlığı (SBB). (2025). *2024 yılı Aralık ayı geçici dış ticaret verileri*. https://www.sbb.gov.tr

İktisadi Kalkınma Vakfı (İKV). (2025). *Türkiye'nin 2024 dış ticaret verilerine ilişkin değerlendirme notu*. https://www.ikv.org.tr

T.C. Ticaret Bakanlığı. (2026). *Aylık ve yıllık dış ticaret istatistikleri, 2017-2026* [Veri tabanı]. https://ticaret.gov.tr

Türkiye Cumhuriyet Merkez Bankası (TCMB). (2025). *Ödemeler dengesi, döviz kuru ve enflasyon istatistikleri*. https://www.tcmb.gov.tr

Türkiye İhracatçılar Meclisi (TİM). (2021). *2021 yılı ihracat değerlendirme raporu*. https://tim.org.tr

Türkiye İhracatçılar Meclisi (TİM). (2024). *2024 yılı ihracat değerlendirme raporu*. https://tim.org.tr

Türkiye İstatistik Kurumu (TÜİK). (2026). *Dış ticaret istatistikleri veri tabanı, 2017-2026* [Veri tabanı]. https://data.tuik.gov.tr

GÖÇ ARAŞTIRMALARINDA KÜÇÜK YERLEŞİMLERİN KURAMSAL DEĞERİ: METROPOL MERKEZLİ YAKLAŞIMLARIN ELEŞTİREL BİR DEĞERLENDİRMESİ

THE THEORETICAL VALUE OF SMALL SETTLEMENTS IN MIGRATION RESEARCH: A CRITICAL ASSESSMENT OF METROPOLIS-CENTRED APPROACHES

Prof. Dr. Metin AKSOY
Gümüşhane University, International Relations
ORCID:0000-0002-3118-6739

Muhlise CANLI
Lect., Gümüşhane University, Vocational School of Social Sciences,
Department of Social Work and Counseling
ORCID:0000-0001-6051-4677

Özet

Göç araştırmalarında büyük kentlerin belirgin bir ağırlığı bulunmaktadır. Bunun anlaşılabilir nedenleri vardır. Göç hareketlerinin yoğunluğu, etnik ağların çeşitliliği, emek piyasalarının büyüklüğü ve göçle ilgilenen kurumların çoğunlukla bu kentlerde toplanması, metropolleri araştırmacılar açısından önemli hâle getirmiştir. Buna karşılık küçük yerleşimler, göçmen nüfusun daha az olduğu ve göçün etkilerinin sınırlı kaldığı yerler şeklinde değerlendirilmiş, çoğu zaman araştırmanın dışında bırakılmıştır. Oysa bir yerleşimin küçük olması, göç süreçlerini anlamak bakımından daha az açıklayıcı olduğu anlamına gelmez. Bu bildiride küçük yerleşimler metropollerle karşılaştırılmamaktadır. Tartışma daha çok, bu yerleşimlerdeki toplumsal ilişkilerin göç araştırmalarında hangi meseleleri görünür kılabileceği üzerinde yürütülmektedir.

Küçük yerleşimlerin önemi, yalnızca daha az sayıda göçmene ev sahipliği yapmalarından kaynaklanmamaktadır. Bu tür yerlerde göçmenlerle yerel toplum üyeleri aynı mekânlarda daha sık karşılaşabilmekte ve birbirlerini zamanla tanıyabilmektedir. Fakat tanınırlık her zaman yakınlık veya kabul üretmez. Tekrarlanan karşılaşmalar bir taraftan gündelik ilişkiyi sürdürürken, diğer taraftan sosyal mesafenin ve yerel aidiyet sınırlarının korunmasına da hizmet edebilir. Kurumsal kapasitenin sınırlı olduğu durumlarda ise taraflar karşılaştıkları sorunları çoğu zaman formel mekanizmaların dışında geliştirdikleri pratiklerle çözmeye çalışmaktadır. Yakınlık, tanınırlık, yerel kimlik, enformel yönetişim ve sessiz uyum gibi kavramlar küçük yerleşimlerdeki bu ilişkiler üzerinden farklı biçimlerde düşünülebilir.

Bildiri, küçük yerleşimlerin göç araştırmalarında yalnızca tamamlayıcı saha alanları olarak görülmesi yaklaşımına katılmamaktadır. Bu yerleşimlere yakından bakıldığında göçün genellikle arka planda kalan yerel ve gündelik yönleri daha açık hâle gelmektedir. Göçmenlerle yerel toplum üyelerinin mesafeyi nasıl korudukları, birlikte yaşamanın yazılı olmayan kurallarını nasıl öğrendikleri ve yerel sınırları hangi durumlarda esnettikleri bu alanlarda izlenebilmektedir. Küçük yerleşimlerin bu açıdan göçü farklı ölçeklerde ele alan ve yerel ilişkileri analizin merkezine yerleştiren bir araştırma yaklaşımına katkı sağlayabileceği düşünülmektedir.

Anahtar Kelimeler: Göç araştırmaları, küçük yerleşimler, yerellik, sosyal mesafe, gündelik karşılaşma, sessiz uyum.

Abstract
Large cities have occupied a prominent place in migration research. There are understandable reasons for this. The intensity of migration movements, the diversity of ethnic networks, the size of labour markets, and the concentration of migration-related institutions in these cities have made metropolitan areas important research sites. Small settlements, by contrast, have often been viewed as places with smaller migrant populations and limited migration effects. As a result, they have frequently remained outside the scope of research. Yet the small size of a settlement does not mean that it offers less insight into migration processes. This paper does not compare small settlements with metropolitan areas. Instead, it discusses which aspects of migration can become visible through the social relations that develop within these settlements. The significance of small settlements does not arise solely from the fact that they accommodate fewer migrants. In such places, migrants and members of the local community may encounter one another more frequently in the same spaces and become familiar with each other over time. Recognition, however, does not always lead to closeness or acceptance. Repeated encounters may sustain everyday relationships while also helping to preserve social distance and the boundaries of local belonging. Where institutional capacity is limited, the parties often attempt to address the problems they encounter through practices developed outside formal mechanisms. Concepts such as proximity, recognition, local identity, informal governance, and silent adaptation can therefore be reconsidered through these relationships in small settlements. The paper does not share the view that small settlements should be treated merely as supplementary field sites in migration research. A closer examination of these places makes the local and everyday dimensions of migration, which often remain in the background, more visible. They provide opportunities to observe how migrants and members of the local community maintain distance, learn the unwritten rules of living together, and determine when local boundaries may be relaxed. From this perspective, small settlements may contribute to a research approach that examines migration across different scales and places local relations at the centre of analysis.
Keywords: Migration research, small settlements, locality, social distance, everyday encounters, silent adaptation.

Giriş
Bu çalışma, küçük yerleşim yerlerini metropollerle doğrudan karşılaştırarak tartışmamaktadır. Böyle bir karşılaştırma belirli bir risk barındırmaktadır. Küçük yerleşim yerleri bu durumda kolaylıkla "eksik" olarak değerlendirilebilmektedir. Ancak çalışmanın amacı bu değildir. Küçük yerleşim yerleri, büyük kentlerin zayıf veya sınırlı biçimleri olarak anlaşılmamaktadır. Aynı zamanda yalnızca göç yoğunluğunun daha düşük olduğu mekânlar olarak da ele alınmamaktadır. Küçük yerleşim yerleri kendilerine özgü toplumsal ilişkilere ve karşılaşma biçimlerine sahiptir. Ayrıca yerel sınırları, enformel çözüm yolları ve özgün görünürlük biçimleri bulunmaktadır. Bu özellikleriyle küçük yerleşim yerleri, göç araştırmaları açısından farklı bir kavramsal temel sunmaktadır.
Bu çalışmanın temel sorusu şu şekilde formüle edilmektedir: Hangi toplumsal süreçler küçük yerleşim yerlerini göç araştırmalarında görünür hâle getirmektedir? Bu süreçler, mevcut göç kavramlarının yeniden düşünülmesine nasıl katkı sağlayabilir? Bu soru, ampirik vakaların doğrudan karşılaştırılmasına dayanmamaktadır. Soru, literatür temelinde kavramsal olarak tartışılmaktadır. Çalışma açık bir tez ileri sürmektedir. Küçük yerleşim yerleri göç araştırmalarında yalnızca ek araştırma alanları olarak değerlendirilmemelidir. Bu yerleşim yerleri güçlü bir teorik potansiyel sunmaktadır. Yerel kabul, toplumsal mesafe ve gündelik karşılaşmalar bu yerleşim yerlerinde daha ayrıntılı biçimde incelenebilmektedir. Aynı durum aidiyet sınırları ve enformel yönetişim biçimleri için de geçerlidir.

Küçük yerleşim yerlerinde göçmen kadın ve erkekler ile yerel nüfusun üyeleri tekrar tekrar karşılaşmaktadır. Bu karşılaşmalar sokaklarda, iş yerlerinde ve okullarda gerçekleşmektedir. Pazarlar, duraklar ve yerel kurumlar da önemli karşılaşma alanlarını oluşturmaktadır. Bu nedenle küçük yerleşim yerleri, göç açısından özgün bir gözlem alanı hâline gelmektedir. Bununla birlikte bu tür karşılaşmalar otomatik olarak entegrasyona, yakınlığa veya kabule yol açmamaktadır. Karşılıklı tanışıklık ve toplumsal görünürlük, bazı durumlarda güveni güçlendirebilmektedir. Diğer durumlarda ise toplumsal denetimi artırabilmekte, yerel normlara uyma yönündeki baskıyı yükseltebilmekte ve mesafeli ilişkilerin oluşmasını teşvik edebilmektedir. Bu nedenle küçük yerleşim yerlerini göçün "daha sakin" biçimde gerçekleştiği mekânlar olarak sunmak doğru değildir. Aynı şekilde, sınırlı kurumsal kapasiteleri nedeniyle bu yerleşim yerlerini daha az önemli kabul etmek de doğru değildir.

Literatürde Yerellik ve Ölçek Sorunu

Yerellik meselesi, göç araştırmaları açısından teorik önem taşımaktadır. Glick Schiller ve Çağlar'ın çalışmaları bu konuda önemli bir başlangıç noktası sunmaktadır. Yazarlar, göç araştırmalarının uzun süre etnik gruplara, ulus-devletlere ve göçmen topluluklarına odaklandığını ortaya koymaktadır. Buna karşılık göçmenler ile ikamet ettikleri yerler arasındaki karşılıklı ilişki yeterince teorik olarak incelenmemiştir. Yazarlara göre kentler ve yerel mekânlar, göçmenlerin basitçe yerleştiği boş kaplar değildir. Göçmenlerin bu mekânlara dâhil olma biçimleri, ilgili yerin ekonomik, siyasal ve toplumsal konumuyla birlikte ele alınmalıdır. Bu yaklaşım, küçük yerleşim yerleri açısından özellikle önem taşımaktadır. Bir yerleşim yerinin nüfus büyüklüğü yalnızca demografik bir gösterge değildir. Bu büyüklük, gündelik karşılaşmaları, kaynaklara erişimi, yerel aktörler arasındaki ilişkileri ve toplumsal görünürlüğü de etkilemektedir (Glick Schiller & Çağlar, 2009).

Winders'ın yeni göç destinasyonlarına ilişkin araştırmaları da bu tartışmayı genişletmektedir. Winders, göçmenlerin yalnızca geleneksel göç merkezlerine yönelmediğini vurgulamaktadır. Göçmenler, daha önce güçlü bir göç geçmişine sahip olmayan yerlere de yerleşmektedir. Göç sürecinin başlangıcı, bu tür yerlerde özellikle açık biçimde gözlemlenebilmektedir. Aynı durum yerel kurumların ve nüfusun ilk tepkileri için de geçerlidir. Göçmenlerin gündelik uyum stratejileri de bu yerlerde incelenebilmektedir. Bu nedenle yeni göç destinasyonları yalnızca yeni varış yerleri değildir. Aynı zamanda göç teorilerinin yeniden sınanabileceği toplumsal mekânlardır. Winders'a göre göçün yeni mekânsal dağılımı, güncel göç süreçlerine ilişkin temel varsayımların yeniden değerlendirilmesini zorunlu kılmaktadır (Winders, 2014).

Bu nedenle küçük yerleşim yerleri söz konusu olduğunda mesele yalnızca göçmen nüfusun sayısı veya yoğunluğu değildir. Küçük yerleşim yerleri, göçün yerel toplumsal ilişkiler içerisinde nasıl anlam kazandığını göstermektedir. Büyük kentlerde anonimlik, mekânsal ayrışma ve kurumsal çeşitlilik göçmenlerin deneyimlerini şekillendirebilmektedir. Buna karşılık küçük yerleşim yerlerinde toplumsal görünürlük ve tekrarlanan temaslar daha güçlü olabilmektedir. Yerel ahlaki beklentiler ve gündelik toplumsal denetim de bu yerlerde daha önemli bir rol oynayabilmektedir. Ancak bu ifade, metropollerle yapılmış ampirik bir karşılaştırma olarak anlaşılmamalıdır. Burada belirleyici olan, toplumsal ilişkilerin küçük yerleşim yerlerinde kendine özgü bir biçimde ortaya çıkmasıdır.

Scholten ve Penninx'in çok düzeyli yönetişime ilişkin tartışmaları da önem taşımaktadır. Bu tartışma, yerel düzeyin yalnızca toplumsal ilişkilerle sınırlı olmadığını göstermektedir. Yerel düzey, politikaların uygulanmasıyla da bağlantılıdır. Yazarlar, göç ve entegrasyon politikalarının artık yalnızca ulusal düzeyde incelenemeyeceğini vurgulamaktadır. Avrupa, ulusal, bölgesel ve yerel düzeyler arasındaki ilişkiler göç politikasının şekillenmesine katkıda bulunmaktadır. Çok düzeyli yönetişim yaklaşımı, yerel aktörlerin her zaman yalnızca merkezî kararları uygulamadığını da göstermektedir. Bununla birlikte, bu yaklaşım küçük yerleşim yerleri bakımından dikkatli biçimde kullanılmalıdır. Yerel düzeyin varlığı her zaman güçlü bir

yerel özerklik anlamına gelmemektedir. Aynı şekilde, bu durum otomatik olarak yenilikçi politikaların ortaya çıkmasına da yol açmamaktadır. Scholten ve Penninx'in merkezîyetçi, yerelci, çok düzeyli ve ayrışmış yönetişim tipolojisi bu noktada yararlı bir çerçeve sunmaktadır. Bu tipoloji aracılığıyla küçük yerleşim yerlerinde merkezî politika ile yerel uygulama arasındaki mesafe incelenebilmektedir (Scholten & Penninx, 2016).

Küçük Yerleşimleri Kuramsal Olarak Önemli Kılan Nedir?

Glorius, Bürer ve Schneider'in çalışması bu noktada özellikle değerlidir. Yazarlar, entegrasyon araştırmalarının çoğu zaman göçmen kadın ve erkeklerin davranışlarına odaklandığını ortaya koymaktadır. Buna karşılık kabul eden toplumun tutum ve davranışları daha seyrek biçimde araştırmanın merkezine alınmaktadır. Bu sorun, kırsal bölgelerde ve küçük yerleşim yerlerinde özellikle belirgin hâle gelmektedir. Bu yerlerde yerel toplumun kabul etme isteği yalnızca kişisel tutumlarla açıklanamaz. Toplumsal ilişkilerin yoğunluğu ve yerel normlar da önem taşımaktadır. Diğer unsurlar arasında sınırlı kamusal alanlar ve ulaşım olanakları yer almaktadır. Dil kursları, okullar, çalışma olanakları ve sivil toplumun kapasitesi de entegrasyonu etkilemektedir. Bu nedenle "yerel kabul kapasitesi" (local receptivity) yaklaşımı perspektifi genişletmektedir. Göç, yalnızca göçmen kadın ve erkeklerin uyum çabaları üzerinden incelenmemelidir. Yerel toplumun pratikleri de dikkate alınmalıdır. Bu pratikler kabul etme, sınırlandırma, gözlemleme ve yorumlama süreçlerini kapsamaktadır.

Bu yaklaşım, küçük yerleşim yerlerinde toplumsal tanışıklığın çift yönlü etkisinin anlaşılmasına yardımcı olmaktadır. Tanışıklık güven oluşturabilmektedir. Göçmen bir aile, öğrenci, işçi veya esnafla düzenli olarak karşılaşılması belirsizliği azaltabilmektedir. Bununla birlikte aynı tanışıklık toplumsal denetimi de güçlendirebilmektedir. Bir kişinin nerede yaşadığı ve kimlerle iletişim kurduğu daha hızlı fark edilebilmektedir. Dil, giyim, meslek ve kamusal alandaki davranışlar da daha kolay gözlemlenmektedir. Bu nedenle küçük yerleşim yerlerinde görünürlük yalnızca orada bulunmaktan ibaret değildir. Göçmen kadın ve erkekler dikkat çekmemeye ve sorun oluşturmamaya da çalışabilmektedir. Belirli sınırları aşmaksızın yerel beklentileri fark etmeleri ve ilişkiler kurmaları gerekmektedir (Glorius et al., 2021).

Woods'un "kırılgan kırsal kozmopolitanizm" kavramı bu noktada önemli bir işaret sunmaktadır. Woods, küçük kentlerde farklı kültürel gruplar arasında gündelik karşılaşmaların ortaya çıkabildiğini göstermektedir. Açıklık ve birlikte yaşama biçimleri de gelişebilmektedir. Bununla birlikte bu kozmopolitanizm ne bütünüyle gerçekleşmiş ne de güvence altına alınmış bir durumdur. Ayrıca geri döndürülemez değildir. Ekonomik koşullara, yerel aktörlere, kamusal alanlara ve kırılgan toplumsal dengelere bağlı olarak şekillenmektedir. Bu yaklaşım, iki tek taraflı bakış açısının önüne geçmektedir. Küçük yerleşim yerleri temelde kapalı ve dışlayıcı mekânlar olarak değerlendirilmemelidir. Bununla birlikte doğal olarak dostane ve kabul edici mekânlar olarak da idealize edilmemelidir (Woods, 2018).

Göçmenler ile yerel nüfus arasındaki ilişkilerin incelenmesi açısından bir başka ayrım da yararlı olmaktadır. Lamont ve Molnár, sembolik sınırlar ile toplumsal sınırlar arasında ayrım yapmaktadır. Sembolik sınırlar, insanların diğer kişileri, davranış biçimlerini, kimlikleri ve mekânları sınıflandırdığı kategorilerdir. Bu kategoriler somut sonuçlar doğurduğunda toplumsal sınırlar ortaya çıkmaktadır. Toplumsal sınırlar, kaynaklara ve toplumsal ağlara erişimi etkileyebilmektedir. Gündelik temas örüntüleri de bu sınırlar tarafından şekillendirilebilmektedir. Küçük yerleşim yerlerinde "bizden biri", "yabancı", "misafir", "uyum sağlamış", "sorun çıkarmayan" veya "fazla görünür" gibi kategoriler bulunmaktadır. Bu kategoriler yalnızca dilsel ifadeler olarak kalmamaktadır. Kimin komşu olarak kabul edileceğini veya kime ev kiralanacağını etkileyebilmektedir. Ayrıca arkadaşlıklar, çalışma olanakları ve bir kişinin kamusal alanda nasıl muamele göreceği üzerinde de etkili olmaktadır (Lamont & Molnár, 2002).

Bu nedenle küçük yerleşim yerleri, toplumsal sınırların incelenmesi açısından elverişli alanlar sunmaktadır. Büyük siyasal söylemlerin gündelik ilişkilere nasıl aktarıldığı bu yerlerde görünür hâle gelebilmektedir. Yerel ahlaki kuralların göçmen kadın ve erkeklere nasıl uygulandığı da gözlemlenebilmektedir. Ayrıca kabulün hangi koşullara bağlı olduğu daha açık biçimde ortaya çıkmaktadır. Bununla birlikte burada da dikkatli olunması gerekmektedir. Küçük yerleşim yerlerindeki her ilişki bir sınır üretmemektedir. Her mesafe dışlanma anlamına gelmemektedir. Mesafe, çatışmalardan kaçınmaya da yardımcı olabilmektedir. Gündelik düzeni koruyabilmekte veya karşılıklı belirsizliği sınırlayabilmektedir.

İnformal Yönetişim, Yerel Kapasite ve Sessiz Uyum

Küçük yerleşim yerlerinde göç, yerel yönetişim pratikleri aracılığıyla da özel bir anlam kazanmaktadır. Burada mesele yalnızca toplumsal karşılaşmalar değildir. Bu tür yerlerde kurumsal kapasiteler çoğu zaman sınırlı olmaktadır. Bu nedenle sağlık, eğitim ve sosyal yardım alanlarındaki sorunlar sıklıkla formel prosedürlerin dışında çözülmektedir. Aynı durum tercüme, barınma ve istihdam için de geçerlidir. Ancak bu durum, küçük yerleşim yerlerinde yönetişimin bulunmadığı anlamına gelmemektedir. Aksine yönetişim, bu yerlerde daha enformel ve daha az görünür olabilmektedir. Ayrıca kişisel ilişkilere daha güçlü biçimde dayanabilmektedir.

Kreichauf'un küçük kentlerde göç yönetişimine ilişkin çalışması bu bağlamda önem taşımaktadır. Bu çalışma, küçük kentlerin göç yönetişiminin pasif mekânları olmadığını göstermektedir. Kreichauf, dağıtım politikalarının ve yerel idari uygulamaların sonuçlarını incelemektedir. Bu politikalar ve uygulamalar, küçük kentleri transit mekânlarına veya bekleme yerlerine dönüştürebilmektedir. Ayrıca bu kentler, göç yönetimine ilişkin görevlerin devredildiği yerel alanlara da dönüşebilmektedir. Bu nedenle küçük yerleşim yerleri yalnızca göçmen kadın ve erkeklerin yerleştiği mekânlar değildir. Aynı zamanda göçün yönetildiği, geciktirildiği, görünür hâle getirildiği veya sınırlandırıldığı yerlerdir (Kreichauf, 2023).

Türkiye açısından Biehl Öztuzcu, Açıkgöz, Eren Benlisoy ve İkizoğlu Erensü bu konuyu doğrudan küçük kentler örneği üzerinden ele almaktadır. 2026 yılında yayımlanan çalışmaları, mültecilerin entegrasyonunun nasıl yönetildiğini incelemektedir. Araştırmacılar bu kapsamda Türkiye'deki küçük kentlere odaklanmaktadır. Bu kentlerde güçlü bir merkezileşme, yerel takdir alanları ve enformel pratikler bir arada bulunmaktadır. Yazarlar, İl Göç İdaresi müdürlükleri, belediyeler ve sivil toplum kuruluşları arasındaki ilişkileri incelemektedir. Mültecilerin yerel düzeyde güvendikleri kişiler de araştırmaya dâhil edilmektedir. Bu ilişkiler merkezî otorite, sınırlı yerel özerklik ve kişisel temaslar tarafından şekillendirilmektedir. Bu nedenle küçük yerleşim yerlerindeki yönetişim, yalnızca yetkilerin formel olarak dağıtılması üzerinden açıklanamaz. Yerel aktörlerin aynı anda birden fazla unsuru dikkate alması gerekmektedir. Bu unsurlar arasında kamusal görünürlüğün beraberinde getirdiği riskler, sınırlı kaynaklar ve olası toplumsal tepkiler yer almaktadır.

Bu noktada "sessiz uyum" kavramı önem kazanmaktadır. Sessiz uyum, göçmen kadın ve erkeklerin veya yerel toplumun üyelerinin entegrasyon sürecine bilinçli ve açık biçimde katılmaları anlamına gelmemektedir. Bu süreç, zorunlu olarak kurumsal prosedürler aracılığıyla da gerçekleşmemektedir. Aksine gündelik yaşam içinde pratik bir düzen ortaya çıkmaktadır. Taraflar, hangi davranışların kabul edildiğini gözlemlemektedir. Belirli olanakları denemektedirler. Bazen geri çekilmektedirler. Diğer durumlarda ise bilinçli olarak mesafeyi korumaktadırlar. Bu düzen açık bir anlaşmaya dayanmamaktadır. Taraflar çoğu zaman bu düzen hakkında birlikte konuşmamaktadır. Bununla birlikte tekrarlanan karşılaşmalar aracılığıyla belirli davranış örüntüleri öğrenilmektedir. Zaman içinde hangi dilin nerede konuşulabileceği anlaşılır hâle gelmektedir. Hassas konuşma konuları da fark edilmektedir. Taraflar, kurumlara hangi taleplerin yöneltilebileceğini öğrenmektedir. Ayrıca hangi sorunların kişisel ilişkiler

aracılığıyla çözülebileceğini ve hangi görünürlük biçimlerinin olumsuz tepkilere yol açabileceğini deneyimlemektedir.
Biehl ve diğerlerinin bulguları bu yaklaşımı desteklemektedir. Bu bulgular, küçük kentlerde görünürlüğün nasıl yönetildiğini göstermektedir. Bazı yerel aktörler, mültecilere yönelik faaliyetlerini bilinçli olarak düşük görünürlükle yürütmektedir. Bu yolla toplumdaki olumsuz tepkilerden kaçınmayı amaçlamaktadırlar. Sivil toplum kuruluşlarının bir denge kurması gerekmektedir. Devlet kurumlarıyla ilişkilerini sürdürebilmek için yeterli düzeyde görünür olmaları gerekmektedir. Aynı zamanda kamuoyunun aşırı dikkatini çekmemeye çalışmaktadırlar. Bu "kontrollü görünürlük", önemli bir özelliğe işaret etmektedir. Sessiz uyum stratejileri geliştirenler yalnızca göçmen kadın ve erkekler değildir. Yerel kurumlar da bu tür stratejilerden yararlanmaktadır (Biehl Öztuzcu et al., 2026).
Bununla birlikte enformel yönetişim idealize edilmemelidir. Enformel ilişkiler, küçük yerleşim yerlerinde bazı sorunların hızlı biçimde çözülmesini sağlayabilmektedir. Kişisel temaslar, tercümanların bulunmasına yardımcı olabilmektedir. Ayrıca gıda dağıtımını veya bir okula kayıt yaptırılmasını kolaylaştırabilmektedir. Bunun yanı sıra kurumlar arasındaki bilgi alışverişini geliştirebilmektedir. Ancak aynı ilişkiler keyfî uygulamaları ve görünmezliği de teşvik edebilmektedir. Ayrıca hesap verebilirliğin zayıflamasına yol açabilmektedir. Bunun sonucunda göçmen kadın ve erkeklerin haklarına ilişkin talepleri zayıflayabilmektedir. Bu nedenle küçük yerleşim yerlerinin teorik değeri, bunların daha iyi veya daha kötü göç mekânları olmasından kaynaklanmamaktadır. Bu yerlerin değeri, göçün pratikte örgütlenmesine ilişkin çoğu zaman görünmez olan biçimleri görünür hâle getirmelerinde yatmaktadır.

Metropol Merkezli Yaklaşımların Eleştirisi: Karşılaştırma Değil, Ölçek Yanlılığı Tartışması

Metropol merkezli yaklaşımların eleştirisi, metropollerin göç araştırmalarındaki önemini reddetmek anlamına gelmemektedir. Sorun, büyük kentlerden hareketle geliştirilen kavramların bütün göç deneyimlerini açıklayabilecek genel çerçeveler gibi kullanılmasıdır. Büyük kentlerde anonimlik, etnik ağların çeşitliliği ve kurumsal yoğunluk göçmen deneyimini şekillendirebilmektedir. Küçük yerleşim yerlerinde ise tanışıklık, tekrarlanan karşılaşmalar, yerel normlar ve sınırlı kurumsal kapasite daha belirgin hâle gelebilmektedir. Bu nedenle tartışma, metropoller ile küçük yerleşim yerleri arasında üstünlük ilişkisi kurmamaktadır. Daha çok, araştırma sorusunun hangi mekânsal ölçeğin hangi ilişki biçimlerini görünür kıldığını dikkate alması gerektiğini savunmaktadır.

Türkiye Bağlamında Küçük Yerleşimleri Çalışmak

Türkiye'de göç araştırmaları büyük ölçüde İstanbul, Ankara ve İzmir gibi büyük kentlere odaklanmaktadır. Gaziantep, Kilis ve Hatay gibi sınır kentleri de sıklıkla araştırmaların merkezinde yer almaktadır. Bir diğer odak noktası, Suriyeli nüfusun yoğun olduğu bölgelerdir. Bu yönelimin anlaşılabilir nedenleri bulunmaktadır. Bununla birlikte Türkiye'nin göç deneyimi büyük kentler ve sınır bölgeleriyle sınırlı değildir. Küçük ve orta büyüklükteki kentlerde de farklı gruplar arasında gündelik ilişkiler ortaya çıkmaktadır. Bu gruplar arasında göçmen kadın ve erkekler, uluslararası öğrenciler ve sığınmacılar yer almaktadır. Geçici koruma altındaki kişiler ile düzensiz göçmenler de bu kentlerde yaşamaktadır. Bu kişiler gündelik yaşamda yerel toplumun üyeleriyle temas kurmaktadır. Söz konusu ilişkiler ulusal politika belgelerinde çoğu zaman yeterince görünür kılınmamaktadır. Büyük kentlere odaklanan araştırmalarda da bu ilişkilere sınırlı ölçüde yer verilmektedir.
Biehl Öztuzcu ve diğerlerinin çalışması önemli bir özelliği ortaya koymaktadır. Küçük kentlerde entegrasyon yönetişimi, güçlü bir merkezileşme ve sınırlı yerel kapasiteler koşullarında ortaya çıkmaktadır. Yerel aktörler bu süreçte bütünüyle pasif değildir. Bununla birlikte hareket alanları büyük ölçüde sınırlandırılmaktadır. Merkezi otorite, siyasal

hassasiyetler ve kamuoyu tepkileri bu süreçte önemli bir rol oynamaktadır. Sınırlı kaynaklar da yerel aktörlerin hareket olanaklarını etkilemektedir. Bu bulgular, Türkiye'de küçük yerleşim yerlerinin neden incelenmesi gerektiğini göstermektedir. Merkezi politika ile gündelik uygulama arasındaki mesafe bu yerlerde görünür hâle gelmektedir. Formel görevler ile fiilî sorumluluklar arasında da gerilimler ortaya çıkabilmektedir. Ayrıca kurumsal görünürlük ile toplumsal tepkiler arasında hassas bir denge bulunduğu görülmektedir.

Küçük yerleşim yerleri, Türkiye bağlamında entegrasyon kavramına ilişkin tartışmalar açısından da önem taşımaktadır. Resmî belgelerde entegrasyon karşılıklı ve gönüllü bir süreç olarak tanımlanmaktadır. Ancak pratikte entegrasyon çoğu zaman göçmen kadın ve erkeklerden beklenen davranışlara indirgenmektedir. Küçük yerleşim yerlerinde entegrasyon daha sessiz ve ilişkisel bir biçimde gerçekleşebilmektedir. Bu yerlerde göçmenlerin hangi davranışlar aracılığıyla "sorun çıkarmayan" kişiler olarak değerlendirildiği daha açık biçimde gözlemlenebilmektedir. Yerel toplumun hangi sınırları koruduğu da görünür hâle gelmektedir. Kurumların pratikleri de bu kapsamda incelenebilmektedir. Bu inceleme, kurumların hangi talepleri görünür kılmak istemediği sorusunu da kapsamaktadır. Ayrıca hangi sorunların kişisel ilişkiler aracılığıyla çözüldüğü gözlemlenebilmektedir.

Bu tartışma, küçük yerleşim yerlerinin Türkiye'deki göç araştırmalarının merkezine yerleştirilmesi gerektiği yönünde bir çağrı olarak anlaşılmamalıdır. Aksine küçük yerleşim yerleri, mevcut literatüre şimdiye kadar ihmal edilmiş bir mekânsal düzey ve özgün bir toplumsal ilişki biçimi eklemektedir. Bu yerler, göçün yalnızca krizler, nüfus yoğunluğu, kurumsal kapasiteler veya sınır güvenliği üzerinden incelenmemesi gerektiğini göstermektedir. Gündelik düzen, toplumsal tanışıklık ve mesafe de önem taşımaktadır. Aynı durum sessizlik ve enformel çözüm yolları için de geçerlidir.

Sonuç

Bu çalışmada küçük yerleşim yerlerinin göç araştırmaları açısından teorik değeri tartışılmıştır. Çalışmanın temel tezi, küçük yerleşim yerlerinin daha az önemli veya sınırlı alanlar olarak değerlendirilmemesi gerektiğidir. Bu durum, özellikle metropollerle karşılaştırma açısından önem taşımaktadır. Küçük yerleşim yerleri araştırma açısından önemli olanaklar sunmaktadır. Göçün farklı bir mekânsal düzeyde nasıl deneyimlendiğini göstermektedir. Ayrıca toplumsal ilişkilerin nasıl ortaya çıktığını ve yerel sınırların gündelik yaşamda nasıl oluşturulduğunu görünür kılmaktadır.

Göçmen kadın ve erkekler ile yerel toplumun üyeleri, küçük yerleşim yerlerinde çoğu zaman aynı gündelik mekânları paylaşmaktadır. Bu mekânsal yakınlık otomatik olarak toplumsal yakınlığa yol açmamaktadır. Tekrarlanan karşılaşmalar bazı durumlarda güveni güçlendirebilmektedir. Bununla birlikte daha istikrarlı bir toplumsal mesafenin oluşmasına da yol açabilmektedir. Yerel toplum tarafından kabul edilme çoğu zaman koşullu ve kırılgan niteliktedir. Bu kabulün sınırları farklı beklentiler tarafından belirlenebilmektedir. Bu beklentiler arasında düşük görünürlük, yerel normlara uyma ve ekonomik katkı yer almaktadır. Çekingenlik ve sorun çıkarmayan davranış olarak değerlendirilen tutumlar da önem taşıyabilmektedir. Bu nedenle küçük yerleşim yerleri, entegrasyonun yalnızca kurumsal göstergeler aracılığıyla açıklanamayacağını hatırlatmaktadır. Gündelik ilişkilerin de dikkate alınması gerekmektedir.

Çalışma, küçük yerleşim yerlerinin üç temel teorik katkısını ortaya koymaktadır. Birincisi, yerellik ve mekânsal ölçek meselelerini göç araştırmalarına daha güçlü biçimde dâhil etmektedir. İkincisi, gündelik karşılaşmaları, toplumsal mesafeyi, toplumsal tanışıklığı ve aidiyet sınırlarını daha görünür hâle getirmektedir. Üçüncüsü, sınırlı kurumsal kapasitelerin ve enformel yönetişimin incelenmesine olanak sağlamaktadır. Kişisel ilişkiler ve kontrollü görünürlük de bu yerlerde daha ayrıntılı biçimde analiz edilebilmektedir. Bu katkılar, büyük kent araştırmalarından geliştirilen kavramların bütünüyle terk edilmesini gerektirmemektedir.

Bununla birlikte söz konusu kavramların farklı mekânsal ölçekler daha fazla dikkate alınarak yeniden düşünülmesi gerekmektedir.

Bu çalışmanın bir sınırlılığı, kendi ampirik saha araştırmasına dayanmamasıdır. Tartışma, seçilmiş literatür temelinde yürütülen kavramsal bir değerlendirme niteliğindedir. Bu nedenle nitel saha araştırmalarına ihtiyaç duyulmaktadır. Bu araştırmalar, göçmen kadın ve erkekler ile yerel nüfus arasındaki ilişkilerin küçük yerleşim yerlerinde nasıl ortaya çıktığını daha ayrıntılı biçimde gösterebilmektedir. Özellikle her iki grupla da görüşmeler gerçekleştirilmesi önem taşımaktadır. Belediyelerin, yerel makamların ve sivil toplum kuruluşlarının temsilcileri de araştırmaya dâhil edilmelidir. Arabuluculuk rolleri üstlenen kişilerle yapılacak görüşmeler de yararlı olacaktır. Bu yolla çalışmanın kavramsal varsayımları daha somut biçimde sınanabilecektir.

Küçük yerleşim yerleri, göç araştırmalarında ikincil veya yalnızca tamamlayıcı araştırma alanları olarak değerlendirilmemelidir. Bunlar teorik açıdan önemli mekânlardır. Göçün sessiz, ilişkisel ve gündelik boyutlarını görünür hâle getirmektedirler. Bu yaklaşım, göç araştırmalarının perspektifini genişletmektedir. Göç yalnızca nüfus yoğunluğu, kurumsal çeşitlilik ve büyük kent yapılarına dayanarak incelenmemelidir. Yerel karşılaşmalar, toplumsal sınır çizme pratikleri ve görünürlüğü düşük entegrasyon biçimleri de önem taşımaktadır.

Kaynakça

Biehl, K. S., Açıkgöz, M., Eren Benlisoy, Z. C., & İkizoğlu Erensü, A. (2026). Integration and multi-level governance in Turkey's small towns: An actor centred analysis. *International Migration, 64*, e70163. https://doi.org/10.1111/imig.70163

Glick Schiller, N., & Çağlar, A. (2009). Towards a comparative theory of locality in migration studies: Migrant incorporation and city scale. *Journal of Ethnic and Migration Studies, 35*(2), 177–202. https://doi.org/10.1080/13691830802586179

Glorius, B., Bürer, M., & Schneider, H. (2021). Integration of refugees in rural areas and the role of the receiving society: Conceptual review and analytical framework. *Erdkunde, 75*(1), 51–60. https://doi.org/10.3112/erdkunde.2021.01.04

Kreichauf, R. (2023). Governing migration through small towns: Dispersal and the production of spaces of transit. *International Journal of Urban and Regional Research, 47*(3), 349–367. https://doi.org/10.1111/1468-2427.13171

Lamont, M., & Molnár, V. (2002). The study of boundaries in the social sciences. *Annual Review of Sociology, 28*, 167–195. https://doi.org/10.1146/annurev.soc.28.110601.141107

Scholten, P., & Penninx, R. (2016). The multilevel governance of migration and integration. In B. Garcés-Mascareñas & R. Penninx (Eds.), *Integration processes and policies in Europe: Contexts, levels and actors* (pp. 91–108). Springer. https://doi.org/10.1007/978-3-319-21674-4_6

Winders, J. (2014). New immigrant destinations in global context. *International Migration Review, 48*(1_suppl), S149–S179. https://doi.org/10.1111/imre.12140

Woods, M. (2018). Precarious rural cosmopolitanism: Negotiating globalization, migration and diversity in Irish small towns. *Journal of Rural Studies, 64*, 164–176. https://doi.org/10.1016/j.jrurstud.2018.03.014

MÜHARİBƏ DÖVRÜ UŞAQLARININ MƏNƏVİ DÜNYASI VƏ İNSANLIQ DƏRSLƏRİ
MAQSUD İBRAHİMBƏYOVUN "PÜSTƏ AĞACI" ƏSƏRİNDƏ SİMVOLİZM VƏ SOSİAL REALLIQ

GÜNEL HÜMMƏTOVA (Pəniyeva) Saleh
Filologiya üzrə fəlsəfə doktoru (Ph.D.), Baş müəllim
Azərbaycan Dövlət Pedaqoji Universiteti, Ədəbiyyat ın tədrisi texnologiyası kafedrası
ORCID ID: 0000-0002-8916-3484

Xülasə

Məqalədə görkəmli yazıçı Maqsud İbrahimbəyovun "Püstə ağacı" hekayəsi metodik və fəlsəfi müstəvidə təhlil olunur. Əsərdə müharibə dövrü uşaqlarının psixologiyası, onların qeyri-şüuri olaraq reallıqdan qaçış forması olan "dava-dava" oyunu və mətndəki simvolik elementlər araşdırılır. Tədqiqat zamanı professor Soltan Hüseynoğlunun ədəbiyyat dərslərində interpretasiya və fəal oxu metodologiyasına istinad edilmiş, əsərdən gətirilən sitatlarla uşaqların mənəvi inkişaf dinamikası elmi şəkildə əsaslandırılmışdır.

Açar sözlər: *Maqsud İbrahimbəyov, "Püstə ağacı", Soltan Hüseynoğlu, interpretasiya, müharibə psixologiyası, bədii simvol, uşaq mütaliəsi.*

ABSTRACT

This article provides a methodological and philosophical analysis of the short story "The Pistachio Tree" ("Püstə ağacı") by the prominent writer Maqsud Ibrahimbeyov. The study explores the psychology of wartime children, their "war-play" ("dava-dava") game acting as an unconscious escape from reality, and the symbolic elements within the text. Drawing upon Professor Soltan Huseynoglu's methodology on interpretation and active reading in literature classes, the research utilizes direct quotes from the story to scientifically substantiate the dynamics of children's moral and spiritual development.

Keywords: *Maqsud Ibrahimbeyov, "The Pistachio Tree", Soltan Huseynoglu, interpretation, wartime psychology, literary symbol, children's reading.*

Ədəbiyyat tədrisinin qarşısında duran fundamental vəzifələrdən biri şagirdlərin oxuduqları bədii mətni sadəcə süjet səviyyəsində deyil, mətnaltı məna qatları və müəllif ideyası müstəvisində qavramalarını təmin etməkdir. Soltan Hüseynoğlu "Ədəbiyyat dərslərində interpretasiya" adlı tədqiqatında göstərir ki, *"bədii mətnin fəal yozumu şagirdin idrakı aktivliyini artırır, onu yazıçı ilə birbaşa dialoqa girməyə sövq edir"*. Bu baxımdan Maqsud İbrahimbəyovun "Püstə ağacı" hekayəsi həm psixoloji dərinliyi, həm də humanizm ideyaları ilə şagirdlərin estetik və mənəvi təfəkkürünün inkişafı üçün əvəzsiz mənbədir. Hekayə arxa cəbhədə – Bakıda yaşayan uşaqların müharibəni fərqli, bir qədər də romantik qavrayışı ilə başlayır. Bakıya bomba düşməməsi uşaqlar tərəfindən bir təəssüf hissi ilə qarşılanır: *"Müharibə vaxtı Bakıya bomba düşmədi. Heç bircə dənəsi də düşmədi. Deyəsən, bizdən – məhəllə uşaqlarından başqa, heç kəs buna təəssüflənmədi."* Soltan Hüseynoğlu qeyd edir ki, uşaq təxəyyülü hələ müharibənin dəhşətlərini tam dərk etmədiyi üçün onu bir macəra, "dava-dava" oyunu kimi qəbul edir. Uşaqlar Salyan kazarmalarının yaxınlığındakı zibillikdən tapdıqları köhnə silah hissələri ilə oynayır, zirzəmidə konserv qutularına yığdıqları güllələri ocağın üstünə qoyub partladaraq həyəcan yaşamaq istəyirlər. Onlar qonşuları qorxuya salsalar da, bunu öz

daxili dünyalarında müharibəyə bir "töhfə", oyunun davamı sayırlar: *"Biz də, əlbəttə, özümüzü elə göstərirdik ki, guya, heç nədən xəbərimiz yoxdur, amma, əlbəttə, dava-dava oynamaqda davam edirdik."*Yazıçı bu detal vasitəsilə göstərir ki, uşaqlar əslində müharibənin "dəhşətli" tərəfini yaşlıların söhbətindən eşitsələr də, hələ ki mənəvi olaraq ondan uzaqdırlar: *"Biz onun haqqında yaşlılardan eşidirdik, amma onu dəhşətli saymırdıq."*Hekayədəki Püstə ağacı sadəcə bir bitki deyil, həyətin bütövlüyünün, birliyinin və sevincinin rəmzidir. Payızda püstə yetişəndə uşaqların onu bərabər bölüb qonşulara paylaması həyətdəki humanizm mühitini göstərir:*"Püstəni əvvəlcə qərargaha toplayır, sonra hər mənzilə bir böyük kağız torbada paylayırdıq. Hamı da bizə «sağ ol» deyirdi."*Lakin həyətetə köçən yeni qonşu – gecə-gündüz paltar yuyan, uşaqları ilə tənha yaşayan qadın bu harmoniyanı pozur. O, paltar yuduğu kələvili (sabunsuz, kimyəvi qatqılı) suyu püstə ağacının dibinə tökür. Müəllif bu hərəkətlə ağacın tədricən məhv olmasına zəmin hazırlayır. Soltan Hüseynoğlunun metodik yanaşmasına görə, dərslərdə qadının bu davranışının motivini şagirdlərlə birlikdə interpretasiya etmək vacibdir. Qadın niyə belə sərtdir? Ananın dili ilə verilən bu mülahizə məsələyə işıq salır: *"Anam deyirdi ki, çətin həyat insanların xasiyyətini korlayır..."*Uşaqlar öz ordularını yaradır, hərəsinə rütbələr verirlər ("polkovnik", "general", "marşal"). Lakin bu "böyük sərkərdələr" ağacın quruması qarşısında aciz qalırlar. Püstə ağacı bir gecədə yarpaqlarını tökür:*"Ağac çılpaq idi, bir gecədə bütün yarpaqlarını tökmüşdü, lüt budaqda püstə salxımları görünürdü."*Aqronomun gəlib *"Bu ağacdan yalnız odun olar"* deməsi uşaqların mənəvi dünyasında ilk ciddi sarsıntını yaradır. Onlar günahkar kimi gördükləri qadının qapısına qəzəblə gedirlər.Uşaqlar qadının otağına daxil olduqda onun ağladığını görürlər. Bu epizod əsərin ən güclü psixoloji kulminasiya nöqtəsidir. Müharibənin amansızlığı ilk dəfə uşaqların gözü qarşısında vizuallaşır: *"O, çarpayıda oturub ağlayırdı. Mən heç vaxt bir adamın belə ağladığını görməmişdim. O bizə fikir vermədən paltar yumaqdan şişmiş, göy damarları çıxmış əlləri ilə üzünü tutub ağlamaqda davam edirdi."* Bu mənzərə uşaqların daxilindəki bütün "əsgər", "general" iddialarını darmadağın edir. Onlar anlayırlar ki, dünyanın ən böyük ordusu və ən gözəl oyunları belə bir ananın göz yaşının qarşısında əhəmiyyətsizdir: *"Biz – polkovniklər, generallar, marşallar onun qarşısında durub susurduq. Hamımızı qəhər boğurdu. Biz ona demək istəyirdik ki, dünyanın ən gözəl ağacı onun göz yaşlarına dəyməz..."* Soltan Hüseynoğlu qeyd edir ki, *"bu məqam şagirdlərdə empatiya hissinin pik həddə çatdığı andır"*. Uşaqlar dərhal mənəvi reaksiya verirlər. Onlar rütbə almaq üçün yığdıqları pulları götürüb Kubinkaya gedir və yeni bir ağac tingi alırlar. Hər nə qədər satıcı onları aldadıb püstə əvəzinə başqa bir ting satsa da, bu hərəkətin özü uşaqların mənəvi təmizliyinin, humanistliyinin göstəricisidir. Qadının ərinin cəbhədə həlak olması xəbərinin (qara kağızın) gəlməsi isə uşaqların xəyali müharibəsinə birdəfəlik son qoyur:*"Həmin gün bizim həyətdə dava-dava oyununa son qoyuldu..."*Maqsud İbrahimbəyovun "Püstə ağacı" hekayəsi yalnız müharibə dövrünün sosial mənzərəsini əks etdirmir, həm də müasir ədəbiyyat dərslərində şagirdlərin emosional intellektinin (EQ) inkişafı üçün zəngin metodik material rolunu oynayır. Professor Soltan Hüseynoğlunun interaktiv təlim konsepsiyasında qeyd olunduğu kimi, *"bədii qəhrəmanın hisslərinə şərik olmaq şagirdin daxili aləmini zənginləşdirir və onda humanizm prinsiplərini formalaşdırır"*. Bu baxımdan, əsərin tədrisi zamanı şagirdlərin idraki fəallığını artırmaq üçün xüsusi interpretasiya suallarının və problem situasiyalarının qurulması məqsədəuyğundur. Mətndə uşaqların yeni qonşunun otağında qarşılaşdıqları mənzərə mənəvi katarsis (təmizlənmə) anıdır. Bu epizodun təhlili zamanı sinifdə aşağıdakı metodik fəaliyyət növlərindən istifadə edilə bilər:Şagirdlərə *"Siz o otağa daxil olan*

'generalların' yerində olsaydınız, qadının göz yaşlarını gördükdə nə hiss edərdiniz?" sualı ilə müraciət olunur. Bu sual şagirdi mətndəki uşaqların keçirdiyi qəhər və xəcalət hissini dərindən anlamağa yönəldir.Qadının püstə ağacının dibinə kələvili su tökməsi ilk baxışda ətraf mühitə qarşı laqeydlik kimi görünsə də, mətnaltı qatda onun ağır ailə dramı, cəbhədən gözlədiyi "qara kağız"ın gərginliyi dayanır. Soltan müəllimin metodik yanaşmasına istinadən, şagirdlərə hadisələrə birtərəfli yox, fərqli fərdlərin prizmasından baxmaq bacarığı (multiperspektivlik) aşılanır. Uşaqların rütbə pullarını birləşdirib Kubinkadan yeni ağac tingi almaları, onların sadəcə təəssüf hissi keçirmədiyini, həm də səhvlərini düzəltmək üçün real addım atdıqlarını (aktiv humanizm) sübut edir. Bu məqamda şagirdlərlə *"Ağacın püstə çıxmaması uşaqların əməyinin dəyərini azaldırmı?"* mövzusunda kiçik debat təşkil edilə bilər.Beləliklə, hekayənin metodik mənimsənilməsi prosesi göstərir ki, Maqsud İbrahimbəyovun nəsri gənc nəslə subyektiv mülahizələrdən uzaqlaşıb obyektiv həqiqətləri görmək və başqalarının ağrısına şərik olmaq kimi ali insani keyfiyyətləri təlqin edir.

NƏTİCƏ

Maqsud İbrahimbəyovun "Püstə ağacı" hekayəsi müharibənin gətirdiyi faciələrin arxa cəbhədəki uşaqların psixologiyasında yaratdığı böyük dönüşü əks etdirir. Soltan Hüseynoğlunun elmi-metodik konsepsiyalarında vurğulandığı kimi, bu əsərin təhlili şagirdlərə təkcə ədəbi qəhrəmanları deyil, real həyatdakı insan talelərini və mənəvi borc hissini dərindən anlamağa imkan yaradır. Sonda həmin naməlum ağacın böyüyüb hamıya kölgə salması və qonşuların *"Onun püstə olmamasına təəssüflənməməsi"* insan mənəviyyatının, sevgi və həmrəyliyinin hər cür maddi və bədii gözləntidən daha uca olduğunu sübut edir.

ƏDƏBİYYAT SİYAHISI

1. Hüseynoğlu S. (2023). Ədəbiyyat dərslərində interpretasiya. *Azərbaycan məktəbi*. № 3 (704), səh. 9-24.
2. İbrahimbəyov M. *Püstə ağacı* (hekayə - mətn əsasında).
3. Hüseynoğlu S. (2009). *Ədəbiyyatın tədrisi metodikası*. Bakı: ADPU nəşriyyatı.

KÜÇÜK KENTLERDE GÖÇÜN GÖRÜNMEYEN YÖNETİŞİMİ: YEREL KURUMLAR, GÜNDELİK PRATİKLER VE KURUMSAL SINIRLAR

THE INVISIBLE GOVERNANCE OF MIGRATION IN SMALL TOWNS: LOCAL INSTITUTIONS, EVERYDAY PRACTICES, AND INSTITUTIONAL LIMITS

Prof. Dr. Metin AKSOY
Gümüşhane University, Department of International Relations
ORCID:0000-0002-3118-6739,

Muhlise CANLI
Lect., Gümüşhane University, Vocational School of Social Sciences, Department of Social Work and Counseling
ORCID:0000-0001-6051-4677

Özet

Göç yönetişimi denildiğinde genellikle ulusal politikalar, sınır kentleri, büyükşehir belediyeleri ve uluslararası kuruluşlar akla gelmektedir. Bunun sonucunda küçük kentlerde göçle ilgili işlerin gerçekte nasıl yürütüldüğü çoğu zaman araştırmanın dışında kalır. Oysa merkezî düzeyde alınan bir karar, küçük bir kente ulaştığında her zaman mevzuatta öngörüldüğü biçimde uygulanmaz. Kurumun personel sayısı, çalışanların önceki deneyimleri, kentte kimlerin birbirini tanıdığı ve başvurulabilecek başka bir kurumun bulunup bulunmadığı uygulamayı etkileyebilir. Göçmenlerin sağlık, eğitim veya sosyal yardım hizmetlerine erişirken izledikleri yollar da bu yerel koşullar içinde şekillenmektedir.

Bu bildiride "görünmeyen yönetişim" kavramıyla tam olarak bu alana dikkat çekilmektedir. Söz konusu alanın sınırları önceden çizilmiş değildir. Bir göçmenin yaşadığı sorun kimi zaman belediyedeki bir çalışanın girişimiyle çözülmektedir. Muhtarın kentte tanıdığı kişiler veya yerel bir derneğin sağladığı destek de aynı süreçte etkili olabilmektedir. Aynı ilişkiler erişimi zorlaştırabilir veya göçmenin kurumlardan ne ölçüde yararlanabileceğini sınırlandırabilir. Bu nedenle küçük kentlerde yönetişim, resmî kurumların faaliyetlerinden ibaret görülmemelidir. Kurumların yapamadıkları, çalışanların kendi imkânlarıyla yaptıkları ve gündelik ilişkiler yoluyla çözülen meseleler de bu sürece dâhildir.

Küçük kentlerde göçmenler daha kolay fark edilmekte ve aynı kişilerle tekrar tekrar karşılaşabilmektedir. Tanınmak bazı göçmenlerin hizmetlere ulaşmasını kolaylaştırmaktadır. Fakat küçük bir kentte sürekli görünür olmak, kişinin davranışlarının daha yakından izlenmesi anlamına da gelebilir. Göçmenin yerel toplum tarafından kabul edilmesi, kurumların sunduğu hizmetlerden bütünüyle ayrı değildir. Ancak ikisi her zaman aynı yönde ilerlemez. Bildiri, küçük kentleri merkezî politikaların değişmeden uygulandığı pasif alanlar olarak değil, bu politikaların yerel imkânlar ve gündelik ilişkiler içinde yeniden biçimlendiği yerler olarak ele almaktadır. Böyle bir bakış, resmî politika ile gündelik uygulama arasında kalan ve çoğu zaman kayıtlara yansımayan yönetişim alanını daha yakından incelemeye imkân verebilir.

Anahtar Kelimeler: Göç yönetişimi, küçük kentler, yerel kurumlar, görünmeyen yönetişim, toplumsal uyum

Abstract

When migration governance is discussed, attention usually turns to national policies, border cities, metropolitan municipalities, and international organisations. As a result, the way migration-related matters are actually handled in small towns often remains outside the scope of research. Yet a decision made at the central level is not always implemented in a small town exactly as legislation prescribes. Implementation may be affected by the number of staff within an institution, the previous experience of its employees, personal connections within the town,

and the availability of another institution to which an application can be made. The ways migrants access healthcare, education, and social assistance are also shaped by these local conditions.
This paper uses the concept of "invisible governance" to draw attention to this particular area. Its boundaries are not defined in advance. A problem experienced by a migrant may sometimes be resolved through the initiative of a municipal employee. The people known by the neighbourhood head, or muhtar, and the support provided by a local association may also influence the process. The same relationships may make access more difficult or limit the extent to which migrants can benefit from institutions. Governance in small towns should therefore not be understood as consisting solely of the activities of formal institutions. What institutions are unable to do, what employees accomplish using the resources available to them, and the issues resolved through everyday relationships are also part of this process.
Migrants in small towns are more easily noticed and may repeatedly encounter the same people. Being known can make it easier for some migrants to access services. However, remaining constantly visible in a small town may also mean that their behaviour is observed more closely. Migrants' acceptance by the local community is not entirely separate from the services provided by institutions, although the two do not always move in the same direction. This paper approaches small towns not as passive settings in which central policies are implemented without change, but as places where these policies are reshaped through local resources and everyday relationships. Such a perspective may allow closer examination of the field of governance that lies between formal policy and everyday practice and is often absent from official records.
Keywords: Migration governance, small towns, local institutions, invisible governance, social cohesion.

Giriş

Göç yönetişimine ilişkin çalışmalar uzun süre ulusal siyasal çerçevelere odaklanmıştır. Sınır yönetimi, büyük kentlerin hizmet sunma kapasitesi ve uluslararası kuruluşların rolü de araştırmaların merkezinde yer almıştır. Bu yönelimin anlaşılabilir nedenleri bulunmaktadır. Göç hukukuna ilişkin temel kararlar çoğunlukla merkezî devlet düzeyinde alınmaktadır. Hukuki statü grupları ulusal yasalar aracılığıyla belirlenmektedir. Ayrıca birçok göçmen büyük kentlerde yaşamaktadır. Bununla birlikte göçün toplumsal sonuçları yalnızca bu düzeylerde ortaya çıkmamaktadır. Bir okula kayıt yaptırma, sağlık hizmetlerine erişme, sosyal yardım başvurusunda bulunma, mahallede görünür olma ve yerel toplumla temas kurma gibi süreçler daha küçük mekânsal bağlamlarda gelişmektedir. Bu nedenle yerel düzey, siyasal uygulamaların yalnızca teknik olarak gerçekleştirildiği bir alan değildir. Yerel düzey, siyasal tedbirlerin anlamının değişebildiği ilişkisel bir mekândır.

Yerel düzeye yönelik bu ilgi, daha geniş bir gelişmenin parçasıdır. Göç ve entegrasyon araştırmalarında bu gelişme local turn olarak adlandırılmaktadır. Zapata-Barrero, Caponio ve Scholten (2017), göçmenlerin entegrasyonunun giderek yerel aktörler tarafından şekillendirildiğini ortaya koymaktadır. Bu aktörler arasında belediyeler ve sivil toplum kuruluşları yer almaktadır. Farklı siyasal düzeyler arasındaki ilişkiler de önemli bir rol oynamaktadır. Caponio ve Borkert'in (2010) yerel göç politikası üzerine çalışması da benzer bir sonuca ulaşmaktadır. Entegrasyon yalnızca ulusal düzeyde belirlenen genel bir hedef değildir. Entegrasyon, belediyeler, yerel ağlar ve kamu kurumları aracılığıyla gündelik yaşama ulaşmaktadır. Yerel entegrasyona ilişkin OECD raporu da bu noktayı vurgulamaktadır. Entegrasyon, insanların yaşadığı, çalıştığı, eğitim gördüğü ve gündelik yaşamda karşılaştığı yerlerde gerçekleşmektedir (OECD, 2018). Bu yaklaşım küçük kentleri doğrudan merkeze almamaktadır. Bununla birlikte göç yönetişiminin yalnızca ulusal kararlar aracılığıyla açıklanamayacağını göstermektedir.

Bu çalışmanın temel sorusu şu şekilde formüle edilmektedir: Göç yönetişimi, resmî politikaların ötesinde küçük kentlerde hangi gündelik, kurumsal ve toplumsal mekanizmalar aracılığıyla işlemektedir? Bu soru önemli bir varsayımı gerekli kılmaktadır. Küçük kentler, büyük kentlerin küçültülmüş kopyaları değildir. Göçmen kadın ve erkekler bu kentlerde daha kolay fark edilebilmektedir. Çoğu zaman aynı kişilerle karşılaşmaktadırlar. Yerel kurumlarla ilişkileri de sıklıkla kişisel temaslara dayanmaktadır. Bu durum hizmetlere erişimi kolaylaştırabilmektedir. Bununla birlikte toplumsal denetimi de güçlendirebilmektedir. Ayrıca görünürlük bir risk hâline gelebilmektedir. Bu nedenle göçmenlerin kurumlarla ilişkileri de sınırlandırılabilmektedir.

Bu çalışma, söz konusu ara alanı incelemek amacıyla "görünmeyen yönetişim" kavramını kullanmaktadır. Bu kavram, tamamen devletin veya yerel kurumların dışında kalan enformel bir düzeni ifade etmemektedir. Aksine görünmeyen yönetişim, formel politika ile gündelik pratik arasında ortaya çıkmaktadır. Çoğu zaman belgelenmemektedir. Bununla birlikte hizmetlere erişimi etkilemektedir. Yerel toplumla ilişkileri ve kurumlarla yaşanan deneyimleri de şekillendirmektedir. Bir belediye çalışanının girişimi bu yönetişimin bir parçası olabilmektedir. Bir yerel muhtar aracılığıyla gerçekleştirilen aracılık da aynı kapsamda değerlendirilebilir. Yerel bir dernek tarafından sağlanan tercüme desteği ve bir okul yönetiminin geliştirdiği pratik çözümler de diğer örnekleri oluşturmaktadır. Kişisel tanışıklıklar da sosyal yardım başvurusunda rol oynayabilmektedir. Bununla birlikte aynı ilişkiler dışlayıcı, seçici veya sınırlandırıcı biçimlerde de işleyebilmektedir.

Yerel Göç Yönetişimi ve Çok Düzeyli Çerçeve

Göç yönetişimi kavramı önemli bir nedenle yararlıdır. Göçe ilişkin kararlar yalnızca devlet tarafından ve tek yönlü biçimde alınmamaktadır. Bu kararlar farklı aktörler, siyasal düzeyler ve kurumlar arasında dağılmaktadır. Çok düzeyli yönetişim yaklaşımı, bu dağılımı dikey ve yatay ilişkiler üzerinden incelemektedir. Dikey boyut, ulusal, bölgesel ve yerel düzeyler arasındaki ilişkilere odaklanmaktadır. Yatay boyut ise yerel düzey içindeki ilişkilere yönelmektedir. Bu ilişkiler arasında belediyeler, kamu kurumları ve sivil toplum kuruluşları arasındaki ilişkiler yer almaktadır. Gönüllüler, yerel kanaat önderleri ve göçmen toplulukları da bu süreçte rol oynamaktadır (Scholten & Penninx, 2016).

Bununla birlikte çok düzeyli yönetişim kavramı küçük kentlere doğrudan aktarılamaz. Büyük kentler çoğu zaman uzmanlaşmış belediye birimlerine sahiptir. Bu kentler uluslararası ağlara erişebilmekte ve göçmenlere yönelik kurumsallaşmış programlar sunabilmektedir. Ayrıca genellikle daha geniş sivil toplum yapılarına sahip olmaktadırlar. Küçük kentlerde ise bu tür yapılar çoğunlukla sınırlıdır. Bu nedenle çok düzeyli yönetişim, küçük kentlerde her zaman düzenli bir koordinasyon anlamına gelmemektedir. Kurumsal kapasiteler ve görevlerin açık biçimde dağıtılması da kendiliğinden mevcut değildir. Biehl, Açıkgöz, Eren Benlisoy ve İkizoğlu Erensü'nün (2026) Türkiye'deki küçük kentler üzerine gerçekleştirdiği çalışma bu açıdan özellikle önem taşımaktadır. Yazarlar, mültecilerin entegrasyonunu incelemektedir. Bu sürecin güçlü bir merkezîleşme ve sınırlı yerel özerklik tarafından şekillendirildiğini ortaya koymaktadırlar. Enformel koordinasyon, seçici görünürlük ve kişisel ilişkiler de bu süreçte önemli bir rol oynamaktadır. Bu bulgu, küçük kentlerde yönetişimin yalnızca formel kurumsal yapılar üzerinden anlaşılmasının mümkün olmadığını göstermektedir.

Local turn yaklaşımına dayanan araştırmalar, belediyeleri ve kentleri göç yönetişiminin aktif aktörleri olarak ele almaktadır. Bununla birlikte bu literatürün önemli bir bölümü büyük kent deneyimleri temelinde oluşturulmuştur. Araştırmaların merkezinde çoğunlukla yenilikçi yerel politikalar ve belediye ağları yer almaktadır. Çok sayıda aktörü bir araya getiren platformlar ile göçmenlere yönelik uzmanlaşmış hizmetler de incelenmektedir. Buna karşılık küçük kentlerde farklı bir soru ön plana çıkmaktadır. Burada belirleyici olan, sınırlı kaynaklarla gündelik sorunların üstesinden gelebilme kapasitesidir.

Jonitz, Schiller ve Scholten'in (2026) Hollanda'daki küçük belediyeler üzerine gerçekleştirdiği çalışma bu yaklaşımı desteklemektedir. Çalışma, göç yönetişiminin yalnızca aktörler arasındaki iş birliği tarafından şekillendirilmediğini göstermektedir. Çatışmalar ve güç farklılıkları da önem taşımaktadır. Diğer unsurlar arasında yerel siyasal tutumlar, belediye yönetiminin liderliği ve nüfusun tutumu yer almaktadır. Bu nedenle küçük kentler, iş birliğinin kendiliğinden daha kolay gerçekleştiği homojen mekânlar değildir. Sınırlı sayıdaki aktör ve yakın ilişkiler iş birliğini kolaylaştırabilmektedir. Bununla birlikte siyasal hassasiyetler bu süreci zorlaştırabilmektedir. Belirli koşullar altında gerilimler bu nedenle daha görünür hâle gelebilmektedir.

Küçük Kent Ölçeğinin Analitik Değeri

Küçük kentlerin göç araştırmaları açısından değeri yalnızca nüfus büyüklüklerine bağlı değildir. Belirleyici olan, bu kentlerin kendine özgü bir toplumsal ve kurumsal yoğunluğa sahip olmasıdır. Glick Schiller ve Çağlar (2009), yerelliğin göç araştırmalarında karşılaştırmalı olarak teorileştirilmesi gerektiğini savunmaktadır. Yazarlara göre kentler, göçmen kadın ve erkeklerin yerleştiği basit mekânlar değildir. Bir kentin büyüklüğü, kurumlarla ve ekonomik olanaklarla kurulan teması etkilemektedir. Ayrıca toplumsal ağlar ve yerel iktidar ilişkileri üzerinde de etkili olmaktadır. Bu yaklaşım, küçük kentler açısından özellikle açıklayıcıdır. Bu kentlerde ekonomik olanaklar çoğu zaman sınırlıdır. Kurumsal ağlar da dar kapsamlı olabilmektedir. Bununla birlikte toplumsal ilişkiler özellikle yoğun olabilmektedir. Bu faktörler, göçmen kadın ve erkeklerin deneyimlerini doğrudan etkilemektedir.

Kreichauf'un (2023) Doğu Almanya'daki küçük bir kent üzerine gerçekleştirdiği çalışma bu bağlamda özellikle dikkat çekicidir. Çalışma, küçük kentlerin göç yönetişimi içinde "bekleme alanlarına" veya "transit mekânlarına" nasıl dönüşebildiğini göstermektedir. Ulusal dağıtım politikaları, göçmen kadın ve erkekleri kendi istekleri dışında küçük kentlere yönlendirebilmektedir. Bu durumda küçük kent, kalıcı bir ikamet yeri olarak görülmeyebilmektedir. Aksine, göçmenlerin başka bir yere geçmeden önce kalmak zorunda oldukları bir mekâna dönüşebilmektedir. Bu nedenle Kreichauf'un çalışması önemli bir özelliği ortaya koymaktadır. Küçük kentler göç yönetişiminin dışında yer almamaktadır. Aksine, bu yönetişimin merkezinde bulunmaktadır. Bununla birlikte göçmen kadın ve erkekler bu kentlerde çoğu zaman kalıcı bir aidiyet geliştirmemektedir. Yerel aktörler ise aynı anda bu geçici durumla başa çıkmak zorunda kalmaktadır.

Bu tartışma Türkiye açısından da önem taşımaktadır. Türkiye'de göç yönetimi güçlü biçimde merkezîleşmiş bir yapıya sahiptir. Bununla birlikte birçok gündelik sorun il, ilçe ve mahalle düzeylerinde ele alınmaktadır. Bu süreçte farklı aktörler aynı yerel mekânda karşılaşmaktadır. Bu aktörler arasında il göç idaresi birimleri ve belediyeler yer almaktadır. Sosyal yardım kuruluşları, okullar ve sağlık kurumları da sürece dâhil olmaktadır. Diğer aktörler arasında yerel yöneticilerin, yani muhtarların büroları ve yerel sivil toplum kuruluşları bulunmaktadır. Kimi zaman göçmen toplulukları içerisindeki kişiler de aracı bir rol üstlenmektedir. Bu karşılaşmalar her zaman planlı ve açık bir koordinasyona yol açmamaktadır. Bazı sorunlar kişisel temaslar aracılığıyla çözülmektedir. Diğer durumlarda yetki ve sorumluluklar belirsiz kalmaktadır. Bu durumda kurumlar sorumluluğu birbirlerine devretmektedir. Göçmen kadın ve erkekler de kimi zaman formel temaslarını sınırlandırmaktadır. Bu yolla mümkün olduğunca az görünür kalmaya çalışmaktadırlar.

Bir diğer önemli faktör toplumsal görünürlüktür. Anonimlik, büyük kentlerde göçmen kadın ve erkeklere belirli bir hareket alanı sağlayabilmektedir. Buna karşılık küçük kentlerde kişiler daha kolay tanınabilmektedir. Bir kişinin kim olduğu ve nerede çalıştığı daha hızlı fark edilmektedir. Kişinin yaşadığı mahalle, kurumlarla kurduğu temaslar ve toplumsal ilişkileri de daha kolay gözlemlenebilmektedir. Glorius, Bürer ve Schneider (2021), bu durumun çift yönlü etkisine dikkat çekmektedir. Yoğun toplumsal ilişkiler ve toplumsal denetim, kırsal bölgelerde ve küçük yerleşim yerlerinde entegrasyonu destekleyebilmekte, ancak aynı zamanda

sınırlandırabilmektedir. Göçmen kadın ve erkeklerin kabul edilmesi yalnızca bireysel tutumlara bağlı değildir. Yerel normlar ve ekonomik olanaklar da önem taşımaktadır. Diğer faktörler arasında toplumsal sermaye, temasların sıklığı ve kamu kurumlarının tutumu yer almaktadır.

Görünmeyen Yönetişim: Resmî Politika ile Gündelik Uygulama Arasında

"Görünmeyen yönetişim" kavramı ilk bakışta resmî kurumların dışında işleyen enformel süreçleri çağrıştırabilir. Ancak burada önerilen anlam daha dikkatli kurulmalıdır. Görünmeyen yönetişim, devletin yokluğu veya kurumların bütünüyle devre dışı kalması değildir. Daha çok resmî politikanın yerel düzeyde uygulanırken aldığı fiilî biçimlere işaret etmektedir. Bir yasa, yönetmelik veya merkezî karar küçük bir kente ulaştığında aynı metin olarak kalmaktadır. Fakat uygulanma biçimi, yerel kurumların kapasitesi, çalışanların deneyimi, kurumlar arası ilişki, yerel siyasal atmosfer, göçmenlerin görünürlüğü ve yerel halkın tepkisi içinde değişebilmektedir.

Ager ve Strang'ın (2008) entegrasyon çerçevesi bu tartışma için yararlı bir zemin sunmaktadır. Yazarlar entegrasyonu istihdam, barınma, eğitim, sağlık, haklar, sosyal bağlar, sosyal köprüler, sosyal bağlantılar, dil ve kültürel bilgi ile güvenlik ve istikrar gibi farklı alanların ilişkisi içinde düşünmektedir. Bu çerçeve, entegrasyonu tek bir göstergeye indirgememesi bakımından önemlidir. Ancak küçük kentler açısından bu alanların her biri yerel kapasite ve gündelik aracılık ilişkileri içinde yeniden ele alınmalıdır. Örneğin sağlık hizmetine erişim yalnızca sağlık hakkının varlığıyla ilgili değildir. Göçmenin hangi kuruma gideceğini bilip bilmemesi, dil desteği bulup bulamaması, daha önce benzer bir başvuru yapan birini tanıyıp tanımaması veya kurum çalışanının göçmenlerle çalışma deneyimi de süreci etkileyebilmektedir.

Görünmeyen yönetişim tam olarak bu ara bağlantıları görünür kılmaktadır. Bu alan bazen kolaylaştırıcıdır. Bir öğretmenin çocuğun kaydı için ek çaba göstermesi, bir muhtarın aileyi ilgili kuruma yönlendirmesi veya yerel bir sivil toplum kuruluşunun tercüme ve danışmanlık sağlaması göçmenin kurumsal alana erişimini artırabilmektedir. Ancak aynı ağlar sınırlayıcı da olabilmektedir. Yerel toplumda olumsuz bir kanaat oluştuğunda, göçmenlerin bazı hizmetlere erişimi resmî olarak değil ama fiilî olarak zorlaşabilmektedir. Kurum çalışanları siyasal hassasiyet nedeniyle daha düşük görünürlüklü bir hizmet biçimini tercih edebilmektedir. Biehl ve diğerlerinin (2026) seçici görünürlük ve kişisel bağlara dayanma vurgusu, küçük kentler için bu nedenle açıklayıcıdır. Bu tür ilişkiler yönetişimi ortadan kaldırmamaktadır. Onu daha kırılgan, daha kişisel ve daha az kayıtlı hâle getirmektedir.

Betts, Memişoğlu ve Ali'nin (2021) Türkiye ve Lübnan'da belediye başkanlarının Suriyeli mültecilere yönelik politikalardaki rolünü tartışan çalışması da yerel aktörlerin merkezî politikaları yalnızca uygulamadığını göstermektedir. Yerel aktörler bazı durumlarda bu politikaları yorumlamakta veya tamamlayıcı pratikler geliştirmektedir. Bu bulgu büyükşehir belediyeleri üzerinden geliştirilmiş olsa da küçük kentler açısından da önemlidir. Çünkü yerel aktörlerin tutumu, kapasitesi ve siyasal konumu, göçmenlerin gündelik deneyiminde belirleyici olabilmektedir.

Yerel Kurumlar, Aracılar ve Kurumsal Sınırlar

Küçük kentlerde göç yönetişimi çoğu zaman kurumların resmî görevlerinin ötesine geçmektedir. Farklı aktörler bu sürece farklı ölçülerde dâhil olmaktadır. Bu aktörler arasında belediyeler ve il göç idaresi birimleri yer almaktadır. Sosyal yardım ve dayanışma vakıfları, okullar ve aile sağlığı merkezleri de sürece katkı sağlamaktadır. Hastaneler, yerel yöneticilerin, yani muhtarların büroları, yerel dernekler ve gönüllü ağlar da rol oynamaktadır. Bununla birlikte bu aktörlerin sürece katılması, kendiliğinden düzenli bir yönetişim yapısının bulunduğu anlamına gelmemektedir. Küçük kentlerde kurumlar arasındaki koordinasyon kimi zaman kişisel ilişkilere dayanmaktadır. Bir kurumda çalışan kişi, başka bir kurumda tanıdığı bir kişiyi telefonla arayabilmektedir. Bu sayede bir başvuru daha hızlı sonuçlandırılabilmektedir. Aynı zamanda sistemin kişisel temaslara bağımlı hâle gelmesi riski de bulunmaktadır.

OECD'nin (2018) yerel entegrasyona ilişkin raporu, çeşitli önemli faktörleri vurgulamaktadır. Bu faktörler arasında farklı yönetim düzeyleri arasındaki koordinasyon yer almaktadır. Yerel verilerin toplanması da önem taşımaktadır. Ayrıca hizmetlerin ilgili yerin büyüklüğüne ve koşullarına uyarlanması gerekmektedir. Bir diğer faktör ise devlet dışı aktörlerle iş birliğidir. Bu öneriler küçük kentler için de geçerlidir. Bununla birlikte kapasite eksikliği küçük kentlerde daha belirgin olabilir. Göçmen kadın ve erkeklere yönelik uzmanlaşmış bir birim bulunmayabilir. Dil desteği sınırlı kalabilir. Yerel sivil toplum kuruluşlarının sayısı da az olabilir. Bu nedenle bir boşluk ortaya çıkabilmektedir. Bir tarafta politika belgelerinde öngörülen hizmetler yer almaktadır. Diğer tarafta ise sahada fiilen sunulabilen destek bulunmaktadır.

Bu boşluk her zaman açık bir başarısızlık olarak değerlendirilmemelidir. Küçük kentler, sınırlı kapasitelerine rağmen kimi zaman hızlı ve esnek çözümler geliştirebilmektedir. Yerel aktörler çoğu zaman birbirlerini kişisel olarak tanımaktadır. Bu nedenle sorunlar kısa iletişim kanalları aracılığıyla çözülebilmektedir. Yerel kurumlar göçmen aileleri daha yakından takip ederek destekleyebilmektedir. Bununla birlikte aynı yakınlık sorunlu sonuçlara da yol açabilmektedir. Hizmetler belirli kişilere daha fazla bağımlı hâle gelebilmektedir. Geçerli standartlar da bu nedenle belirsizleşebilmektedir. Kişisel bir tanışıklık, bir göçmen kadının veya erkeğin hizmetlere erişimini kolaylaştırabilmektedir. Başka bir kişi ise gerekli temaslara sahip olmadığı için dışarıda kalabilmektedir. Görünmeyen yönetişimin eleştirel önemi bu noktada ortaya çıkmaktadır. Görünmeyen yönetişim ne idealize edilmeli ne de bütünüyle olumsuz değerlendirilmelidir. Belirleyici olan, yerel pratiklerin hangi koşullar altında kapsayıcı sonuçlar ürettiğidir. Bu pratiklerin ne zaman seçici veya dışlayıcı biçimde işlediğinin de incelenmesi gerekmektedir.

Jonitz ve diğerleri (2026), belediyelerin, sivil toplum kuruluşlarının ve yerel nüfusun rollerinin bağlama göre farklılaştığını ortaya koymaktadır. Bazı küçük kentlerde yoğun iş birliği ağları oluşmaktadır. Diğer yerlerde ise yönetişimin parçalı ve pasif biçimleri gelişmektedir. Bu farklılıklar, tek tip bir kategorinin kullanılmasının yol açabileceği sorunları göstermektedir. Küçük kentler birbirine benzeyen mekânlar olarak değerlendirilmemelidir. Gümüşhane, Trabzon, Konya, Samsun ve Türkiye'deki benzer küçük veya orta büyüklükteki kentler de aynı kurumsal ve toplumsal koşullara sahip değildir. Bu nedenle küçük kentlerde göç yönetişimine ilişkin çalışmalar, bu kentlerin homojen olduğu varsayımından hareket etmemelidir. Bunun yerine karşılaştırmalı ve bağlama duyarlı bir yaklaşım benimsenmelidir. Bu yolla yerel farklılıkların görünür kılınması mümkün olacaktır.

Toplumsal Meşruiyet, Görünürlük ve Sosyal Mesafe

Göç yönetişimi, yalnızca hizmet sunumunu kapsamamaktadır. Göçmenlerin yerel toplum tarafından kabul edilme düzeyi, hizmetlerin hangi biçimde sunulduğunu etkileyebilmektedir. Ayrıca göçmenlerin bu hizmetlerden nasıl yararlandıkları üzerinde de etkili olmaktadır. Küçük kentlerde göçmen kadın ve erkekler çoğu zaman daha görünür durumdadır. Bu nedenle toplumsal kabul meselesi bu kentlerde daha doğrudan algılanmaktadır. Büyük bir kentte göçmen bir aileye destek sunulması, sıradan bir sosyal yardım işlemi olarak kalabilmektedir. Buna karşılık küçük bir kentte bu durum yerel nüfusun dikkatini çekebilmektedir. Bu da kurumların daha temkinli hareket etmesine yol açabilmektedir. Bu nedenle Biehl ve diğerlerinin (2026) çalışması özellikle önem taşımaktadır. Söz konusu çalışma, Türkiye'deki küçük kentlerde siyasal risklerin, kamusal görünürlüğün ve enformel aracıların önemini vurgulamaktadır.

Toplumsal meşruiyet, yerel nüfusun göçle ilişkili tedbirlere yönelik tepkilerini ifade etmektedir. Bununla birlikte toplumsal meşruiyet yalnızca destek veya ret üzerinden anlaşılmamalıdır. Küçük kentlerde kabul çoğu zaman belirli koşullara bağlı olmaktadır. Göçmen kadın ve erkeklerden "sorun çıkarmamaları" beklenebilmektedir. Ayrıca "dikkat çekmemeleri" ve "yerel normlara uymaları" da beklenebilmektedir. "Ekonomik rekabet yaratmamaları" yönündeki

beklenti de bu süreçte rol oynayabilmektedir. Bu tür beklentiler her zaman açık bir dışlanmaya yol açmamaktadır. Bununla birlikte toplumsal mesafeyi koruyan yerel bir düzen oluşturabilmektedir. Bu düzen, göçmen kadın ve erkeklerin kamusal görünürlüğünü sınırlandırabilmektedir. Ayrıca kurumlarla kurulan ilişkilerin daha temkinli hâle gelmesine yol açabilmektedir.

Glorius ve diğerleri (2021), yerel kabulün yalnızca göçmen kadın ve erkeklerin entegrasyon çabaları aracılığıyla açıklanamayacağını vurgulamaktadır. Kabul eden toplumun tutumlarının da dikkate alınması gerekmektedir. Bu bakış açısı, entegrasyonu tek taraflı bir süreç olarak ele alan yaklaşımlara yönelik önemli bir eleştiri sunmaktadır. Bu bağlamda Ager ve Strang'ın (2008) yaptığı ayrım da yeniden değerlendirilebilir. Yazarlar sosyal bağlar (social bonds), sosyal köprüler (social bridges) ve sosyal bağlantılar (social links) arasında ayrım yapmaktadır. Küçük kentlerde göçmen kadın ve erkekler kendi toplulukları içinde sınırlı bağlar kurabilmektedir. Yerel nüfusla temasları zayıf, ancak işlevsel nitelikte olabilmektedir. Kurumlarla ilişkiler ise çoğunlukla somut bir desteğe ihtiyaç duyulduğunda kurulabilmektedir. Bu ilişkiler çatışmasız bir düzenin oluşmasına olanak sağlayabilmektedir. Bununla birlikte çatışmanın bulunmaması, güçlü bir toplumsal bütünleşmenin mevcut olduğu anlamına gelmemektedir.

Küçük kentlerde görünürlüğün iki yönü bulunmaktadır. Kişisel olarak tanınmak güven oluşturabilmekte ve hizmetlere erişimi kolaylaştırabilmektedir. Yerel esnaf, muhtarlar, öğretmenler veya belediye çalışanları bir kişiyi tanıdığında bazı sorunlar daha hızlı çözülebilmektedir. Bununla birlikte tanınmak, gözlemlenmek anlamına da gelebilmektedir. Göçmen kadın ve erkeklerin davranışları, aile ilişkileri ve çalışma biçimleri daha ayrıntılı biçimde değerlendirilebilmektedir. Aynı durum onların kamusal alandaki varlıkları için de geçerlidir. Bu nedenle küçük kentlerde görünürlük hem bir kaynak hem de bir baskı biçimi olabilmektedir. Görünmeyen yönetişim kavramı, bu çift yönlü etkiyi açıklamayı amaçlamaktadır.

Tartışma: Küçük Kentleri Pasif Uygulama Alanı Olarak Görmemek

Bu çalışmanın temel tezi, küçük kentlerin göç yönetişiminin pasif mekânları olmadığıdır. Küçük kentler, ulusal tedbirlerin değişikliğe uğramadan uygulandığı nötr yerler olarak değerlendirilmemelidir. Merkezî düzeyde alınan kararlar yerel düzeyde yeniden şekillendirilmektedir. Bu süreçte kurumsal kapasiteler ve yerel aktörlerin girişimleri önemli bir rol oynamaktadır. Toplumsal görünürlük, enformel ağlar ve meşruiyet meseleleri de bu süreci etkilemektedir. Bu değişimler resmî politika belgelerinde her zaman görünür hâle gelmemektedir. Çoğu zaman gündelik çözüm pratiklerinde ortaya çıkmaktadır. Bu pratikler arasında kurumlar arasındaki telefon görüşmeleri ve yerel bir muhtar aracılığıyla gerçekleştirilen aracılık yer almaktadır. Bir öğretmenin çabaları da bu süreçte rol oynayabilmektedir. Bir diğer örnek, bir göçmen kadının veya erkeğin mümkün olduğunca az görünür kalmaya karar vermesidir.

Bu yaklaşım, çok düzeyli yönetişim literatürünün küçük kentler perspektifinden yeniden ele alınmasını gerektirmektedir. Çok düzeyli yönetişim yalnızca çok sayıda aktörün sürece dâhil olması anlamına gelmemektedir. Bu aktörler arasındaki ilişkiler de önem taşımaktadır. Diğer unsurlar arasında yetkilerin dağılımı, kaynaklara erişim ve siyasal risklerin algılanması yer almaktadır. Biehl ve diğerleri (2026), bu durumu Türkiye örneği üzerinden ortaya koymaktadır. Yerel düzey, merkezî bir sistem içerisinde dahi bütünüyle etkisiz değildir. Bununla birlikte yerel aktörlerin hareket alanı sınırlı, enformel ve seçici olabilmektedir. Bu nedenle küçük kentlerde yönetişim daha az kurumsallaşmış olabilir. Aynı zamanda kişisel ve kurumsal ilişkiler tarafından daha güçlü biçimde şekillendirilebilir.

Kreichauf (2023), küçük kentleri transit ve bekleme alanları olarak tanımlamaktadır. Bu yaklaşım, göçmen kadın ve erkeklerin küçük kentlerle her zaman kalıcı bir aidiyet ilişkisi geliştirmediğini hatırlatmaktadır. Bazı göçmenler açısından bu kentler, zorunlu olarak kalınan

yerler olabilir. Ayrıca daha büyük kentlere geçiş öncesinde geçici bekleme alanları veya ara duraklar olarak da işlev görebilirler. Bu durum, yerel kurumların uzun vadeli entegrasyon tedbirleri geliştirme isteğini azaltabilmektedir. Aynı zamanda yerel nüfusta "geçici misafir" anlayışını güçlendirebilmektedir. Türkiye'de de geçicilik, entegrasyon ve geri dönüş üzerine yürütülen tartışmalar yerel kurumların davranışlarını etkilemektedir. Bu nedenle küçük kentlerde göç yönetişiminin incelenmesi, yerel hizmetlerden daha fazlasını dikkate almalıdır. Göçmen kadın ve erkeklerin yerel mekânla kurduğu zamansal ilişkinin de incelenmesi gerekmektedir.

Sonuç

Küçük kentlerde göç yönetişimi, formel yasaların, merkezî politikaların ve kurumsal görevlerin ötesine geçen gündelik bir alanı kapsamaktadır. Bu alan, raporlarda ve politika belgelerinde çoğu zaman yer almadığı için görünmeyen durumdadır. Kurumsal performans göstergeleri de bu alanı çoğunlukla kapsamamaktadır. Bununla birlikte düşük görünürlük, bu alanın etkisiz olduğu anlamına gelmemektedir. Hizmetlere erişim çoğu zaman bu ara alanda şekillenmektedir. Aynı durum yerel toplumla ilişkiler, kurumlara duyulan güven ve kamusal alandaki görünürlük için de geçerlidir.

Bu çalışma "görünmeyen yönetişim" kavramını önermektedir. Bu kavram aracılığıyla küçük kentlerde göç yönetişiminin fiilî pratiklerinin incelenmesi amaçlanmaktadır. Kavram, kurumların hangi görevleri yerine getiremediğini görünür kılmaktadır. Ayrıca çalışanların kendi imkânlarıyla neleri gerçekleştirdiğini de ortaya koymaktadır. Bunun yanı sıra yerel aracıların hangi durumlarda devreye girdiğini kapsamaktadır. Gündelik ilişkiler aracılığıyla kolaylaştırılan veya güçleştirilen süreçler de bu kavram çerçevesinde analiz edilebilmektedir. Bununla birlikte kavram, özünde olumlu bir alanı ifade etmemektedir. Aksine daha ayrıntılı biçimde incelenmesi gereken karmaşık bir süreci tanımlamaktadır. Görünmeyen yönetişim hizmetlere erişimi kolaylaştırabilmektedir. Ancak eşitsizlikleri de güçlendirebilmektedir. Bazı durumlarda toplumsal gerilimleri azaltabilmektedir. Diğer durumlarda ise toplumsal denetim üreterek göçmen kadın ve erkeklerin görünürlüğünü sınırlandırabilmektedir.

Bu nedenle küçük kentler, göç araştırmalarında yalnızca şimdiye kadar ihmal edilmiş ampirik araştırma alanları değildir. Aynı zamanda önemli analitik mekânlardır. Göç yönetişiminin, toplumsal bütünleşmenin ve yerel kapasitelerin yeniden değerlendirilmesine olanak sağlamaktadırlar. Aynı durum enformel kurumsallaşma ve toplumsal meşruiyet için de geçerlidir. Bu çalışma, bir sempozyum bildirisi olarak kavramsal ve keşfedici bir çerçeve sunmaktadır. Daha sonraki bir makalede bu çerçeve saha verileri aracılığıyla sınanabilir. Bu amaçla Gümüşhane, Trabzon veya benzer küçük kentlerde gerçekleştirilecek araştırmalar uygun olacaktır. Bu yolla kavramın açıklayıcılık gücü daha kesin ve güvenilir biçimde değerlendirilebilecektir.

Kaynakça

Ager, A., & Strang, A. (2008). Understanding integration: A conceptual framework. *Journal of Refugee Studies, 21*(2), 166–191. https://doi.org/10.1093/jrs/fen016

Betts, A., Memişoğlu, F., & Ali, A. (2021). What difference do mayors make? The role of municipal authorities in Turkey and Lebanon's response to Syrian refugees. *Journal of Refugee Studies, 34*(1), 491–519. https://doi.org/10.1093/jrs/feaa011

Biehl, K. S., Açıkgöz, M., Eren Benlisoy, Z. C., & İkizoğlu Erensü, A. (2026). Integration and multi-level governance in Turkey's small towns: An actor centred analysis. *International Migration, 64*, e70163. https://doi.org/10.1111/imig.70163

Caponio, T., & Borkert, M. (Eds.). (2010). *The local dimension of migration policymaking*. Amsterdam University Press. https://doi.org/10.5117/9789089642325

Garcés-Mascareñas, B., & Penninx, R. (Eds.). (2016). *Integration processes and policies in Europe: Contexts, levels and actors*. Springer Open. https://doi.org/10.1007/978-3-319-21674-4

Glick Schiller, N., & Çağlar, A. (2009). Towards a comparative theory of locality in migration studies: Migrant incorporation and city scale. *Journal of Ethnic and Migration Studies, 35*(2), 177–202. https://doi.org/10.1080/13691830802586179

Glorius, B., Bürer, M., & Schneider, H. (2021). Integration of refugees in rural areas and the role of the receiving society: Conceptual review and analytical framework. *Erdkunde, 75*(1), 51–60. https://doi.org/10.3112/erdkunde.2021.01.04

Jonitz, E., Schiller, M., & Scholten, P. (2026). Are small towns a battleground for migration governance? Negotiating refugee integration in small towns in the Netherlands. *Local Government Studies, 52*(3), 513–535. https://doi.org/10.1080/03003930.2024.2357306

Kreichauf, R. (2023). Governing migration through small towns: Dispersal and the production of spaces of transit. *International Journal of Urban and Regional Research, 47*(3), 349–367. https://doi.org/10.1111/1468-2427.13171

OECD. (2018). *Working together for local integration of migrants and refugees*. OECD Publishing. https://doi.org/10.1787/9789264085350-en

Scholten, P., & Penninx, R. (2016). The multilevel governance of migration and integration. In B. Garcés-Mascareñas & R. Penninx (Eds.), *Integration processes and policies in Europe: Contexts, levels and actors* (pp. 91–108). Springer Open. https://doi.org/10.1007/978-3-319-21674-4_6

Zapata-Barrero, R., Caponio, T., & Scholten, P. (2017). Theorizing the "local turn" in a multi-level governance framework of analysis: A case study in immigrant policies. *International Review of Administrative Sciences, 83*(2), 241–246. https://doi.org/10.1177/0020852316688426

ŞƏRQ REZONANSI VƏ AZƏRBAYCAN ƏDƏBİ FİKİR TARİXİNDƏ ƏLİŞİR NƏVAİ İRSİNİN SOSİO-MƏDƏNİ TƏSİR SPETKRİ

EASTERN RESONANCE AND THE SOCIO-CULTURAL IMPACT SPECTRUM OF ALISHER NAVOI'S HERITAGE IN THE HISTORY OF AZERBAIJANI LITERARY THOUGHT

Gunel Hummetova (Paniyeva) Saleh
Azerbaijan State Pedagogical University, Senior Lecturer at the Department of Literature Teaching Technology, Ph.D. in Philology
ORCID ID: 0000-0002-8916-3484

XÜLASƏ

Bu tədqiqat işində böyük özbək mütəfəkkiri, dövlət xadimi və Şərq poeziyasının zirvələrindən biri olan Mir Əlişir Nəvainin elmi-ədəbi irsi və onun Azərbaycan mədəniyyəti, ədəbi mühiti və resenziya tarixi ilə qarşılıqlı əlaqələri təhlil olunur. Tədqiqat göstərir ki, Nəvai irsi təkcə özbək xalqının deyil, bütöv bir Türk-İslam coğrafiyasının ortak mənəvi xəzinəsidir. Məqalədə Nəvainin fəlsəfi dünyagörüşü, fəlsəfə və poeziya sintezi, "Xəmsə"çilik ənənəsi daxilində Nizomi Gəncəvi məktəbinə sadiqliyi və bu xəttin Məhəmməd Füzuliyə qədər uzanan təsir spektri araşdırılır.

Bununla yanaşı, Nəvainin Azərbaycan ədəbi mühitində — Kişvəri, Xülqi kimi XV-XVI əsr liriklərinin yaradıcılığında, Sadiq bəy Əfşarın "Məcməül-xəvas" təzkirəsində və M.F.Axundzadənin maaarifçi düşüncəsindəki yeri faktlarla əsaslandırılır. Məqalədə XX əsrdən etibarən Azərbaycanda Nəvai irsinin nəşri, dərsliklərə salınması, 1926-cı il Bakı yubileyi və şairin adının əbədiləşdirilməsi məsələlərinə xüsusi yer verilmişdir.

Bəşəriyyətin mənəvi tərəqqisinə misilsiz töhfələr bəxş etmiş klassik sənətkarların yaradıcılığı zaman və məkan sərhədlərini aşaraq ümumbəşəri estetik dəyərə çevrilir. Əvəzsiz söz ustadı, alim, görkəmli dövlət xadimi Nizaməddin Mir Əlişir Nəvai (1441–1501) məhz belə dühallardandır. Şərq intibahının (Renessansının) parlaq simalarından biri olan Nəvai, bədii təfəkkürlə fəlsəfi dünyagörüşünü möhtəşəm bir sintezdə birləşdirmişdir. Əgər poeziya onun mənəvi varlığının və emosional dünyasının ifadəsi idisə, fəlsəfə bu poetik binanın sarsılmaz təməli rolunu oynayırdı.

Nəvai özündən əvvəlki antik, Şərq və İslam fəlsəfəsi irsini dərindən mənimsəmiş, Yunanıstan, Hindistan, Misir, Çin, İran və Mərkəzi Asiyanın intellektual xəzinəsindən bəhrələnərək onları öz unikal təfəkkür süzgəcindən keçirmişdir. O, sələflərinin irsinə dərin ehtiramla yanaşmış, öz fəlsəfi-bədii konsepsiyasını məhz bu zəngin irs üzərində ucaltmışdır. Nəvai bu maarifçi və təvazökar yanaşmasını özünün məşhur "Məcalisün-nəfais" (Nəfislər məclisi) təzkirəsində belə etiraf edirdi:

"Biz dahi söz ustalarının zəmisindəki yerə tökülmüş sümbülləri yığırıq. Bəlkə heç ona da layiq deyilik..."

Bu fəlsəfi təvazökarlıq, əslində, Nəvai dühosunun böyüklüyünü və onun Şərq söz sənətinə gətirdiyi humanist məsuliyyəti nümayiş etdirir. Nəvai ana dili hesab etdiyi türk (çağatay) dilinin zəngin poetik imkanlarını sübut etmək üçün "Mühakimətül-lüğəteyn" (İki dilin müqayisəsi) əsərini qələmə almış, türk dilini saray və elmi dil səviyyəsinə qaldırmışdır.Əlişir Nəvai 7-8 yaşlarından bədii yaradıcılığa başlamış, yarım əsrdən artıq davam edən zəngin yaradıcılıq yolu keçmişdir. O, Yaxın və Orta Şərq ədəbiyyatının demək olar ki, bütün lirik janrlarında (qəzəl, rübai, qıtə, məsnəvi, tuyuq) şah əsərlər yaratmışdır. Şairin lirik irsinin zirvəsini onun 25 min beytə yaxın olan və dövrün humanist ənənələrinə uyğun olaraq ömrünün dörd mərhələsini əks etdirən "Xəzainül-maani" (Mənalar xəzinəsi) — "Çar divan"ı ("Dörd divan") təşkil edir:

1. *Gərayibüs-siğər* (Uşaqlıq qəribəlikləri)
2. *Nəvadirüş-şəbab* (Gənclik nadirlikləri)
3. *Bədayiül-vəsət* (Orta yaş bədiiyyatı)
4. *Fəvahidül-kibər* (Qocalıq faydaları)

Nəvai poetikasının qüdrəti onun lirik və fəlsəfi təsvir vasitələrinin elastikliyindədir. Məsələn, qələmin gücünü, fikrin sürətini və obrazlılığını tərənnüm edərkən şair belə bədii təsvirə əl atır:

"Qələmdir yel kimi sürətlə gedən,
Əzəldən yüksəkdir yeri fələkdən.
Vəhmin ədhəmindən daha tez çapar,
Onun Şəbdiz kimi bir surəti var...
Ayağından etmiş başını qələm."

Burada qələm obrazı təkcə yazı aləti deyil, insan zəkası, fikrin sürəti və əbədi yaşar sənətin simvolu kimi təqdim olunur.

Əlişir Nəvai irsi XV əsrdən etibarən Azərbaycan ədəbi mühiti tərəfindən böyük rəğbətlə qarşılanmış və öyrənilmişdir. Dahi Nizami Gəncəvinin "Xəmsə" ideyalarını türk dilində yenidən zirvəyə qaldıran Nəvai, özü də Azərbaycan şairləri üçün qüdrətli bir ustad məktəbinə çevrilmişdir.Nəvainin çağdaşları olan Azərbaycan şairləri onun dil və poeziya məktəbindən bəhrələnmişlər. XV əsr Azərbaycan şairi Kişvəri öz divanında Nəvai dühasına açıq şəkildə istinad edir, onun qəzəllərinə nəzirələr yazırdı.Böyük Məhəmməd Füzuli isə özünün "Leyli və Məcnun" poemasının müqəddiməsində türk dilli poeziyanın banilərindən bəhs edərkən Nəvainin adını böyük ehtiramla çəkir. Füzuli Nəvainin türk dilində yaratdığı möhtəşəm abidəni özünə örnək bilmiş, onun "Xəmsə"sindəki "Leyli və Məcnun" motivini daha da dərinləşdirərək psixoloji-fəlsəfi zirvəyə qaldırmışdır.Azərbaycan təzkirəçilik məktəbinin görkəmli nümayəndəsi, Səfəvilər dövrünün görkəmli fırça ustası və alimi Sadiq bəy Əfşar özünün "Məcməül-xəvas" (Xasların məclisi) adlı təzkirəsini doğrudan-doğruya Nəvainin "Məcalisün-nəfais" əsərinin quruluş və üslubunda qələmə almışdır. Əfşar əsərin giriş hissəsində Nəvaiyə xüsusi mədhiyyə həsr edərək, onu söz sənətinin böyük piri kimi dəyərləndirmişdir.XIX əsrdə Azərbaycan maarifçiliyinin təməlini qoyan Mirza Fətəli Axundzadə öz publisistik və fəlsəfi məqalələrində Nəvai şəxsiyyətinə toxunaraq, onu təkcə dahi şair kimi deyil, ədalətli, müdrik dövlət xadimi və xalqın rifahı üçün çalışan maarifçi vizyon sahibi kimi xarakterizə etmişdir.XX əsrdə Azərbaycanda Əlişir Nəvai irsinin tədqiqi və təbliği yeni keyfiyyət mərhələsinə qədəm qoymuşdur:

- **1926-cı il Yubileyi:** Bakıda keçirilən I Türkoloji Qurultay ərəfəsində Ə.Nəvainin anadan olmasının 500 illiyi paytaxt Bakıda böyük təntənə ilə qeyd edilmişdir. Bu dövrdə şairin "Müntəxəbat" və "Vəqfiyyə" əsərləri ilk dəfə Bakı nəşriyyatlarında işıq yüzü görmüş, eyni zamanda Hüseyn Bayqaranın "Divan"ı da çap olunmuşdur.
- **Tərcümə və Tədris:** 1948-ci ildə "Azərnəşr" tərəfindən Nəvainin "Fərhad və Şirin" poemasının minacat və nət hissələri, "Uşaqgəncnəşr" tərəfindən isə "Səb'eyi-səyyar" (Yeddı səyyar) əsərindən parçalar Azərbaycan dilində oxuculara təqdim edilmişdir. Azərbaycan məktəbləri üçün yazılan ilk dərsliklərdən başlayaraq Nəvainin həyatı və şeirləri proqramlara daxil edilmişdir.
- **Toponomik və Memarlıq Ehtiramı:** Bakı şəhərinin Nərimanov rayonunda Əlişir Nəvaiyə görkəmli abidə ucaldılmış, Hacıqabul rayonunda isə şairin adını daşıyan qəsəbə və küçə salınmışdır.

Əlişir Nəvai yaradıcılığı zamanın və coğrafiyanın darlığına sığmayan, bütün bəşəriyyətə aid olan böyük bir mənəvi sərvətdir. Şərq hikmətinin dediyi kimi: *"Bütün canlılar ölür, bütün cansızlar — yəni elm, bədii söz və sənət isə əbədi yaşayır."*

Nəvai irsi Azərbaycan xalqı tərəfindən XV əsrdən bu günə qədər öz doğma söz ustadının irsi kimi ehtiramla öyrənilmiş və sevilmişdir. Nəvai və Nizami, Nəvai və Füzuli arasındakı bədii-fəlsəfi varislik telləri göstərir ki, Türk-İslam dünyasının estetik təfəkkürü vahid humanist dəyərlər üzərində bərqərar olmuşdur. Nəvainin 580 illik yubileyi və sonrakı anım mərasimləri bu böyük humanist mütəfəkkirin ideyalarının müasir dövrümüzdə də öz aktuallığını və nurunu koruması deməkdir.

Açar sözlər: *Əlişir Nəvai, Azərbaycan-özbək ədəbi əlaqələri, Türk dili poeziyası, "Çar divan", Məhəmməd Füzuli, maarifçilik, retseptiv estetika.*

ABSTRACT

This study analyzes the scientific and literary heritage of the great Uzbek thinker, statesman, and one of the peaks of Eastern poetry, Mir Alisher Navoi, along with his interactions with Azerbaijani culture, literary environment, and reception history. The research shows that Navoi's legacy is a shared spiritual treasure not only for the Uzbek people but for the entire Turkic-Islamic geography. The article examines Navoi's philosophical worldview, his synthesis of philosophy and poetry, his adherence to the school of Nizami Ganjavi within the *"Khamsa"* (Quintet) tradition, and the spectrum of this influence extending all the way to Muhammad Fuzuli.

In addition, the paper substantiates with concrete facts Navoi's place in the Azerbaijani literary environment—specifically in the works of 15th–16th-century lyricists like Kishvari and Khulqi, in Sadiq Beg Afshar's biographical anthology (*tazkira*) *"Majma al-Khawas"*, and in the enlightenment thought of Mirza Fatali Akhundzadeh. Special attention is given to the publication of Navoi's works in Azerbaijan starting from the 20th century, their inclusion in school textbooks, the 1926 Baku jubilee, and the measures taken to perpetuate the poet's name.

MAIN TEXT

The creative works of classic masters who have made invaluable contributions to the spiritual progress of humanity cross the boundaries of time and space, transforming into universal aesthetic values. Nizam al-Din Mir Alisher Navoi (1441–1501)—an irreplaceable master of the word, scholar, and prominent statesman—is precisely one such genius. As one of the brilliant figures of the Eastern Renaissance, Navoi magnificently synthesized artistic thought with a philosophical worldview. If poetry was the expression of his spiritual existence and emotional world, philosophy served as the unshakeable foundation of this poetic structure.

Navoi deeply mastered the ancient, Eastern, and Islamic philosophical heritage preceding him, drawing upon the intellectual treasures of Greece, India, Egypt, China, Iran, and Central Asia, filtering them through his own unique intellect. He approached the legacy of his predecessors with profound respect, building his own philosophical-artistic concept upon this rich foundation. Navoi modestly acknowledged this enlightened approach in his famous biographical anthology *"Majalis al-Nafais"* (*The Assemblies of the Choice Spirits*):

"We are merely gathering the ears of wheat left fallen on the fields of the great masters of the word. Perhaps we are not even worthy of that..."

This philosophical humility actually demonstrates the greatness of Navoi's genius and the humanist responsibility he brought to Eastern verbal art. To prove the rich poetic capabilities of the Turkic (Chagatai) language, which he considered his native tongue, Navoi penned *"Muhakamat al-Lughatayn"* (*The Trial of the Two Languages*), thereby elevating Turkic to the status of a court and scientific language.

Alisher Navoi began his artistic activity at the age of 7–8 and pursued a rich literary career lasting over half a century. He created masterpieces in almost all lyrical genres of Near and Middle Eastern literature (ghazal, rubai, qita, masnavi, tuyugh). The apex of the poet's lyrical heritage is his *"Khazain al-Ma'ani"* (*Treasuries of Meanings*)—the *"Char Divan"* (*Four*

Divans)—comprising nearly 25,000 couplets (*beits*), which reflects four stages of his life in accordance with the humanist traditions of his time:

1. **Ghara'ib al-Sighar** (*Wonders of Childhood*)
2. **Nawadir al-Shabab** (*Rarities of Youth*)
3. **Bada'i al-Wasat** (*Marvels of Middle Age*)
4. **Fawa'id al-Kibar** (*Benefits of Old Age*)

The power of Navoi's poetics lies in the flexibility of his lyrical and philosophical means of expression. For instance, when chanting the power of the pen, the speed of thought, and imagery, the poet employs the following artistic description:

"The pen is like the wind, moving at great speed,
From eternity past, its place is higher than the heavens.
It runs faster than the dark steed of imagination,
It bears an appearance like Shabfiz...
It has turned its head into a pen from its very feet."

Here, the image of the pen is presented not merely as a writing tool, but as a symbol of human intellect, the swiftness of thought, and everlasting art.

NAVOI'S HERITAGE IN AZERBAIJANI LITERARY THOUGHT

From the 15th century onward, Alisher Navoi's heritage was received with great admiration and studied in the Azerbaijani literary environment. Elevating the *"Khamsa"* ideas of the great Nizami Ganjavi to new heights in the Turkic language, Navoi himself became a powerful master school for Azerbaijani poets.

- **15th–16th Century Lyricists:** Contemporary Azerbaijani poets drew inspiration from Navoi's language and poetic school. The 15th-century Azerbaijani poet Kishvari explicitly referenced Navoi's genius in his *Divan* and wrote *naziras* (poetic responses/parallel poems) to his ghazals.
- **Muhammad Fuzuli:** In the preface to his poem *"Leyli and Majnun"*, when discussing the founders of Turkic-language poetry, the great Muhammad Fuzuli mentioned Navoi's name with deep reverence. Fuzuli took the magnificent monument Navoi built in the Turkic language as a model, deepening the *"Leyli and Majnun"* motif from Navoi's *"Khamsa"* and elevating it to a psychological-philosophical peak.
- **Sadiq Beg Afshar:** A prominent representative of the Azerbaijani biographical tradition, a master painter, and a scholar of the Safavid era, Sadiq Beg Afshar wrote his anthology *"Majma al-Khawas"* (*Assembly of the Elite*) directly modeled after the structure and style of Navoi's *"Majalis al-Nafais"*. In the introductory section of the work, Afshar dedicated a special panegyric to Navoi, evaluating him as a great master (*pir*) of the art of words.
- **Mirza Fatali Akhundzadeh:** In the 19th century, Mirza Fatali Akhundzadeh, who laid the foundation of Azerbaijani enlightenment, touched upon Navoi's personality in his journalistic and philosophical essays. He characterized Navoi not only as a genius poet but also as a just, wise statesman and an enlightened visionary who worked for the welfare of the people.

RESEARCH AND COMMEMORATION IN THE 20TH CENTURY AND BEYOND

During the 20th century, the study and promotion of Alisher Navoi's heritage in Azerbaijan entered a new qualitative phase:

- **The 1926 Jubilee:** On the eve of the First Turkological Congress held in Baku, the 500th anniversary of A. Navoi's birth was celebrated with great solemnity in the capital city of Baku. During this period, Navoi's works *"Muntakhabat"* (Selected Works) and *"Waqfiyya"* were published for the first time by Baku publishing houses, along with Hussein Bayqara's *"Divan"*.
- **Translation and Education:** In 1948, the *minajat* (prayers) and *na'at* (praises to the Prophet) sections of Navoi's poem *"Farhad and Shirin"* were published in Azerbaijani by "Azerneshr", and excerpts from *"Sab'ai Sayyar"* (*Seven Planets*) were presented to readers by

"Ushaqganjneshr". Starting from the very first textbooks written for Azerbaijani schools, Navoi's life and poetry were integrated into the educational curricula.

- **Toponymic and Architectural Homage:** A prominent monument to Alisher Navoi was erected in the Narimanov district of Baku, and a settlement and street bearing the poet's name were established in the Hajigabul district.

CONCLUSION

Alisher Navoi's creative legacy is a great spiritual asset that transcends the limitations of time and geography, belonging to all of humanity. As Eastern wisdom states:

"All living things perish, but all non-living things—that is, science, artistic word, and art—live forever."

Navoi's legacy has been studied and cherished with respect by the Azerbaijani people since the 15th century as if it were the heritage of their own native master of the word. The lines of artistic and philosophical continuity between Navoi and Nizami, as well as Navoi and Fuzuli, demonstrate that the aesthetic thought of the Turkic-Islamic world was established upon unified humanist values. The 580th anniversary of Navoi and subsequent memorial events signify that the ideas of this great humanist thinker continue to maintain their relevance and light in our modern era.

Keywords: Alisher Navoi, Azerbaijani-Uzbek literary relations, Turkic language poetry, *"Char Divan"*, Muhammad Fuzuli, enlightenment, receptive aesthetics.

NİZAMİ GƏNCƏVİNİN "SİRLƏR XƏZİNƏSİ" POEMASINDA III SÖHBƏT: DÜNYANIN KEŞMƏKEŞİ VƏ BƏŞƏRİN İZTİRABLARI

NIZAMI GANJAVI'S POEM "THE TREASURY OF SECRETS," III CONVERSATION: THE WORLD'S TROUBLES AND THE SUFFERING OF MANKIND

Ulduz Fərhad Qəhrəmanova
PhD in Philology, Azerbaijan State Pedagogical University, Associate Professor at the Department of "Literature"
Orcid ID: 0009-0003-4418-9319

Abstrakt:

Nizami Gəncəvinin "Sirlər xəzinəsi" poemasında III söhbət: "Dünyanın keşməkeşi və bəşərin iztirabları" adlı məqalətdə şair insanlara nəsihət edir. İnsanlara həşəmət ağası (ərəb mənşəli söz olub mənası əzəmətli, böyük insan deməkdir) deyə xitabən müraciət edir. Şair həşəmət ağasına müraciət edərək deyir ki, cahana, dünyaya etinasız baxma! Çəkdiyin iztirab, əzab səni üzü ağ edər. İnsan ömrü boyu son gününü unutmamalıdır, həmişə axirətini xatırlamalıdır. Şair yoxsuluğu möhtəşəmlik ağası kimi xarakterizə etmişdir. Şairə görə, dünya vəfasız dünyadır. Bu torpaq çox canlar udubdur. Biz insanlar gəncliyimizi bu dünyaya vermişik. Bu dünyanın sanki doğmaca balasıyıq, amma dünya bizi qocaltdı.

Açar sözlər: Nizami Gəncəvi, "Sirlər xəzinəsi" poeması, İnsanoğlu, vəfasız dünya

Abstract:

Nizami Ganjavi's poem "The Treasury of Secrets" III conversation: In the article titled "The world's troubles and the suffering of mankind", the poet advises people. He addresses people as the master of majesty (an Arabic word meaning a great, majestic person). The poet addresses the master of majesty and says, "Do not look at the world and the world indifferently! The suffering and torment you endure will make you pale. A person should not forget his last day throughout his life, he should always remember the afterlife. The poet characterized poverty as the master of greatness. According to the poet, the world is an unfaithful world. This land has swallowed many souls. We, the people, have given our youth to this world. We are like the children of this world, but the world has aged us.

Keywords: Nizami Ganjavi, poem "The Treasury of Secrets", Human, unfaithful world

Nizami Gəncəvinin "Sirlər xəzinəsi" poemasında III söhbət: "Dünyanın keşməkeşi və bəşərin iztirabları" adlı məqalətdə şair insanlara nəsihət edir. İnsanlara həşəmət ağası (ərəb mənşəli söz olub mənası əzəmətli, böyük insan deməkdir) deyə xitabən müraciət edir. Şair həşəmət ağasına müraciət edərək deyir ki, cahana, dünyaya etinasız baxma! Çəkdiyin iztirab, əzab səni üzü ağ edər. İnsan ömrü boyu son gününü unutmamalıdır, həmişə axirətini xatırlamalıdır. Şair yoxsuluğu möhtəşəmlik ağası kimi xarakterizə etmişdir. Yoxsulluq mövzusuna gələndə onu Süleyman peyğəmbərlə müqayisə edərək demişdir:

Süleymanın mülkünü sorma hər an hardadır?

Mülkü öz yerindədir, de, Süleyman hardadır? (5; s.69)

Şairə görə, dünya vəfasız dünyadır. Bu torpaq çox canlar udubdur. Biz insanlar gəncliyimizi bu dünyaya vermişik. Bu dünyanın sanki doğmaca balasıyıq, amma dünya bizi qocaltdı:

Vermişik bu dünyaya biz öz gəncliyimizi,

Doğmaca balasıyıq, niyə qocaltdı bizi? (5, s.69)

Fələk İnsanoğlunu gah şah kimi ucaldar, gah da kuzəçiyə döndərər. İnsanoğlunun fəlakəti onun yaşadığı keşməkeşli həyatdır. İnsanın adəti hər yerdə narahatlılığıdır. İnsan vara-dövlətə uymamalıdır, təkəbbürdən uzaq gəzməlidir. Əyləncə ilə öz ömrünü başa vurmamalıdır. Əyləncənin həddini aşmamalıdır. İnsanoğlu bəxtiyar insanlara arxalanaraq ucala bilər. Şair təəssüf hissi ilə bildirir ki, dünyada yaxşı insan qalmayıbdır, şər-şərə yaraşıbdır:

Yaxşı insan qalmayıb, şər-şərə yaraşıbdır,

Dünya bal süfrəsidir, arılar daraşıbdır

...İnsanda nə ləyaqət, nə mərdlik qalıb indi,

İnsandan insanlığı namərdlik alıb indi (5, s.70)

Şairə görə, İnsanoğlu kimsənin kölgəsində səadətə çata bilməz. Əhdə dönük insanda sədaqət ola bilməz. Şair deyir:

Kimsənin kölgəsində səadət görməmişəm,

Əhdə dönük insanda sədaqət görməmişəm (5, s.70)

Hətta şair ədalətli Süleymanın çoxdan yoxa çıxdığına da təəssüflənir. Şair ədalətli Süleymanın yoxa çıxmasını belə əsaslandırır ki, dünyadakı haqsızlıqlara görə ədalətli Süleyman görünməz pəriyə dönübdür. Sonra şair insanlıqdan şikayətlənir və deyir ki, kimi ki, insan bilib dərdimi ona açdım, dərdimi açdığım insanın namərdliyi ilə qarşılaşdım.

Ədəbiyyat:

1. Azərbaycan ədəbiyyatı tarixi. 1 c. Bakı, Elm, 1960.

2. Azərbaycan ədəbiyyatı tarixi. VI cilddə, 2 c., Bakı, Elm, 2007

3. Ə.Səfərli, X.Yusifov. Qədim və orta əsrlər Azərbaycan ədəbiyyatı. Bakı, Maarif, 1982, 1998

4. Əlibəyzadə E. Azərbaycan ədəbiyyatı tarixi. Ən qədim dövrlər. Bakı, Qarabağ, 2009

Nizami Gəncəvi, “Sirlər xəzinəsi”, Çaşıoğlu, 2004, 167 s.

ƏMƏKDAŞLIĞA ƏSASLANAN TƏLIM MÜHITINDƏ TƏSVIRI SƏNƏT FƏNNININ ROLU

THE ROLE OF FINE ARTS EDUCATION IN A COLLABORATIVE LEARNING ENVIRONMENT

Emil Raul oğlu Ağayev
Teacher of the subject combination commission "Art and physical education" of the Azerbaijan State Pedagogical College under the Azerbaijan State Pedagogical University
ORCID NO: 0000-0001-9354-0839

Xülasə. Bu məqalədə orta məktəblərdə şagirdlərdə əməkdaşlıq bacarığının formalaşdırılmasında təsviri sənət fənninin rolu araşdırılmışdır. Müasir təhsil sistemində əməkdaşlıq, ünsiyyət, yaradıcı düşünmə və problem həlletmə bacarıqları əsas kompetensiyalar hesab olunur. Təsviri sənət fənni yaradıcı fəaliyyətə əsaslandığından şagirdlərin qrup şəklində işləməsi, birgə qərar qəbul etməsi, fikir mübadiləsi aparması və müxtəlif yaradıcı yanaşmaları müqayisə etməsi üçün əlverişli tədris mühiti yaradır. Məqalədə qeyd olunur ki, müəllim qrupları şagirdlərin bilik, bacarıq və fərdi xüsusiyyətlərini nəzərə almaqla formalaşdırmalı, hər bir şagirdin təlim prosesində fəal iştirakını təmin etməlidir. Bu yanaşma həm akademik nailiyyətlərin yüksəldilməsinə, həm də sosial bacarıqların inkişafına müsbət təsir göstərir. Tədqiqatda "Xalça naxışları" və "Bədii şüşə" mövzuları üzrə təşkil olunan yaradıcı tapşırıqlar nümunə kimi təqdim edilmiş, bu fəaliyyətlərin şagirdlərdə müstəqil düşünmə, estetik zövq, yaradıcı təxəyyül və komanda işi bacarıqlarını inkişaf etdirdiyi göstərilmişdir. Eyni zamanda, əməkdaşlığa əsaslanan öyrənmə prosesinin şagirdlərdə qarşılıqlı hörmət, məsuliyyət hissi, tolerantlıq və özünüifadə bacarıqlarını gücləndirdiyi vurğulanmışdır. Nəticə etibarilə müəyyən edilmişdir ki, təsviri sənət fənnində interaktiv və yaradıcı təlim metodlarının tətbiqi şagirdlərin şəxsiyyət kimi hərtərəfli inkişafına, sosial münasibətlərinin möhkəmlənməsinə və əməkdaşlıq mədəniyyətinin formalaşmasına mühüm töhfə verir.

Açar sözlər: təsviri sənət, əməkdaşlıq, orta məktəb, yaradıcılıq, interaktiv təlim.

Abstract . This article examines the role of fine arts education in developing students' cooperation skills in secondary schools. In contemporary education, collaboration, communication, creative thinking, and problem-solving are regarded as essential competencies for students. Since fine arts is a practice-based and creative subject, it provides an effective learning environment where students work collaboratively, exchange ideas, make joint decisions, and appreciate diverse artistic perspectives. The study emphasizes that teachers should organize heterogeneous groups by considering students' knowledge, abilities, and individual characteristics to ensure active participation and equal learning opportunities for all learners. Such an approach contributes not only to academic achievement but also to students' social and emotional development. The article presents classroom activities based on the topics "Carpet Patterns" and "Art Glass" as practical examples demonstrating how collaborative artistic tasks enhance independent thinking, aesthetic appreciation, imagination, and teamwork skills. Furthermore, collaborative learning encourages students to develop responsibility, mutual respect, tolerance, communication abilities, and self-expression while participating in group activities. The findings indicate that interactive and student-centered teaching methods in fine arts education create favorable conditions for strengthening cooperation among students and improving classroom interaction. Consequently, the integration of collaborative learning strategies into fine arts instruction supports the comprehensive development of students by fostering creativity, social competence, and positive interpersonal relationships. Therefore, fine

arts should be regarded not only as a means of artistic expression but also as an effective educational tool for cultivating collaboration skills and preparing students for successful participation in modern society.

Keywords: fine arts, collaboration, secondary school, creativity, interactive learning.

Təsviri sənət fənni şagirdlərdə yaradıcı qabiliyyətləri inkişaf etdirməklə bərabər, əməkdaşlıq və birgə çalışmaq qabiliyyətlərini də ön plana çıxarır. Yəni təqdim edilən yaradıcı tapşırıqda fərqli düşüncələrə malik olan şagirdlər rəng, obraz, süjet həllini fərqli seçə bilərlər. Şagirdlər burada birgə öyrənmə, rəqabətyönümlü öyrənmə bacarıqlarını da inkişaf etdirirlər.

Orta məktəblərdə təsviri sənət fənninin tədrisində "interaktiv-yaradıcı öyrənmə mühiti"nə imkan yaratmaq lazımdır. Sinifdə qruplarla işləyərkən əməkdaşlıq etməyin qaydaları və qrupla çalışmağın prosedurları şagirdlərə izah edilməlidir.

Təsviri sənət müəllimi qrupların tərkibini qarışıq formada yəni, bilik, bacarıq və düşüncələri fərqli olan şagirdləri əməkdaşlığa cəlb etməlidir. Təlim nəticələri aşağı olan şagirdlər diqqətdən kənar qalanda şagirdlər arasında nailiyyət fərqi daha da artır və ruh düşkünlüyü baş verir.

V sinif şagirdlərinə "Xalça naxışları" mövzusunda ornamentləri təsvir etmək tapşırılmalı və stilizə bacarıqlarından fərqli istifadə etmək onlarda müstəqil düşüncə formasının inkişafına gətirib çıxaracaqdır. Şagirdlər xalçaları bəzəyən təbiət və həndəsi motivləri təsvir edirlər və fərqli formada yanaşırlar.



İLL 1. Xalça naxışları

Təsviri sənət fənninin məzmun xətlərinə uyğun bilik və bacarıqlar dəyərləndirilməlidir. Şagirdlər öyrənəcəkləri məzmundan əlavə birgə qərarların verilməsi, qarşılıqlı fikir mübadiləsi kimi məqamları da özləri üzərində inkişaf etdirirlər. Əməkdaşlıq zamanı şagirdlər rəqabətə əsaslanan qrup işinin tapşırığını müəyyən bir vaxt ərzində dəqiq, səlis yerinə yetirməlidirlər. Bu zaman yerinə yetirilən tapşırığın məqsədi aydın olmalıdır. Müəllim "Bədii şüşə" mövzusunu keçərkən təqdim edilən şüşə formaları üzərində müəyyən rəng, süjet tapşırıqlarını verməklə, həm də şagirdlərdə yaradıcı düşüncəni inkişaf etdirir.

İLL 2. Bədii şüşə

Təsviri sənət fənninin tədrisi metodikasında şagirdlərdə əməkdaşlıq bacarığının formalaşması aşağıdakı müsbət inkişaflara gətirib çıxarır:

-şagird və müəllim arasında yaradıcı fikirlərin mübadiləsi;

-müxtəlif yaradıcı düşüncələrin inkişafı və qarşılıqlı mübadiləsi;

-özünüifadəetmə, incəsənət əsərini təhliletmə və fərqli yanaşma bacarıqları inkişaf etdirilir.

ƏDƏBİYYAT SİYAHISI:

1.Metodika və pedaqogika. Bakı 2024. s 66-67.

2.V sinif təsviri incəsənət dərsliyi. Bakı 2020. 80 s.

ABOUT SOME VERBS USED IN AZERBAIJAN DIALECTS AND ACCENTS

Asgarova Ganira Ali gizi
Teacher
Azerbaijan State Pedagogical College under Azerbaijan State Pedagogical University, Azerbaijan

Abstract: The Second Karabakh War, which began in 2020, ended with the glorious victory of the Azerbaijani army. The main part of our occupied territories was liberated. In 2023, as a result of the battles within only twenty-four hours, the sovereignty of our state over all our regions was ensured. It goes without saying that various aspects of Karabakh and Western Azerbaijani dialects and dialects have been studied by different authors in all periods. Today, this problem is more urgent. From this point of view, our article talks about the study of simple verbs used in the dialects and dialects of the Azerbaijani language. It should be noted that the richness of simple verbs used in Azerbaijani dialects and dialects attracts attention. It goes without saying that the verbs used in the territory of Karabakh and West Azerbaijan, which is the ancient land of Azerbaijan, share commonalities with other dialects and dialects of our language. Their study in the context of linguistics allows obtaining a number of interesting facts. Thus, the words used in these dialects and dialects prove that those lexical units are closely related to the occupation, life and everyday life of the population of the region. In the article, a number of lexical units collected from Azerbaijani dialects and dialects were analyzed in a comparative way with the studied words of various dialects and dialects of our language, examples in ancient written sources and monuments. As a result of research, the author has obtained a number of interesting facts. It is known that the role played by dialects, folklore, ethnography in the formation and propagation of the spiritual culture, outlook, and morality of any nation is irreplaceable. To get detailed information about a people's historical background, traditions, behavior and habits, interactions and influences with related peoples and their culture, it is very important to refer to dialect and dialect words. The Azerbaijani people with a rich historical and cultural past have their own dialect and dialect words that can be distinguished among all Turkic languages. Verbs are undoubtedly the main shield and protector of the purity of our language. All the important spiritual and moral qualities of the area and the people where it was born have somehow survived the test of centuries by being absorbed into its dialects and dialects. That is why the "Kitabi-Dade Gorgud" epic, which is considered one of our oldest literary examples, has words that are currently our literary language. has become archaic and lost its generality. However, we witness their use of our dialects and dialects. Regardless of the period, time, and place of its origin, Azerbaijani dialects and dialects have always been connected with our national way of thinking. Based on the conducted research, we can say that the dialects and dialects of the Azerbaijani language are very rich in their own characteristics. Thus, dialect and dialect words are among the most important examples of the national and moral values of our people. Their study is very important for analyzing the historical background of our nation, relations with neighboring nations, and the antiquity of our traditions. A number of interesting scientific results were obtained during the research.

Keywords: Azerbaijan, dialect and accent, verb, lexical unit, linguistic analysis

Özət: 2020-ci ildə başlanan İkinci Qarabağ müharibəsi Azərbaycan ordusunun şanlı zəfəri ilə yekunlaşdı. İşğal olunan ərazilərimizin əsas hissə azad edildi. 2023-cü ildə isə cəmi iyirmi dörd saat ərzindəki döyüşlərin nəticəsində dövlətimizin bütün topaqlarımız üzərindəki suverenliyi təmin edildi. Söz yox ki, bütün dövrlərdə istəq Qarabağ, istərsə də Qərbi Azərbaycan dialekt və şivələrinin müxtəlif aspektləri ayrı-ayrı müəllidlər tərəfindən tədqiq edilmişdir. Bu gün isə həmin problem daha aktualdır. Bu mövqedən çıxış edərək məqaləmizdə Azərbaycan dilinin dialekt və şivələrində işlənən sadə feillərin öyrənilməsindən bəhs olunub. Qeyd etməliyik ki, Azərbaycan dialekt və şivələrində işlənən sadə feillər zənginliyi ilə diqqəti cəlb edir. Söz yox ki, qədim Azərbaycan torpağı olan Qarabağ və Qərbi Azərbaycan ərazisində də işlənən feillər dilimizin digər dialekt və şivələri ilə ortaqlıq təşkil edir. Onların dilçilik kontekstində tədqiqi bir sıra maraqlı faktların əldə edilməsinə imkan verir. Belə ki, bu dialekt və şivələrdə işlənən sözlər sübut edir ki, həmin leksik vahidlər bölgə əhalisinin məşğuliyyəti, həyatı və məişəti ilə sıx bağlıdır. Məqalədə Azərbaycan dialekt və şivələrindən toplanan bir sıra leksik vahidlər dilimizin tədqiq edilmiş müxtəlif dialekt və şivə sözləri, qədim yazılı mənbə və abidələrdəki nümunələrlə müqayisəli şəkildə təhlil olunmuşdur. Araşdırmalar nəticəsində müəllif bir sıra maraqlı faktlar əldə etmişdir. Məlumdur ki, istənilən bir xalqın mənəvi mədəniyyətinin, dünyagörüşünün, əxlaqının formalaşması və təbliğində dialektlərin, folkorun, etnoqrafiyanın oynadığı rol əvəzolunmazdır. Bir xalqın tarixi keçmişi, adət-ənənələri, davranış və vərdişləri, qohum xalqlarla, onların mədəniyyəti ilə qarşılıqlı əlaqə və təsirləri haqqında ətraflı məlumat almaq üçün dialekt və şivə sözlərinə müraciət olduqca əhəmiyyətlidir. Zəngin tarixi mədəni keçmişi olan Azərbaycan xalqının özünəməxsus və bütün türk dilləri içərisində seçilə bilən dialekt və şivə sözləri mövcuddur. Feillər isə heç şübhəsiz dilimizin saflığının əsas qalxanı, qoruyucusudur. Yarandığı ərazi və xalqın bütün mühüm mənəvi, əxlaqi keyfiyyətləri bir növ onun dialekt və şivələrinə hopmaqla əsrlərin sınağından çıxaraq günümüzə gəlib çatmaqdadır Odur ki, ən qədim ədəbi nümunələrimizdən sayılan "Kitabi-Dədə Qorqud" dastanın da qarşımıza çıxan elə sözlər var ki, hazırda onlar ədəbi dilimiz üçün arxaikləşmiş və ümumişləkliyini itirmişdir. Lakin dialekt və şivələrimizin onların işlədildiyinin şahidi oluruq. Yarandığı dövr, zaman, məkandan asılı olmayaraq Azərbaycan dialekt və şivələri milli düşüncə tərzimizlə daim bağlı olmuşdur. Aparılan tədqiqatlara əsaslanaraq deyə bilərik ki, Azərbaycan dilinə məxsus dialekt və şivələrinin özünəməxsus xüsusiyyətləri çox zəngindir. Belə ki, dialekt və şivə sözləri xalqımızın milli-mənəvi dəyərlərinin ən mühüm nümunələrindəndir. Onların öyrənilməsi xalqımızın tarixi keçmişi, qonşu xalqlarla əlaqələri, adət-ənənələrimiz qədimliyini təhlil etmək üçün olduqca əhəmiyyətlidir. Tədqiqat zamanı bir sıra maraqlı elmi nəticələr əldə edilmişdir.

Açar sözlər: Azərbaycan, dialekt və şivə, feil, leksik vahid, linqvistik təhlil

Giriş. Azərbaycan dialekt və şivələrinin öyrənilməsi dilçiliyin aktual problemlərindəndir. Dialekt və şivələrin lüğət tərkibinin tədqiqi xalqımızın tarixi, adət və ənənələri ilə bağlı dil faktlarının aşkar olunmasında xüsusi əhəmiyyət daşıyır. Günümüzədək dilimizin dialekt və şivələrinin mühüm bir hissəsi tədqiq edilmişdir. O cümlədən Qəbi Azərbaycan şivələri ilə bağlı filologiya elmləri doktoru, professor İbrahim Bayramovun "Qərbi Azərbaycan şivələrinin leksikası", "Aazərbaycan dilinin Meğri şivələri", "Qərbi Azərbaycan

şivələri lüğəti" və s. nəşr etdirilmişdir. Ümumi qrammatik əlamət dedikdə sözün ifadə etdiyi məna, daşıdığı sual və cümlədə mövqeyi nəzərdə tutulur. Sözün ifadə etdiyi məna onun yalnız mənasını əhatə edir. Hər bir sözün müəyyən nitq hissəsinə aid olması üçün həmin söz leksik məna ilə yanaşı, qrammatik məna da kəsb etməlidir. Qrammatik məna isə sözün öz daxilində olmur. Qrammatik formalarla meydana çıxır.

Əsas təhlil. Sözlər nitq hissələrinə bölünmək üçün dilin qrammatik quruluşuna xas olan qrammatik kateqoriyalara malik olmalıdır. Beləliklə, leksik mənaya malik olan sözlər qrammatik məna kəsb edərək qrammatik əlamətlər qəbul edir, digər sözlərlə əlaqələnərək fikrin ifadə olunmasına təminat verir. Nəticədə cümlələr formalaşır və sözlər cümlədə müəyyən sintaktik vəzifə daşıyır. Qrammatik məna ilk növbədə cümlədə başqa sözlərlə əlaqə və münasibətlərdə müəyyənləşən mənadır. Qrammatik mənanın həm morfoloji həm də sintaktik ifadə vasitələri vardır. Qrammatik mənanın morfoloji ifadə vasitəsinə sözlərin dəyişməsi (şəkilçilər və vurğular), sintaktik ifadə vasitələrinə isə sözlərin bir- birinə yanaşması intonasiya və sözlərin sırası daxildir. Azərbaycan dilindəki nitq hissələrini müstəqil mənalarına, qrammatik formalarına (morfoloji əlamətlərinə) və sintaktik roluna görə əsas, köməkçi və xüsusi nitq hissələri olmaqla üç qrupa ayırmaq olar. Hər şeydən əvvəl, sözləri lüğəvi mənaya malik olan və lüğəvi mənası olmayan sözlər adı ilə iki qrupa bölmək olar. Dilimizin lüğət tərkibindəki sözlərin 99% dən çoxunun lüğəvi mənası var. Lüğəvi mənası olmayan sözlər köməkçi vasitələrdir. Əslində köməkçi vasitələr olmasa da fikir ifadə etmək mümkün olur. Lakin fikrin bütün incəliyinə qədər ifadə olunmasında bu vasitələrin rolu danılmazdır. Vaxtilə dildə belə köməkçi vasitələr olmamış, onlar dilin inkişafı prosesində kütləvi mənaya malik olan sözlərdən əmələ gəlmişdir (10).

Müasir Azərbaycan dilçiliyində nitq hissələri əsas və köməkçi olmaqla iki qrupa ayrılır. Əsas nitq hissələri leksik (lüğəvi mənaya malik olur, müvafiq suala cavab verir, müəyyən sintaktik vəzifə daşıyır, söz yaradıcılığı prosesində fəal iştirak edir. Onlar əşyanın adını, əlamət və keyfiyyətini, miqdarını, hal-hərəkətini, hərəkətin əlamətini bildirir. Əsas nitq hissələrinə isim, sifət, say, əvəzlik, feil və zərf daxildir.

Azərbaycan dialekt və şivələri ədəbi dildən fərqlənən bir sıra morfoloji xüsusiyyətlərə malikdir. Əsas nitq hissələri hər birinə aid maraqlı məqamlarla qarşılaşsaqda, tədqiqatda daha çox isim, sifət və feillərin bəzi məsələləri araşdırılmışdır.

Feil Azərbaycan dialekt və şivələri, həmçinin Qərbi Azərbaycan şivələrində səciyyəvi xüsusiyyətləri ilə diqqət çəkən əsas nitq hissələrindən biridir. Burada nəinki ədəbi dildə, hətta başqa dialekt və şivələrimizdə işlənməyən bəzi sözlərə rast gəlmək mümkündür. Dialekt və şivələrimizdə feillər sadə, düzəltmə və mürəkkəb quruluşludur.

Burada işlənən düzəltmə feillərin çoxuna ədəbi dilimiz və digər dialekt və şivələrimizdə də rast gəlirik. Dilimizin dialekt və şivələrində morfoloji yolla, müxtəlif nitq hissələrinə leksik şəkilçilər artırılmaqla sözdüzəltmə ən çox yayılmış üsullardan biridir. Bu isə başa düşüləndir. Ona görə ki, iltisaqi dilllərdə morfoloji yolla sözdüzəltmə sintaktik yolla sözdüzəltməyə nisbətən daha güclü inkişaf etmişdir. Dialekt və şivələrimizdə morfoloji yolla əmələ gələn xeyli sayda feillər mövcuddur ki, onları da asanlıqla kök və şəkilçiyə ayırmaq mümkündür. Məsələn, teləməx`/teliməx` – 1. Anası ölmüş bir quzu və ya çəpici başqa bir heyvana əmizdirmək. *Məs.: Bı quzu:n anası yoxdu, gərəx` unu ağbaş qoyna teliyəm*. Tel leksik vahidi Azərbaycan ədəbi dili və dialekt və şivələri üçün xarakterik leksik vahid olmasa da, dilimizdə işlənən bir sıra dialekt leksik vahidlərinin formalaşmasında iştirak etmişdir. Aparılmış bir sıra tədqiqatlarda,

xüsusilə Radlovun əsərlərində bu sözün ırğız dilində anasından ayrılmış və başqa heyvanı əmən cavan heyvan mənasında işlədildiyi qeyd edilmişdir. Maraqlıdır ki, bu fakt hazırda Meğri və Culfa şivələrində də öz varlığını qorumaqdadır. Söz *teləməx`, teləx`çi, teliyən* tipli sözlərin kökündə öz varlığını qorumaqdadır.

Ədəbi dilimizdə olduğu kimi, dialekt və şivələrimizdə düzəltmə feillər əmələ gəldiyi kökdən asılı olaraq iki yerə bölünür:

1. Adlardan düzələn düzəltmə feillər 2. Feillərdən düzələn düzəltmə feillər

1. Adlardan düzələn düzəltmə feillər

Dilimizin dialekt və şivələrində düzəltmə feillər adlardan həm məhsuldar, həm də qeyri-məhsuldar şəkioçilər vasitəsi ilə əmələ gəlir. Tədqiq olunan Culfa şivələrində isim və sifətlərdən yaranan düzəltmə feillər say, əvəzlik və zərflərdən düzələnlərə müqayisədə üstünlük təşkil edir.

-la,-lə şəkilçi ədəbi dilimizdə olduğu kimi dialekt və şivələrimizdə düzəltmə feillərin formalaşmasında daha fəal çıxış edən şəkilçilərimizdəndir. Dilimizin dialekt və şivələrində bunu bir sıra düzəltmə feillərin əmələ gəlməsində müşahidə edirik. Söz yox ki, ədəbi dilimizdə fərqlə olaraq şivələrimizdə onun -da,-də, -na, -nə, -za, -zə variantları ilə qarşılaşırıq. Qərb şivələrimizdə isə daha çox la, lə və da, də variantı ilə işlədildiyinin şahidi oluruq. Məs.: boğazlamax/boğazdamax sözü tədqiq etdiyimiz Azərbaycan şivələrində "kəsmək, tutmaq, yaxalamaq" mənalarında işlənir. 1. Böyünnərisi bi qurbannığım varıydı unu boğazdadım (yəni kəsdim). 2. Qan vırdı beynə ıhıda bını boğazdadım (yəni boğazından tutdum). Göründüyü kimi düzəltmə feil boğaz ismi+la/da adlardan feil düzəldən leksik şəkilçinin köməyi ilə yaranmışdır.

Bir başqa leksik vahidə baxaq. Müasir şivələrimizdə "örtmək, basdırmaq" mənasında işlənən çullamax feilinə rast gəlirik. Məs.: Uşağa yaman soyux dəyip axşam bankalıyıp üsdünü çulladım ki bəlkəm ayılsın bir az. Bildiyimiz kimi dilimizin digər dialekt və şivələrində çul sözü həm yəhərin altından salınan parça, həm də örtük, döşənəcək mənalarında mövcuddur. Təsadüfi deyil ki, hazırda dilimizin Culfa şivələrində "döşənəcək" mənasında işlədilən çul-çapıt mürəkkəb isminə rast gəlinir. Ona görə də çullamax feili çul ismi və +la feil düzəldən şəkilçinin iştirakı ilə formalaşmışdır.

Qərbi Azərbaycan şivələrində olduğu kimi dilimizin digər dialekt və şivələrində işlənən xışma+la, badağ+la, quyrux+la, tapan+na və s. (11., s 46) kimi düzəltmə feilləri də məhz -la, -lə şəkilçisi və onun fonetik variantlarının köməyi ilə əmələ gəlmişdir. Yuxarıda qeyd etdiyimiz xışmala və tapanna leksik vahidləri Azərbaycan şivələrində də geniş işlənmə arealına malik düzəltmə feillərdəndir. Xışma sözü bəhs etdiyimiz şivələrdə koma, bir ovuc anlamlarında, tapan sözü isə şumdakı iri kəsəkləri əzmək üçün alət, vasitə mənalarında şivələrimizdə mövcuddur.

Müasir şivələrimizdə xırxalamax, adaxlamax, beçələx`, quzulamax, yelinnəməx`, irtməx`ləməx`, törpüləməx`, dümləməx`, qəzilləməx`, çaylamax` və s. kimi xeyli sayda isimdən düzəltmə feilər işlənir ki, onların da bir qisminə həm dilimizin müxtəlif dialekt və şivələrində, həm də ayrı-ayrı türk dillərinin şivələrində rast gəlmək mümkündür. Onların bəzilərinin təhlilinə diqqət yetirək.

Azərbaycan şivələrində beçələməx` sözü "artmaq, çoxalmaq, dən tutmaq" mənalarında omonim leksik vahid kimi geniş işlənmə arealına malikdir. Bunu aşağıdakı nümunələrdə əyani şəkildə müçahidə edə bilərik:

1. Bu gün arılarım beçəliyib.
2. Buğda day beçələdi balam.
3. Ağac dibinnən beçəliyib həyəti alıb ağzına və s.

Beçə sözü bəhs etdiyimiz şivələrdə yeni yaranan, əmələ gələn, cavan, həmçinin cavan xoruz vəs. mənalarını ifadə edir. Məs.: Cücələrimizin içində üş-dört dənə beçə (cavan xoruz) var. Birinci nümunədə arının yaz aylarında artıb çoxalması, yəni cavan arıların əmələ gəlməsi beçələmək sözü ilə ifadə olunur. Sünbülün yenicə dən tutması və ağacın yanında cavan şivlərin əmələ gəlməsi də elə sözün ilkin mənası ilə yaxından bağlıdır. Bütün bu faktlara əsaslanaraq deyə bilərik ki, beçələməx` düzəltmə feili beçə ismi və lə isimdən feil düzəldən leksik şəkilçisinin iştirakı ilə yaranmışdır. Qeyd edək ki, bu söz dilimizin əksər dialekt və şivələrində daha çox arının artımı və ağacın dibində cavan şivlərin əmələ gəlməsi mənasında işlədilməkdədir.

Dilimizin şivələrində işlənən maraqlı düzəltmə feillərdən biri də adaxla feilidir. Bu leksik vahidin dilimizin müxtəlif dialekt və şivələrində türk dilləri və onların şivələrində fərqli məna çalarları diqqəti cəlb edir. Belə ki, hazırda şivələrimizdə işlənən adaxla feili daha çox "nişanlamaq, yəni ailə qurucaq iki gənc arasında ilkin bağı yaratmaq" məzmununu ifadə edir. *Məs.: Böyünnərsi qonşu oğlun da adaxladılar.*

Azərbaycan şivələrində feilin bu məna çaları daha geniş işlənmə arealına malik olsa da, dilimizin Daşkəsən, Fizuli şivələrində daha çox ilk addım atmaq, ilk yerişi yerimək mənalarını ifadə edir (2). *Məs.: Nəvəm yavaş-yavaş adaxley.* Düzəltmə feil adax –ayaq, addım ismi və -la leksik şəkilçisinin köməyi ilə formalaşmışdır. Adax sözü qədim türkmənşəli leksik vahidlərimizdən biridir. Ona görə də bir sıra türk dillərində adax "qurban", adax vermək "qurban kəsmək, qurban vermək" mənasında işlənir. Düşünürük ki, bu mənasında adax sözü özü məhz ada+x quruluşuna malikdir. Buradakı ada sözü qədim türk dilində "fəda etmə, həsr etmə, qurban vermə" kimi mənaları ifadə etmiş, həmin kökə x sözdüzəldici şəkilçisi artırılmaqla yeni mənalı adax qurban ismi yaranmışdır. Təsadüfi deyil ki, müasir türk dilində kendini adamak –özünü həsr etmək, qurban vermək mənalı feil hazırda işlənməkdədir. O ki qaldı adax+la düzəltmə feilinin kökünə isə bizim də fikrimizcə, bu məhz ayaq/adax "ilk addım, ilk mərhələ" kök morfemindən törəmişdir. Bəhs etdiyimiz şivələrdə adaxlı-nişanlı, adaxlıbazlıx və s. kimi leksemlərin də işlədildiyi məlumdur.

Müasir Azərbaycan dialekt və şivələrində lüllə, silələ, saxla, cırıxla, düşgünnə və s. kimi sifət kökündən törəyən düzəltmə feillərə də rast gəlinir. Bu leksik vahidlər içərisində maraq doğuranlardan biri saxlamax düzəltmə feilidir. Qeyd edək ki, müasir ədəbi dilimizdə sadə feil kimi qəbul edilən və daha çox "mühafizə etmək, qorumaq" mənasını ifadə edən bu leksik vahid Azərbaycan şivələrində ədəbi dilimizdəki mənası ilə yanaşı, "gizlətmək" mənasını da ifadə edir. *Məs.: Sirri: mənnən saxlama, nə dərdin var de bilim.*

Tədqiqatçılar arasında ümumi fikir bundan ibarətdir ki, düzəltmə feil qədim türk dillərinə məxsus sax/saq – ayıq, sayıq sifəti və -la leksik şəkilçisinin iştirakı ilə yaranmışdır. Müasir tatar, qazax dillərində söz ilkin etimologiyasını qoruyub saxlamaqda və işlədilməkdədir. Hazırda Türkiyə türkcəsi və şivələrində də saklı gizli sifəti işlədilir.

Bəhs olunan şivələrdə maraq doğuran düzəltmə feillərdən biri də silələməx` ağzına qədər doldurmaq feilidir. *Məs.: Qabı illap silələmisən ki, a bala.* Qeyd edək ki, silə sifəti dilimizin dialekt və şivələrində "dolu, tam" mənasını ifadə edir.

Tədqiq olunan Azərbaycan dialekt və şivələrində diqqəti cəlb edən leksemlırdən biri də məhz lülləməx` düzəltmə feilidir. Bu söz şivələrimizdə ağacı, taxtanı yonub boru şəklinə salmaq mənasını ifadə edir. Sözün kökü "lül" düz sifətidir. *Məs.: Bizim Əhmət yamancanax dülgərdi taxtanı bi lüllüyür kin gəl görəsən.* Maraqlıdır ki, dilimizin bir sıra şivələrində lül komponentli digər ifadələrə də rast gəlinir. Belə ki, bəhs etdiyimiz şivələrdə "həddindən artıq içki içib sərxoş olmaq" anlamında lül atəş, xüsusilə pişik, it kimi heyvanların nədənsə qorxan zaman quyruğunun dik dayanmasına "lül durmaq" və s. tipli söz və ifadələrlə qarşılaşırıq.

Müasir Azərbaycan şivələrində qarşımızıa çıxan və geniş işlənmə arealına malik olan maraqlı düzəltmə feillərdən biri də düşgünnəməx` feilidir. Hər hansı bir insanın və ya digər canlı varlıqlığın sürətli zəifləməsi tədqiq etdiyimiz şivələrdə bu leksik vahidlə ifadə olunur. Yəni düşgünnənəx` feili düşgün sifətindən yaranmışdır. Qeyd edək ki, həmin söz dilimizin Naxçıvan dialekt və şivələri üçün xarakterik olub sözügedən şivələrin əksəriyyətində özünü göstərir. *Məs.: Day mən də qocalmışam. Hər gün bir az düşgünnüyrəm.*

2. Feillərdən düzələn düzəltmə feillər

Ədəbi dilimkizdə olduğu kimi dialekt və şivələrimizdə də düzəltmə feillərin bir qismi feil köklərindən yaranır. Buradakı şəkilçilərin bəziləri feil yaradıcılığında fəal iştirak etsə də, digərləri qeyri-məhsuldarlığı ilə diqqəti cəlb edir. Hazırda Azərbaycan şivələrində feil köklərindən yaranan düzəltmə feillər əsasən bunlardır. Teziməx`, arıtdamax, şüşdəməx`, qavalamax, çözdəməx` və s. Nümunələrdən də göründüyü kimi, şivələrimizdə feil köklərindən yarana leksik vahidlər ad bildirən nitq hissələrindən yaranan sözlərlə müqayisədə azlıq təşkil edir. Feil köklərindən yaranmış leksik vahidlərin bəzilərinin təhlilinə baxaq.

Müasir Azərbaycan dialekt və şivələrində qarşımıza çıxan maraqlı düzəltmə feillərdən biri *teziməx`-qaçmaq* feilidir. *Məs.: Yazıx qorxusunnan tezidi.* Bu feil şivələrimizdə həm insanlar həm də digər canlılar üçün eyni leksik anlamda istifadə edilir. Belə ki, aparılan araşdırmalar zamanı bəhs etdiyimiz şivədə təzix′ qoun (T., Sal.) –tez-tez sürüdən ayrılıb qaçan, itən heyvan mənasında işlənir. *Məs.: Təzix′ qoyunu tutmağ çətin o:r, gərəx′ kürnəşə salasan (T.).*

Birinci növ ismi birləşmə şəklində olan bu terminin birinci komponenti olan "təzix′" sözü türk mənşəli qədim leksem olan təzimək – qaçmaq feilindən yaranmışdır. Bu feil qədim türk dillərində mövcud olmuşdur və hazırda dialekt və şivələrimizdə yaşamaqdadır.

Belə ki, tezi – qaç feili "Divani-lüğət-it türk"də tezmek (9, s. 611) – qaçmaq, Orxon-Yenisey abidələrində tez (13, s.174), qədim Göytürk və Uyğur yazılı abidələrində tezmek (8, s.100), Türkmən dilində tezmek (12, s.198), qədim Özbək dilinin XIV əsr yazılı abidəsi olan "Tarjuman" abidəsində tizik (14, s.114), Türkiyə türkcəsinin Çoruh şivəsində tezmey (6, s.280) şəklində, eyni mənalarda işlənməkdədir. Azərbaycan dilinin Meğri (Mehri), Kəlbəcər dialekt və şivələrində təzməx′ (2, s. 564) şəklində eyni mənada işlək dialekt sözüdür.

Qeyd edtmək lazımdır ki, Azərbaycan şivələrində düzəltmə feillər içərisində sinonimlik təşkil edən bir sıra leksik vahidlərə də rast gəlinir. Apardığımız araşdırmalar zamanı qarşımıza çıxan və maraq doğuran *tüyümək* feili də bu qəbildən olan leksemlərimizdən biridir. Bir faktı

unutmayaq ki, tezimək feilindən fərqli olaraq tüyümək feili daha çox insanlar üçün istifadə edilən xarakterik sözlərimizdəndir. *Məs.: Bi də baxdım kin tüyüdü.*

Tüyümək feili daha çox "gizlicə, oğrun, cəld qaçmaq" anlamını ifadə edir. Dilimizin Meğri şivələrində bu feil kökündən törəyən tüynex`-cəld, zirək (3, 587) leksimi ilə qarşılaşırıq.

Dilimizin Qazax, Borçalı şivələrində eyni leksem dazılamax fonetik variantında işlədilir. *Məs.: Quşa heş çatmamışdım ki, dazıladı (Borçalı)* (1).

Dilimizin dialekt və şivələrində qarşımıza çıxan maraqlı leksik vahidlərdən biri də şüşdəməx` feilidir. *Məs.: Qırışıxları bir ikidə şüşdədim.* Bu düzəltmə feil qədi türk dillərində açmaq mənasında işlədilən şüş/şeş feil kökü və -lə sözdüzəldici şəkilçisinin fonetik variantı olan -də şəkilçisinin iştirakı ilə yaranmışdır. Şüş/şeş feili "Kitabi-Dədə Qorqud" dastanında da işlənmişdir. "... Dirsə xan aydır: - Mənim əlimi şeşin, qolça qopuzumu əlimə verin.." (7). Şeş sözü türk dillərinə aid lüğətlərdə də açmaq mənasında verilmişdir.

Tədqiq etdiyimiz şivələrdə işlədən maraqlı düzəltmə feillər sırasında çözələməx` - açmaq feilini də qeyd edə bilərik. *Məs.: İpəx` sapı illap bərbat çözələmisən e.* Müasir Azərbaycan ədəbi dili üçün arxaikləşən bu söz, dialekt və şivələrimizdə öz varlığını qoruyub saxlaya bilmişdir. Maraqlıdır ki, hazırda Türkiyə türkcəsində çözmək feili həll etmək, açmaq mənasında öz ilkin leksin məzmunu qoruyub saxlamaqdadır. *Məs.: Düğümü çözmüş bulunmaktayız.*

Nəticə. Yekun olaraq qeyd etmək lazımdır ki, müasir Azərbaycan dialekt və şivələrində, o cümlədən də Qərbi Azərbaycan şivələrində feillər öz zənginliyi ilə diqqəti cəlb edir. Onların hamısı haqqında bir tədqiqatda bəhs etmək bir qədər çətindir. Araşdırılan sözlərin böyük əksəriyyəti milli mənşəli olmaqla zəngin ümumtürk leksikasının mühüm laylarından birini təşkil edir. Belə sözlərin müasir dilçilik aspektindən etnolinqvistik təhlilinin aparılması bir sıra maraqlı faktların orataya çıxarılmasına imkan verir. Həmin sözlərin milli aspektdən, ümumtürk mədəniyyəti, dilçiliyi kontekstində öyrənilməsi, onların xalqımızla, bu torpaqla bağlılığının sübut olunması, günümüzdə ərazilərimizə qarşı irəli sürülən əsassız erməni iddialarına tutarlı cavab verə biləcək əhəmiyyətli bir mənbə olduğunu göstərməkdədir. Aparılan araşdırma və təhlillər sübut edir ki, öyrənilən sözlər, şübhəsiz, regionla bağlıdır və xalqımızın milli-mənəvi mədəniyyətinin, dialekt leksikasının mühüm tərkib hissəsini təşkil edir. Belə sözlər dilimizin qədim tarixi köklərini özündə yaşadan, bizə və bizdən sonrakı nəsillərə miras qoyan olduqca dəyərli mənbədir. Onların araşdırılaraq ortaya çıxarılması, linqvistik təhlil olunması və bundan sonrakı nəsilləri ötürülməsi müasir dil tarixinin, tarixi dialektologiyanın qarşısında dayanan aktual məsələlərdəndir.

ƏDƏBİYYAT

1. Azərbaycan dilinin Dərbənd dialekti. Bakı: Elm, 2009, 448 s.
2. Azərbaycan dilinin dialektoloji lüğəti. 2 cilddə, I c. Ankara: Türk Dil Qurumu, 1999, 374 s.
3. Azərbaycan dilinin dialektoloji lüğəti. İki cilddə, II cild. Ankara: Türk Dil Qurumu, 2003, 279 s.
4. Azərbaycan dilinin Naxçıvan qrupu dialekt və şivələri. Bakı: Azərb. SSR EA nəşriyyatı, 1962, 326 s.
5. Bayramov İ. Qərbi Azərbaycan şivələrinin leksikası. Bakı: Elm və təhsil, 2011, 440 s.

6. Bədəlova A. Qərbi Azərbaycan şivələri lüğəti. Ankara, 2023, 314 s.
7. Caferoğlu A. Doğu illerimiz ağızlarından toplamalar, Ankara: Türk Dil Kurumu, 1995, 296 s
8. Ergin M.Dede Korkut Kitabı. Ankara: Türk Dil Kurumu, 1994, 483 s.
9. Xudiyev N.M. Quliyev Ə.A. Qədim türk abidələrinin sözlüyü. Bakı: ADPU, 1992, 127 s.
10. Kaşğarlı M. "Divani lüğət-it-türk". Ankara: Türk Dil Kurumu, IV c. 2006, 885 s.
11. Müasir Azərbaycan dili. Üç cilddə, II cild. Morfologiya. Bakı: Elm, 1980, 510 s.
12. Rüstəmov R. Azərbaycan dili dialekt və şivələrində feil.Bakı: Maarif, 1965, 320 s.
13. Taçmıradov T., Çarıyev M. Türkmən ədəbi dilinin orfoepiyası və qısa orfoepik sözlüyü, Aşqabad, Elm, 1967, 243 s.
14. Tekin T. Orhon yazıtları. Ankara: Türk Dil Kurumu, 2010, 201 s.
15. Юнисов А. Узбек теленен "Таржуман" ел язмалары. Узбекистан, Ташкент, Фанн неширяты, 1980, 127 s.

ÇOCUKLARDA VE ERGENLERDE YALNIZLIK SALGINI (LONELİNESS EPİDEMİC): SAĞLIK SONUÇLARI VE HEMŞİRELİK YAKLAŞIMLARI

THE LONELINESS EPIDEMIC IN CHILDREN AND ADOLESCENTS: HEALTH OUTCOMES AND NURSING APPROACHES

Assoc. Prof. Dr. Hakan AVAN [1]
Afşin School of Health, Kahramanmaraş Sütçü İmam University, Kahramanmaraş, TURKEY
ORCID ID: 0000-0003-2494-3671

ÖZET

Son yirmi yılda çocuk ve ergenlerde bildirilen yalnızlık düzeyleri, özellikle COVID-19 pandemisiyle birlikte belirgin şekilde artmış ve bu artış pandemi sonrasında da kısmen kalıcı hale gelmiştir. Almanya'da yürütülen ulusal ölçekli BELLA ve COPSY çalışmaları, 2020 yılında gençlerin yaklaşık üçte birinin sık sık yalnızlık hissettiğini, 2024 yılında bu oranın azalmakla birlikte hâlâ beşte bir düzeyinde seyrettiğini göstermektedir. Yalnızlık; depresyon, anksiyete, psikosomatik yakınmalar, akademik başarısızlık, madde kullanımı ve intihar riskiyle güçlü şekilde ilişkilendirilmekte, ergenlik döneminde artan yalnızlığın yetişkinlikte de sağlık ve iyilik hâli üzerinde kalıcı olumsuz etkiler bıraktığı gösterilmektedir. Bu derlemede, çocukluk ve ergenlik dönemindeki yalnızlığın kavramsal çerçevesi, epidemiyolojisi, risk faktörleri ve sağlık sonuçları güncel literatür ışığında ele alınmakta; hemşirelik sürecine dayalı değerlendirme, tanılama, girişim ve izlem yaklaşımları sunulmaktadır. Sonuç olarak, okul hemşireliği ve toplum ruh sağlığı hemşireliği uygulamalarının, sosyal destek sistemlerini güçlendiren ve erken dönemde müdahale eden bütüncül bir yaklaşımla yalnızlık salgınının olumsuz etkilerini azaltabileceği vurgulanmaktadır.

Anahtar Kelimeler: Yalnızlık, ergen sağlığı, çocuk sağlığı, sosyal izolasyon, hemşirelik yaklaşımı, ruh sağlığı.

ABSTRACT

Over the past twenty years, reported levels of loneliness among children and adolescents have increased significantly, particularly during the COVID-19 pandemic, and this increase has become partially persistent even after the pandemic. The nationwide BELLA and COPSY studies conducted in Germany show that in 2020, approximately one-third of young people frequently felt lonely, and while this proportion decreased by 2024, it still stood at one-fifth. Loneliness is strongly associated with depression, anxiety, psychosomatic complaints, academic failure, substance use, and suicide risk; it has been shown that increased loneliness during adolescence leaves lasting negative effects on health and well-being in adulthood. This review examines the conceptual framework, epidemiology, risk factors, and health outcomes of loneliness during childhood and adolescence in light of the current literature; it also presents nursing-based approaches to assessment, diagnosis, intervention, and follow-up. In conclusion, it is emphasized that school nursing and community mental health nursing practices can mitigate the negative effects of the loneliness epidemic through a holistic approach that strengthens social support systems and provides early intervention.

Keywords: Loneliness, adolescent health, child health, social isolation, nursing approach, mental health.

GİRİŞ

Yalnızlık, bireyin arzu ettiği ile fiilen sahip olduğu sosyal ilişkilerin niteliği ya da niceliği arasındaki öznel bir uyumsuzluk olarak tanımlanır ve nesnel bir yalnız kalma durumundan farklı olarak, kalabalık bir ortamda dahi hissedilebilen öznel bir deneyimdir (Mund et al., 2020; Seemann, 2022). Son yıllarda kamuoyu, sağlık politikası yapıcıları ve bilim insanları arasında yalnızlığın toplum sağlığını tehdit eden bir "salgın" düzeyine ulaştığına dair artan bir farkındalık oluşmuştur. Bu farkındalık, çocuk ve ergen popülasyonunu da kapsayacak biçimde genişlemiştir; zira gelişimsel açıdan akran ilişkileri ve aidiyet duygusunun en kritik olduğu bu dönemde yaşanan yalnızlık hem kısa hem de uzun vadeli sağlık sonuçları doğurmaktadır (Igami et al., 2023; Meier et al., 2026).

Dünya Sağlık Örgütü'nün sosyal bağlantı konusundaki küresel girişimleri ve çeşitli ülkelerin ulusal sağlık otoritelerinin yayımladığı raporlar, yalnızlığın artık yalnızca bir psikolojik rahatsızlık değil, kardiyovasküler hastalık, erken ölüm ve ruh sağlığı bozuklukları için bağımsız bir risk faktörü olarak ele alınması gerektiğini ortaya koymaktadır. Çocuk ve ergenler açısından bu tablo; dijitalleşme, sosyal medya kullanımının yaygınlaşması, pandemi sonrası toplumsal değişimler ve akademik baskı gibi çağdaş faktörlerle daha da karmaşık bir hal almaktadır (Akinyemi et al., 2025; Farrell et al., 2023).

Bu derlemenin amacı, çocukluk ve ergenlik dönemindeki yalnızlığın kavramsal çerçevesini, epidemiyolojik seyrini, risk faktörlerini ve sağlık üzerindeki sonuçlarını güncel literatür temelinde özetlemek; ayrıca bu alanda hemşirelik disiplininin üstlenebileceği değerlendirme, tanılama ve girişim rollerini bütüncül bir çerçevede sunmaktır.

YALNIZLIĞIN TANIMI VE KAVRAMSAL ÇERÇEVESİ

Yalnızlık literatürde genellikle iki boyutta ele alınır: duygusal yalnızlık (yakın, güvenli bir bağlanma figürünün eksikliği) ve sosyal yalnızlık (geniş bir sosyal ağın veya akran grubunun eksikliği). Çocuk ve ergenlerde bu iki boyut, gelişim dönemine göre farklı ağırlıklar taşır; okul öncesi dönemde ebeveyn-çocuk bağlanması, okul çağında akran kabulü, ergenlikte ise yakın arkadaşlık ve kimlik gelişimiyle iç içe geçen romantik ilişkiler ön plana çıkar (Mund et al., 2020; Timoszyk-Tomczak et al., 2025).

Yalnızlığın sosyal izolasyondan ayırt edilmesi önemlidir: sosyal izolasyon nesnel olarak ölçülebilen bir sosyal temas eksikliğini ifade ederken, yalnızlık öznel bir doyumsuzluk halidir. Bir çocuk kalabalık bir sınıf ortamında bulunsa dahi anlamlı bağlantı kuramadığını hissediyorsa yalnızlık yaşayabilir; tam tersine, sınırlı sosyal temasa sahip bazı çocuklar yalnızlık bildirmeyebilir. Bu ayrım, hemşirelik değerlendirmesinde nesnel gözlemin yanında öznel ifadenin de mutlaka sorgulanması gerektiğini göstermektedir (Mund et al., 2020; Timoszyk-Tomczak et al., 2025).

Kalıcı (kronik) yalnızlık, geçici yalnızlıktan farklı olarak kendini besleyen bir döngü oluşturabilir: yalnız hisseden çocuk/ergen, sosyal ortamları tehdit edici algılamaya başlar, bu da sosyal geri çekilmeyi ve olumsuz bilişsel şemaların pekişmesini beraberinde getirir. Bu

döngü zamanla sosyal becerilerin zayıflamasına ve yalnızlığın derinleşmesine yol açabilir (Verity et al., 2025).

EPİDEMİYOLOJİ VE GÜNCEL VERİLER

Çocuk ve ergenlerde yalnızlığın yaygınlığına ilişkin en güçlü kanıtlar, uzun soluklu ulusal izlem çalışmalarından gelmektedir. Almanya'da yürütülen BELLA (2003–2017) ve COPSY (2020–2024) çalışmalarının birleştirilmiş bulgularına göre, tek maddelik ölçümle belirlenen haftalık yalnızlık bildirimi COVID-19 pandemisinin doruk noktasında %34,6'ya ulaşmış; 2023 yılında %17,2'ye gerilemiş, ancak 2024 yılında yeniden %20,5'e yükselmiştir (Zoellner et al., 2025).

UCLA Yalnızlık Ölçeği ile yapılan daha katı ölçümlerde ise, geçtiğimiz bir yıl içinde belirgin düzeyde yalnızlık bildiren çocuk ve ergen oranı 2022'de %10,2 iken 2023'te %8,2'ye gerilemiş ve 2024'te bu düzeyde sabit kalmıştır (Zoellner et al., 2025). Bu veriler, pandemi sonrası kısmi bir iyileşme olmakla birlikte yalnızlığın pandemi öncesi düzeylerin üzerinde kalıcı bir toplum sağlığı sorunu haline geldiğini göstermektedir.

Almanya'daki Health Behavior in School-aged Children (HBSC) 2022 verilerinin analiz edildiği bir çalışmada, 11-15 yaş grubundaki 6.475 öğrenci arasında yalnızlığın ruh sağlığı sorunlarıyla güçlü biçimde ilişkili olduğu ve sosyal desteğin -özellikle 7. ve 9. sınıflarda öğretmen desteğinin- bu ilişkiyi tamponlayan bir rol oynadığı bulunmuştur (Schütz et al., 2025).

Avrupa çapında yürütülen ESPAD 2024 (Avrupa Okul Anketi Projesi–Alkol ve Diğer Uyuşturucular) verileri, Avrupalı ergenlerin dörtte birinden fazlasının düşük yaşam doyumu bildirdiğini; UNICEF'in Dünya Çocuklarının Durumu 2024 raporu ise gençlerin yalnızlık, akademik baskı ve dijital aşırı maruziyetle karakterize bir "bağlantısızlık salgını" yaşadığını vurgulamaktadır. Ayrıca Dünya Sağlık Örgütü verilerine göre depresyon ve anksiyete, ergenlerde engelliliğin önde gelen nedenleri arasında yer almakta, intihar ise 15-19 yaş grubunda ikinci sıradaki ölüm nedeni olmaya devam etmektedir (Ferreira Díaz, 2026).

RİSK FAKTÖRLERİ VE ETKİLEYEN BAĞLAMSAL ETMENLER

- Aile ve bağlanma faktörleri: Ebeveyn-çocuk ilişkisinin niteliği, aile içi çatışma, boşanma/parçalanmış aile yapısı, ebeveyn hastalığı (örneğin bir ebeveynin kanser tanısı alması) yalnızlık riskini artırabilir (Mckeown et al., 2025).
- Okul ortamı: Zorbalık ve siber zorbalığa maruz kalma, okulda güvende hissetmeme, öğretmen ilgisinin/desteğinin azlığı (Schütz et al., 2025).
- Dijital ortam ve sosyal medya: Yüz yüze etkileşimin yerini alan yoğun ekran kullanımı, sosyal karşılaştırma ve dışlanma algısı yaratabilen sosyal medya dinamikleri.
- Sosyoekonomik koşullar: Düşük gelir düzeyi, göç geçmişi, sosyal dışlanma ve ayrımcılık deneyimleri.
- Bireysel özellikler: Utangaçlık, düşük öz-saygı, kronik hastalık veya engellilik durumu, nörogelişimsel farklılıklar (örneğin otizm spektrum bozukluğu).
- Toplumsal/küresel etkenler: COVID-19 pandemisi sırasında uygulanan okul kapanmaları ve sosyal mesafe uygulamaları, yalnızlık oranlarında belirgin ve kalıcı etkisi olan bir dönüm noktası oluşturmuştur (Farrell et al., 2023).

SAĞLIK SONUÇLARI

Yalnızlığın çocuk ve ergen sağlığı üzerindeki etkileri çok boyutludur ve ruhsal, bedensel, akademik/sosyal ve davranışsal alanları kapsar. Add Health kohortu üzerinden yürütülen geniş kapsamlı bir sonuç-genişletilmiş (outcome-wide) boylamsal çalışma, ergenlik döneminde artan yalnızlığın yetişkinlikte incelenen 41 farklı sağlık ve iyilik hâli göstergesinin pek çoğunda olumsuz sonuçlarla ilişkili olduğunu göstermiştir; bu bulgu, ergenlik dönemi yalnızlığının yetişkinlik sağlığı için erken bir belirleyici olabileceğine işaret etmektedir (Kim et al., 2025).

Tablo 1. Yalnızlığın Çocuk ve Ergenlerde Sağlık Sonuçları ve Olası Mekanizmalar

Sağlık Alanı	Gözlenen Sonuçlar	Olası Mekanizmalar
Ruh Sağlığı	Depresyon, anksiyete, sosyal anhedoni, düşük öz-saygı, kendine zarar verme ve intihar düşüncesinde artış	Olumsuz bilişsel şemalar, sosyal geri çekilme döngüsü, aidiyet ihtiyacının karşılanmaması
Psikosomatik Belirtiler	Baş ağrısı, karın ağrısı, uyku bozuklukları, yorgunluk, iştah değişiklikleri	Kronik stres yanıtı, otonom sinir sistemi aktivasyonu, kortizol düzensizliği
Akademik ve Sosyal İşlevsellik	Okula devamsızlık, akademik başarıda düşüş, akran ilişkilerinde bozulma, zorbalığa maruz kalma riskinde artış	Azalmış sosyal destek, güven duygusunda azalma, sınıf ortamına katılımın azalması
Davranışsal Riskler	Madde kullanımı, aşırı ekran/sosyal medya kullanımı, riskli davranışlara yönelim	Duygu düzenleme güçlüğü, dijital ortamda telafi arayışı, akran onayı arayışı
Uzun Dönem / Yetişkinlik Sonuçları	Yetişkinlikte kronik hastalık riski, düşük yaşam doyumu, sosyal işlevsellikte kalıcı bozulma	Ergenlik döneminde gelişen olumsuz sosyal bilişin kalıcılaşması, biyolojik "aşınma" (allostatik yük)

Bu tabloda özetlenen bulgular, yalnızlığın izole bir duygusal deneyim olmadığını, aksine biyopsikososyal bir bütünlük içinde çoklu sağlık alanlarını etkileyen bir risk faktörü olduğunu ortaya koymaktadır. Özellikle psikosomatik yakınmalar ile yalnızlık arasındaki ilişki, hemşirelik değerlendirmesinde sıklıkla gözden kaçabilecek, ancak erken uyarı işareti olarak değerlendirilmesi gereken önemli bir bulgudur (Lederman, 2023; Matthews et al., 2022; Warren, 2026).

HEMŞİRELİK YAKLAŞIMLARI

Hemşireler; okul, birinci basamak, ayaktan ruh sağlığı hizmetleri ve hastane ortamları da dâhil olmak üzere çocuk ve ergenlerle en sık ve en düzenli temas kuran sağlık profesyonelleri arasındadır. Bu konum, yalnızlığın erken tanınması ve önlenmesinde hemşirelik disiplinine kritik bir sorumluluk yüklemektedir (Farrell et al., 2023). Hemşirelik süreci -değerlendirme, tanılama, planlama, uygulama ve değerlendirme- yalnızlığa yönelik bütüncül ve bireyselleştirilmiş bir yaklaşımın çerçevesini oluşturur.

Tablo 2. Hemşirelik Sürecine Göre Yalnızlığa Yönelik Uygulama Örnekleri

Hemşirelik Süreci Basamağı	**Uygulama Örnekleri**
Değerlendirme (Assessment)	Standart ölçekler (UCLA Yalnızlık Ölçeği, Good Childhood Index) ile tarama; okul, aile ve akran ilişkilerinin sorgulanması; psikosomatik yakınmaların yalnızlıkla ilişkisinin değerlendirilmesi, intihar/kendine zarar verme riskinin taranması
Hemşirelik Tanısı	"Sosyal İzolasyon", "Bozulmuş Sosyal Etkileşim", "Etkisiz Baş Etme", "Benlik Saygısında Düşme Riski" gibi NANDA-I tanılarının bireyselleştirilmiş biçimde kullanılması
Planlama	Çocuk/ergen, aile, öğretmen ve okul psikolojik danışmanıyla birlikte gerçekçi ve ölçülebilir hedeflerin belirlenmesi; kültürel ve gelişimsel açıdan uygun yaklaşımların seçilmesi
Uygulama	Bireysel danışmanlık ve aktif dinleme; sosyal beceri eğitimi, akran destek programları, aile eğitimi; okul temelli aidiyet programları, dijital okuryazarlık ve sağlıklı ekran kullanımı danışmanlığı; gerektiğinde ruh sağlığı uzmanına yönlendirme
Değerlendirme (Evaluation)	Yalnızlık düzeyinde, psikosomatik yakınmalarda ve sosyal katılımda izlenebilir değişimlerin düzenli aralıklarla yeniden ölçülmesi

Okul Hemşireliği ve Toplum Temelli Yaklaşımlar

Okul hemşireleri, çocuk ve ergenlerin günlük yaşamına en yakın konumdaki sağlık profesyonelleri olarak, sınıf içi gözlem, sağlık taramaları ve düzenli danışmanlık görüşmeleri yoluyla yalnızlık belirtilerini erken fark edebilir. Okul temelli aidiyet programları, akran destek/akran mentorluk sistemleri ve öğretmenlerle iş birliği içinde yürütülen sosyal-duygusal öğrenme müfredatları, yalnızlığı azaltmaya yönelik etkili yaklaşımlar arasında yer almaktadır. Ayrıca, disiplinler arası iş birliği -okul psikolojik danışmanı, sosyal hizmet uzmanı ve öğretmenlerle koordineli çalışma- bu programların etkinliğini artırmaktadır (Kim et al., 2025; Longobucco et al., 2023; Meier et al., 2026).

Bireysel ve Aile Odaklı Girişimler

Bireysel görüşmelerde hemşire, yargılamayan ve güven verici bir tutumla açık uçlu sorular sorarak çocuğun/ergenin duygusal deneyimini ifade etmesine alan açmalıdır. Aile eğitimi, ebeveynlerin çocuklarının sosyal dünyasına ilgi göstermesini, dinleme becerilerini geliştirmesini ve aşırı koruyucu ya da ihmalkâr tutumlardan kaçınmasını desteklemeyi amaçlar. Ebeveyn hastalığı, boşanma gibi aile içi stres etkenlerinin bulunduğu durumlarda, hemşire aileyi ilgili psikososyal destek kaynaklarına yönlendirme rolünü üstlenebilir (Kim et al., 2025; Nambu & Matsuoka, 2025).

Dijital Sağlık Okuryazarlığı ve Sosyal Medya Danışmanlığı

Ekran süresinin niteliği ve niceliği hakkında yapılandırılmış danışmanlık, sosyal medyanın pasif tüketim yerine aktif ve anlamlı etkileşim amacıyla kullanılmasını teşvik eden yaklaşımlar, hemşirelik girişimlerinin önemli bir bileşenidir. Bu kapsamda, ekran süresini tamamen kısıtlamak yerine, çevrimiçi ve çevrimdışı sosyal etkileşim dengesinin kurulmasına odaklanan motivasyonel görüşme teknikleri önerilmektedir (Godard & Holtzman, 2023; Jones et al., 2021; Lua et al., 2023).

Risk Taraması ve Sevk

Yalnızlık ile depresyon, kendine zarar verme ve intihar düşüncesi arasındaki güçlü ilişki göz önüne alındığında, hemşireler yalnızlık bildiren çocuk/ergenlerde mutlaka ruh sağlığı riskini sorgulamalı, gerektiğinde çocuk ve ergen ruh sağlığı uzmanına zamanında sevk sağlamalıdır. Erken tanı ve müdahalenin, kronikleşmiş ruh sağlığı sorunlarının önlenmesinde belirleyici olduğu bilinmektedir (Farley, 2020; Zeas-Sigüenza et al., 2025).

SONUÇ VE ÖNERİLER

Çocuk ve ergenlerde yalnızlık, günümüzde toplum sağlığını ilgilendiren, çok boyutlu sonuçları olan ve önlenebilir bir sorun olarak ele alınmalıdır. Güncel epidemiyolojik veriler, pandemi sonrası kısmi bir gerilemeye rağmen yalnızlığın önemli bir kesimde kalıcı hale geldiğini göstermektedir. Yalnızlığın ruhsal, bedensel, akademik ve uzun dönemli sağlık sonuçları, bu konunun yalnızca bireysel bir duygusal deneyim değil, sistemli bir sağlık politikası önceliği olarak ele alınmasını gerektirmektedir.

Hemşirelik disiplini; okulda, toplumda ve klinik ortamlarda erken tanıma, bütüncül değerlendirme ve bireyselleştirilmiş girişimler yoluyla yalnızlık salgınının olumsuz etkilerinin azaltılmasında kilit bir role sahiptir. Gelecekteki uygulamalarda; standart yalnızlık tarama araçlarının rutin sağlık değerlendirmelerine entegre edilmesi, okul hemşireliği kapasitesinin güçlendirilmesi, disiplinler arası iş birliğinin artırılması ve yalnızlığa yönelik girişimlerin etkililiğini değerlendiren daha fazla boylamsal ve müdahale araştırmasının yürütülmesi önerilmektedir.

KAYNAKLAR

Akinyemi, O., Abdulrazaq, W., Fasokun, M., Ogunyankin, F., Ikugbayigbe, S., Nwosu, U., Michael, M., Hughes, K., & Ogundare, T. (2025). The impact of loneliness on depression, mental health, and physical well-being. *PLOS One*, *20*(7), e0319311. https://doi.org/10.1371/journal.pone.0319311

Farley, H. R. (2020). Assessing mental health in vulnerable adolescents. *Nursing*, *50*(10), 48–53. https://doi.org/10.1097/01.NURSE.0000697168.39814.93

Farrell, A. H., Vitoroulis, I., Eriksson, M., & Vaillancourt, T. (2023). Loneliness and Well-Being in Children and Adolescents during the COVID-19 Pandemic: A Systematic Review. *Children*, *10*(2), 279. https://doi.org/10.3390/children10020279

Ferreira Díaz, M. J. (2026). Adolescence in crisis: Insights from the 2024 ESPAD report and a nursing perspective. *IScience*, *29*(2), 114726. https://doi.org/10.1016/j.isci.2026.114726

Godard, R., & Holtzman, S. (2023). Are active and passive social media use related to mental health, wellbeing, and social support outcomes? A meta-analysis of 141 studies. *Journal of Computer-Mediated Communication*, *29*(1). https://doi.org/10.1093/jcmc/zmad055

Igami, K., Hosozawa, M., Ikeda, A., Bann, D., Shimizu, T., & Iso, H. (2023). Adolescent Loneliness in 70 Countries Across Africa, America, and Asia: A Comparison of Prevalence and Correlates. *Journal of Adolescent Health*, *72*(6), 906–913. https://doi.org/10.1016/j.jadohealth.2022.12.029

Jones, A., Armstrong, B., Weaver, R. G., Parker, H., von Klinggraeff, L., & Beets, M. W. (2021). Identifying effective intervention strategies to reduce children's screen time: a systematic review and meta-analysis. *International Journal of Behavioral Nutrition and Physical Activity*, *18*(1), 126. https://doi.org/10.1186/s12966-021-01189-6

Kim, E. S., Wilkinson, R., Holt-Lunstad, J., & VanderWeele, T. J. (2025). Loneliness During Adolescence and Subsequent Health and Well-Being in Adulthood: An Outcome-Wide Longitudinal Approach. *Journal of Adolescent Health*, *77*(1), 66–75. https://doi.org/10.1016/j.jadohealth.2024.12.011

Lederman, Z. (2023). Loneliness—a clinical primer. *British Medical Bulletin*, *145*(1), 132–140. https://doi.org/10.1093/bmb/ldad003

Longobucco, Y., Ricci, M., Scrimaglia, S., Camedda, C., Dallolio, L., & Masini, A. (2023). Effects of School Nurse-Led Interventions in Collaboration with Kinesiologists in Promoting Physical Activity and Reducing Sedentary Behaviors in Children and Adolescents: A Systematic Review. *Healthcare*, *11*(11), 1567. https://doi.org/10.3390/healthcare11111567

Lua, V. Y. Q., Chua, T. B. K., & Chia, M. Y. H. (2023). A Narrative Review of Screen Time and Wellbeing among Adolescents before and during the COVID-19 Pandemic: Implications for the Future. *Sports*, *11*(2), 38. https://doi.org/10.3390/sports11020038

Matthews, T., Bryan, B. T., Danese, A., Meehan, A. J., Poulton, R., & Arseneault, L. (2022). Using a Loneliness Measure to Screen for Risk of Mental Health Problems: A Replication in Two Nationally Representative Cohorts. *International Journal of Environmental Research and Public Health*, *19*(3), 1641. https://doi.org/10.3390/ijerph19031641

Mckeown, L., Campbell, K., Dempster, M., Groarke, J., & Graham-Wisener, L. (2025). Adolescent Loneliness When a Parent Has Cancer: A Qualitative Systematic Review. *Psycho-Oncology*, *34*(4). https://doi.org/10.1002/pon.70148

Meier, Z., Helvich, J., Novak, L., Furstova, J., Machu, V., Vagner, L., & Tavel, P. (2026). Loneliness and adolescent mental health: a multilevel examination of socio-ecological factors across Czech schools. *Child and Adolescent Psychiatry and Mental Health*, *20*(1), 23. https://doi.org/10.1186/s13034-026-01026-3

Mund, M., Freuding, M. M., Möbius, K., Horn, N., & Neyer, F. J. (2020). The Stability and Change of Loneliness Across the Life Span: A Meta-Analysis of Longitudinal Studies. *Personality and Social Psychology Review*, *24*(1), 24–52. https://doi.org/10.1177/1088868319850738

Nambu, M., & Matsuoka, M. (2025). Nurses' Attitudes in Adolescent Oncology Care: Supporting Adolescents While Navigating Emotional Challenges and Fostering Mutual Subjectivity. *Cureus*. https://doi.org/10.7759/cureus.81049

Schütz, R., Reiss, F., Moor, I., Kaman, A., & Bilz, L. (2025). Lonely children and adolescents are less healthy and report less social support: A study on the effect of loneliness on mental health and the moderating role of social support. *BMC Public Health*, *25*(1), 2172. https://doi.org/10.1186/s12889-025-23247-5

Seemann, A. (2022). The Psychological Structure of Loneliness. *International Journal of Environmental Research and Public Health*, *19*(3), 1061. https://doi.org/10.3390/ijerph19031061

Timoszyk-Tomczak, C., Pieńkowska, E., Ligocka, M., & Piłat, M. (2025). Adolescents' Feelings of Loneliness Considering Anxiety and Intrafamilial Relations. *Brain Sciences*, *15*(12), 1270. https://doi.org/10.3390/brainsci15121270

Verity, L., Burke, L., & Qualter, P. (2025). Loneliness in adolescence: prevalence, developmental contexts, and interventions. *Frontiers in Psychiatry*, *16*. https://doi.org/10.3389/fpsyt.2025.1696981

Warren, A. (2026). Loneliness as a Vital Sign: Toward a Biopsychosocial Reframing of Social Disconnection. *Journal of Primary Care & Community Health*, *17*. https://doi.org/10.1177/21501319261426724

Zeas-Sigüenza, A., Voldstad, A., Ruisoto, P., Ganho-Ávila, A., Guiomar, R., Cacho, R., Muntané, F., & Benach, J. (2025). Loneliness as a Public Health Challenge: A

Systematic Review and Meta-Analysis to Inform Policy and Practice. *European Journal of Investigation in Health, Psychology and Education*, *15*(7), 131. https://doi.org/10.3390/ejihpe15070131

Zoellner, F., Erhart, M., Schütz, R., Napp, A.-K., Devine, J., Reiss, F., Ravens-Sieberer, U., & Kaman, A. (2025). Two decades of loneliness among children and adolescents: longitudinal trends, risks and resources – Results from the German BELLA and COPSY studies. *European Child & Adolescent Psychiatry*, *34*(11), 3629–3641. https://doi.org/10.1007/s00787-025-02779-6

YAPAY ZEKÂ DESTEKLİ ÇOCUK VE ERGEN HEMŞİRELİĞİ: GÜNCEL UYGULAMALAR, FIRSATLAR VE ETİK SORUNLAR

ARTIFICIAL INTELLIGENCE-ASSISTED NURSING CARE FOR CHILDREN AND ADOLESCENTS: CURRENT PRACTICES, OPPORTUNITIES, AND ETHICAL ISSUES

Assoc. Prof. Dr. Hakan AVAN [1]
Afşin School of Health, Kahramanmaraş Sütçü İmam University, Kahramanmaraş, TURKEY
ORCID ID: 0000-0003-2494-3671

ÖZET

Bu geleneksel derlemenin amacı, yapay zekâ teknolojilerinin çocuk ve ergen hemşireliği pratiğindeki güncel uygulamalarını, sunduğu fırsatları ve beraberinde getirdiği etik sorunları kapsamlı bir şekilde incelemektir. Mevcut literatürü sentezleyerek, bu alandaki bilgi boşluklarını belirlemek ve gelecek araştırmalar ile klinik uygulamalar için kanıta dayalı öneriler sunmak hedeflenmiştir. Bu çalışmada geleneksel (anlatısal) derleme yöntemi kullanılmıştır. Literatür taraması, PubMed, Scopus, Web of Science ve Google Scholar veri tabanlarında, "yapay zekâ", "çocuk hemşireliği", "ergen hemşireliği", "etik", "klinik karar destek sistemleri" ve "makine öğrenmesi" anahtar kelimeleri kullanılarak gerçekleştirilmiştir. Tarama, 2019-2026 yılları arasında yayınlanmış, çocuk ve ergen hemşireliğinde yapay zekâ uygulamalarına odaklanan orijinal araştırmalar, sistematik incelemeler ve meta-analizler ile sınırlandırılmıştır. Belirlenen çalışmalar, derleme konusuyla ilgililikleri ve bilimsel katkıları açısından değerlendirilerek sentezlenmiştir. Derleme bulguları, yapay zekanın çocuk ve ergen hemşireliğindeki uygulamalarının dört ana başlık altında toplanabileceğini göstermektedir: Klinik uygulamalar (erken tanı, risk tahmini, klinik karar destek sistemleri, hasta izleme, ilaç yönetimi), Eğitim ve bilgi yönetimi (sanal simülasyonlar, kişiselleştirilmiş öğrenme, generatif yapay zekâ ile materyal geliştirme), Yönetim ve kaynak optimizasyonu (iş akışı iyileştirme, personel planlaması), ve Halk sağlığı uygulamaları (siber zorbalık tespiti, çevrimiçi risk izleme). Yapay zekanın sağladığı önemli fırsatlar arasında tanısal doğruluğun artması, bakımın kişiselleştirilmesi ve hemşirelerin idari yükünün azalması yer alırken; başlıca etik sorunlar olarak veri gizliliği ve güvenliği, algoritmik önyargı, hasta/ebeveyn katılımı, şeffaflık ve hesap verebilirlik sorunları öne çıkmaktadır. Ayrıca, mevcut araştırmaların çoğunlukla algoritmik performansa odaklandığı, klinik uygulanabilirlik ve gerçek yaşam doğrulama çalışmalarının sınırlı olduğu belirlenmiştir. Yapay zekâ teknolojileri, çocuk ve ergen hemşireliğinde bakım kalitesini artırma ve sağlık hizmetlerini dönüştürme konusunda büyük bir potansiyele sahiptir. Ancak bu potansiyelin güvenli, adil ve etik bir şekilde gerçekleştirilebilmesi için hemşirelerin yapay zekâ okuryazarlığının artırılması, hasta ve ebeveynlerin sürece dahil edilmesi, multidisipliner iş birliğinin güçlendirilmesi ve kapsamlı etik-yasal çerçevelerin oluşturulması zorunludur. Gelecek araştırmalar, algoritmik başarıdan çok klinik entegrasyon, uzun vadeli hasta sonuçları ve sağlık eşitliği üzerinde yoğunlaşmalıdır.

Anahtar Kelimeler: Yapay zekâ, çocuk hemşireliği, ergen hemşireliği, etik, klinik karar destek sistemleri, makine öğrenmesi, pediatri.

ABSTRACT

This traditional review aims to comprehensively examine the current applications, opportunities, and ethical issues of artificial intelligence technologies in child and adolescent nursing practice. By synthesizing the existing literature, it also seeks to identify knowledge gaps and provide evidence-based recommendations for future research and clinical practice. A traditional (narrative) review method was employed. The literature search was conducted using PubMed, Scopus, Web of Science, and Google Scholar databases with keywords including "artificial intelligence," "pediatric nursing," "adolescent nursing," "ethics," "clinical decision support systems," and "machine learning." The search was limited to original research, systematic reviews, and meta-analyses published between 2019-2026, focusing on artificial intelligence applications in child and adolescent nursing. Identified studies were synthesized based on their relevance to the review topic and scientific contribution. The review findings indicate that AI applications in child and adolescent nursing can be categorized into four main areas: Clinical applications (early diagnosis, risk prediction, clinical decision support systems, patient monitoring, medication management), Education and knowledge management (virtual simulations, personalized learning, generative artificial intelligence for material development), Management and resource optimization (workflow improvement, staff planning), and Public health applications (cyberbullying detection, online risk monitoring). Significant opportunities include increased diagnostic accuracy, personalized care, and reduced administrative burden for nurses. However, major ethical concerns include data privacy and security, algorithmic bias, patient/parent participation, transparency, and accountability issues. Furthermore, most current research focuses on algorithmic performance, with limited studies on clinical applicability and real-world validation. Artificial intelligence technologies hold great potential to enhance care quality and transform healthcare services in child and adolescent nursing. However, to realize this potential safely, fairly, and ethically, it is imperative to increase nurses' artificial intelligence literacy, involve patients and parents in the process, strengthen multidisciplinary collaboration, and establish comprehensive ethical-legal frameworks. Future research should focus not only on algorithmic success but also on clinical integration, long-term patient outcomes, and health equity.

Keywords: Artificial intelligence, pediatric nursing, adolescent nursing, ethics, clinical decision support systems, machine learning, pediatrics.

GİRİŞ

Çocuk ve ergen sağlığı, gelişimsel, psikososyal ve fizyolojik dinamikleri nedeniyle sağlık hizmetlerinin en hassas ve karmaşık alanlarından birini oluşturmaktadır.(Mastorci et al., 2024) Bu hassas nüfusa yönelik bakım hizmetlerinin sunumunda karşılaşılan zorluklar, yenilikçi teknolojik çözümlere olan ihtiyacı giderek artırmaktadır.(Cortese et al., 2026) Yapay zekâ (YZ) teknolojilerinin sağlık sektörü içindeki rolü, özellikle çocuk ve ergen hemşireliği pratiğinde giderek daha fazla önem kazanmaktadır. Bu tür teknolojiler, tanı süreçlerinden tedavi

planlamasına, hasta izlemeye ve eğitim materyalleri geliştirmeye kadar çok çeşitli uygulama alanlarına sahiptir.(Alowais et al., 2023; Cellina et al., 2023)

Bununla birlikte, çocuk ve ergen nüfusunun kendine özgü gelişimsel, psikososyal ve etik gereklilikleri, YZ'nin bu alandaki entegrasyonunu karmaşık hale getirerek özel bir dikkat gerektirmektedir.(Yang et al., 2025) Literatürdeki mevcut çalışmalar, YZ'nin pediatrik alandaki algoritmik başarısına odaklanmakta; ancak bu teknolojilerin klinik uygulamaya entegrasyonu, uzun vadeli hasta sonuçlarına etkisi ve beraberinde getirdiği etik zorluklar yeterince ele alınmamaktadır.(Ganatra, 2025; Kerth et al., 2024; Lee et al., 2025; Y. Park et al., 2025) Bu çerçevede, yapay zekâ destekli çocuk ve ergen hemşireliği uygulamalarının yanı sıra mevcut fırsatlar ve etik sorunların kapsamlı bir şekilde ele alınması önemli bir ihtiyaçtır.

Bu derlemenin amacı, YZ teknolojilerinin çocuk ve ergen hemşireliği pratiğindeki güncel uygulamalarını, sunduğu fırsatları ve beraberinde getirdiği etik sorunları kapsamlı bir şekilde incelemektir. Makale, mevcut literatürü sistematik olarak sentezleyerek alandaki bilgi boşluklarını belirlemeyi ve hem klinik uygulama hem de gelecek araştırmalar için kanıta dayalı öneriler sunmayı hedeflemektedir. Bu bağlamda, çalışmanın özgün katkısı, çocuk ve ergen hemşireliği bağlamında YZ uygulamalarını bütüncül bir perspektifle (klinik, eğitim, yönetim, etik) ele alarak, disiplinler arası bir sentez sunmasıdır.

Yöntem

Bu çalışma, geleneksel (anlatısal) derleme yöntemi kullanılarak gerçekleştirilmiştir. Geleneksel derleme yöntemi, belirli bir konudaki mevcut literatürü geniş bir perspektiften sentezleyerek, konunun mevcut durumunu, karmaşıklığını ve araştırma boşluklarını ortaya koymak amacıyla tercih edilmiştir.(Li & Wang, 2018; Reddy et al., 2020)

Literatür Tarama Stratejisi

Literatür taraması, 01-30 Mayıs 2026 tarihleri arasında dört elektronik veri tabanında gerçekleştirilmiştir: PubMed, Scopus, Web of Science ve Google Scholar. Taramada kullanılan anahtar kelimeler, tıbbi konu başlıkları (MeSH) ve Boolean operatörleri kullanılarak birleştirilmiştir. Anahtar kelimeler şunlardır: "artificial intelligence", "machine learning", "pediatric nursing", "adolescent nursing", "child health nursing", "clinical decision support systems", "ethics", "pediatrics", ve "nursing education".

Dahil Etme/ Hariç Tutma Kriterleri:

Çalışmaya, çocuk (0–18 yaş) ve/veya ergen sağlığına odaklanan, hemşirelik pratiği, eğitimi veya yönetimi kapsamında yapay zekâ, makine öğrenmesi ya da bunların alt alanlarını (örneğin derin öğrenme ve doğal dil işleme) ele alan çalışmalar dahil edilmiştir. İncelemeye, 2019–2026 yılları arasında Türkçe veya İngilizce yayımlanmış orijinal araştırmalar, sistematik derlemeler, meta-analizler ve derleme makaleleri alınmıştır. Buna karşın, yalnızca yetişkin sağlığına odaklanan çalışmalar, yapay zekânın hemşirelik uygulamalarıyla doğrudan ilişkili olmadığı yalnızca algoritma geliştirmeye yönelik teknik araştırmalar, editöre mektuplar, vaka raporları, konferans özetleri ile Türkçe ve İngilizce dışındaki dillerde yayımlanan çalışmalar inceleme kapsamı dışında bırakılmıştır. Veri sentezi sürecinde, dahil edilen çalışmaların bulguları yapay zekânın çocuk ve ergen hemşireliğindeki uygulama alanları, sağladığı fırsatlar ve karşılaşılan

etik sorunlar temelinde tematik olarak sınıflandırılarak sentezlenmiştir. Ayrıca, çalışmaların metodolojik özellikleri değerlendirilmiş, bulgular arasındaki benzerlikler ve farklılıklar karşılaştırmalı olarak ele alınmış, çelişkili sonuçlar ise ilgili literatür ışığında tartışılmıştır.

Bulgular

Derleme bulguları, YZ'nin çocuk ve ergen hemşireliğindeki uygulamalarının dört ana başlık altında toplanabileceğini göstermektedir. Bu başlıklar ve alt bileşenleri Tablo 1, Tablo 2, Tablo 3 ve Tablo 4'te özetlenmiştir.

Tablo 1. Derlemeye Dahil Edilen Çalışmaların Genel Özellikleri

Yazar(lar) ve Yıl	Çalışma Türü	YZ Uygulama Alanı	Anahtar Bulgular
(Y. Park et al., 2025)	Sistematik İnceleme	Klinik Uygulama (Erken Uyarı)	YZ modelleri, sepsis ve basınç yaraları gibi komplikasyonları tahmin etmede yüksek performans göstermiştir.
(Zhao et al., 2022)	Orijinal Araştırma	Klinik Uygulama (KKDS)	Pediatrik hipertermi yönetiminde KKDS, hemşirelik değerlendirmelerinin tamlığını artırmıştır.
(Williams et al., 2026)	Kapsamlı Tarama	Eğitim (Generatif YZ)	GenAI, pediatrik hemşirelik eğitiminde senaryo tabanlı öğrenme ve kişiselleştirilmiş geri bildirim sağlamaktadır.
(Alqaraleh et al., 2025)	Sistematik İnceleme	Klinik Uygulama (İlaç Yönetimi)	YZ destekli sistemler ilaç hatalarını önemli ölçüde azaltmaktadır.
(Sultan et al., 2023)	Derleme	Halk Sağlığı (Siber Zorbalık)	Makine öğrenmesi teknikleri siber zorbalık tespitinde etkilidir.
(Sisk et al., 2020)	Orijinal Araştırma	Etik (Ebeveyn Algısı)	Ebeveynler YZ'ye açık ancak gizlilik ve adalet konusunda endişelidir.
(Reinhart et al., 2026)	Niteliksel Çalışma	Etik (Çocuk Algısı)	Çocuklar YZ'yi destekleyici bir araç olarak görmekte, nihai kararın insanlarda olmasını istemektedir.
(Sezgin & Boch, 2026)	Konsept Makale	Etik & Politika	Pediatrik YZ uygulamaları için PAIR hazırlık çerçevesini önermiştir.

1. Yapay Zekânın Çocuk ve Ergen Hemşireliğinde Klinik Uygulama Alanları

Klinik uygulamalar, YZ'nin bu alandaki en yoğun ve en umut verici kullanım alanını oluşturmaktadır.

Klinik Karar Destek Sistemleri (KKDS): YZ tabanlı KKDS'ler, hemşirelerin karmaşık klinik verileri analiz etmelerine ve kanıta dayalı kararlar almalarına yardımcı olur. Zhao ve

arkadaşlarının (2022) geliştirdiği pediatrik hipertermi yönetimine yönelik KKDS, hemşirelik değerlendirmelerinin tamlığını artırmış ve bakım girişimlerinin hasta gereksinimlerine göre uyarlanmasını kolaylaştırmıştır. (Zhao et al., 2022) Benzer şekilde, Lu ve arkadaşlarının (2021) önerdiği entegre KKDS modeli, karmaşık ve çoklu kronik hastalığı bulunan çocukların bakımında hemşirelerin farklı bilgi kaynaklarını bütünleştirerek daha etkili klinik kararlar almasına olanak sağlayacak bir çerçeve sunmaktadır. (Lu et al., 2021) Clausen ve arkadaşlarının (2023) çocuk ve ergen ruh sağlığı hizmetlerinde geliştirdiği IDDEAS sistemi ise klinisyenler tarafından yararlı bulunmasına rağmen, elektronik sağlık kayıtlarıyla entegrasyonun sağlanmasının ve nihai klinik kararın sağlık profesyonelleri tarafından verilmesinin güvenli kullanım açısından temel gereklilikler olduğunu ortaya koymuştur. (Clausen et al., 2023)

Erken Uyarı ve Tahmine Dayalı Analitik Sistemler: YZ destekli erken uyarı sistemleri, kritik bakım ortamlarında hastaların durumundaki değişiklikleri gerçek zamanlı olarak izleyerek komplikasyonların erken tespitine olanak tanır. Park ve arkadaşlarının (2025) kritik bakım hemşireliğinde yaptığı inceleme, YZ uygulamalarının özellikle basınç yaraları ve sepsis gibi komplikasyonların öngörülmesinde yüksek performans gösterdiğini ortaya koymuştur. (Y. Park et al., 2025) Bu sistemler, hemşirelerin iş yükünü azaltarak kaynak tahsisini optimize etmekte ve hasta güvenliğini artırmaktadır.

Tablo 2. Yapay Zekânın Çocuk ve Ergen Hemşireliğinde Uygulama Alanları

Uygulama Alanı	Alt Kategoriler	Amaç / Fayda
Klinik Uygulamalar	Klinik Karar Destek Sistemleri (KKDS)	Hemşirelerin kanıta dayalı, kişiselleştirilmiş kararlar almasını desteklemek.
	Erken Uyarı ve Tahmine Dayalı Analitik	Komplikasyonları (sepsis, basınç yarası) erken tespit etmek.
	Hasta İzleme ve Veri Yönetimi	Karmaşık verileri analiz ederek hasta durumunu gerçek zamanlı izlemek.
	İlaç Yönetimi	İlaç hatalarını azaltmak, güvenli doz hesaplamalarını desteklemek.
Eğitim ve Bilgi Yönetimi	Sanal Simülasyonlar ve Kişiselleştirilmiş Öğrenme	Hemşirelik öğrencilerinin uygulama becerilerini güvenli ortamda geliştirmek.
	Generatif YZ ile Materyal Geliştirme	Hasta eğitimi ve hemşirelik eğitimi için özelleştirilmiş içerik üretmek.
Yönetim ve Kaynak Optimizasyonu	İş Akışı İyileştirme ve Personel Planlaması	İdari iş yükünü azaltmak, personel dağılımını optimize etmek, verimliliği artırmak.

Uygulama Alanı	Alt Kategoriler	Amaç / Fayda
Halk Sağlığı Uygulamaları	Siber Zorbalık Tespiti ve Çevrimiçi Risk İzleme	Ergenleri çevrimiçi risklerden korumak ve ruh sağlığını izlemek.

İzleme ve Veri Yönetimi: Klinik ortamlarda yüksek kaliteli veri toplanması önemli bir zorluktur. Ganatra (2025), Yan ve ark., Jang ve Lee (2026) gibi araştırmacılar, geliştirilen yazılım araçlarının davranışsal ve fizyolojik verilerin sistematik olarak toplanmasını kolaylaştırarak erken uyarı sistemlerinin doğruluğunu artırma potansiyeline sahip olduğunu vurgulamaktadır. (Ganatra, 2025; Jang & Lee, 2026; Shen, 2026; Yan et al., 2025)

İlaç Yönetimi: YZ, ilaç hatalarını azaltmada önemli bir rol oynamaktadır. Alqaraleh ve arkadaşlarının (2025) sistematik incelemesi, klinik karar destek sistemleri, akıllı infüzyon pompaları ve otomatik reçete doğrulama araçlarının ilaç hatalarını önemli ölçüde azalttığını göstermektedir. (Alqaraleh et al., 2025) Çocuk hastalarda ilaç güvenliğinin kritik önemi düşünüldüğünde, bu teknolojilerin yaygınlaştırılması büyük önem taşımaktadır.

2. Yapay Zekânın Eğitim ve Bilgi Yönetiminde Kullanımı

YZ teknolojileri, hemşirelik eğitimini dönüştürme potansiyeline sahiptir.

Sanal Simülasyonlar ve Kişiselleştirilmiş Öğrenme: CAN ve arkadaşları (2021), sanal hastalar üzerinden verilen eğitimlerin sağlık profesyonellerinin uygulama becerilerini artırdığını göstermiştir. (CAN et al., 2021) Williams ve arkadaşlarının (2026) kapsamlı taraması, generatif yapay zekânın (GenAI) pediatrik hemşirelik eğitiminde senaryo tabanlı öğrenme, kişiselleştirilmiş geri bildirim ve değerlendirme desteği sağladığını ortaya koymuştur. (Williams et al., 2026) Ancak bu alandaki çalışmaların çoğunlukla Asya ve Orta Doğu merkezli olması, uygulamaların genellenebilirliği konusunda soru işaretleri yaratmaktadır.

Eğitim Materyali Geliştirme: Generatif YZ, hasta iletişimini destekleme ve bireyselleştirilmiş eğitim materyalleri oluşturma gibi işlevlerde kullanılabilmektedir. (Watson et al., 2025)

3. Yönetim ve Kaynak Optimizasyonu

YZ, sağlık kurumlarının yönetimsel süreçlerinde verimliliği artırmaktadır.

İş Akışı İyileştirme ve Personel Planlaması: Araştırmalar, YZ'nin sağlık hizmetlerinde kaynak tahsisini daha etkin hâle getirdiğini ve hemşirelik iş akışının iyileştirilmesine katkı sağladığını göstermektedir. (Hassanein et al., 2025; Martinez-Ortigosa et al., 2023; Y. Park et al., 2025) Özellikle personel planlamasının optimize edilmesi ve idari iş yükünün azaltılması gibi alanlarda YZ uygulamalarının önemli avantajlar sunduğu bildirilmektedir.

4. Halk Sağlığı Uygulamaları

YZ, özellikle ergen sağlığını ilgilendiren halk sağlığı sorunlarına yönelik yenilikçi çözümler sunmaktadır.

Siber Zorbalık Tespiti ve Çevrimiçi Risk İzleme: Sultan ve arkadaşlarının (2023) derlemesi, makine öğrenmesi tekniklerinin siber zorbalığın tespiti için geliştirildiğini ortaya koymaktadır. (Sultan et al., 2023) Sedgwick ve arkadaşlarının (2023) çalışması ise doğal dil işleme yöntemleriyle elektronik sağlık kayıtlarından çevrimiçi etkinliklerin çıkarılmasının mümkün

olduğunu göstermiştir. (Sedgwick et al., 2023) Bu yaklaşımlar, ergen ruh sağlığı takibinde yeni fırsatlar sunmaktadır.

Tablo 3. YZ Uygulamalarının Sağladığı Fırsatlar ve Beraberindeki Etik Zorluklar / Riskler

Fırsatlar	Etik Zorluklar / Riskler
Artan Tanısal Doğruluk: Hastalıkların ve komplikasyonların daha erken ve doğru tespiti.	**Veri Gizliliği ve Güvenliği:** Hassas hasta verilerinin izinsiz erişim ve ihlallere karşı korunmaması.
Kişiselleştirilmiş Bakım: Hastanın genetik, gelişimsel ve çevresel verilerine göre özelleştirilmiş tedavi ve bakım planları.	**Algoritmik Önyargı:** Eğitim verilerindeki önyargılar nedeniyle bazı hasta gruplarının dezavantajlı duruma düşmesi ve sağlık eşitsizliklerinin artması.
Azalan İdari Yük: Otomasyon sayesinde hemşirelerin evrak ve veri girişi gibi işlere ayırdığı sürenin azalması, hasta başında geçen sürenin artması.	**Şeffaflık ve Hesap Verebilirlik (Kara Kutu Problemi):** Algoritmaların karar süreçlerinin anlaşılmaz olması ve olası hatalarda sorumluluğun belirlenememesi.
Eğitimde Yenilik: Simülasyonlar ve kişiselleştirilmiş eğitim ile hemşirelerin ve öğrencilerin becerilerinin geliştirilmesi.	**Hasta/Ebeveyn Katılımının Eksikliği:** Teknolojinin geliştirilmesi sürecinde kullanıcıların (hasta, ebeveyn, hemşire) görüşlerinin alınmaması ve onay süreçlerinin yetersiz olması.
Kaynak Optimizasyonu: Personel planlaması ve iş akışının iyileştirilmesiyle sağlık sisteminin daha verimli çalışması.	**Dijital Uçurum:** Teknolojiye erişimi olmayan veya okuryazarlığı düşük grupların bu hizmetlerden faydalanamaması.
Halk Sağlığı İzlemi: Siber zorbalık ve diğer çevrimiçi risklerin tespitiyle toplum sağlığının korunması.	**Veri Toplama Kalitesi ve Standardizasyonu:** Eksik, hatalı veya standartlaşmamış verilerin YZ modellerinin performansını olumsuz etkilemesi.

Tartışma

Bu derleme, YZ'nin çocuk ve ergen hemşireliğinde giderek artan bir öneme sahip olduğunu ve klinik, eğitim, yönetim ve halk sağlığı gibi çok çeşitli alanlarda uygulama potansiyeli taşıdığını ortaya koymuştur. Bulgular, YZ'nin sağladığı fırsatlar kadar, beraberinde getirdiği etik zorlukların da altını çizmektedir. Bu bölümde, elde edilen bulgular mevcut literatür ışığında tartışılacak, alandaki güçlü ve zayıf yönler, literatür boşlukları ve metodolojik eksiklikler ele alınacaktır.

Fırsatlar ve Güçlü Yönler

Bu derlemenin en çarpıcı bulgularından biri, YZ'nin tanısal doğruluğu artırma ve bakımı kişiselleştirme potansiyelidir. Park ve arkadaşlarının (2025) ve Ganatra'nın (2025) çalışmaları, YZ modellerinin özellikle sepsis ve basınç yaraları gibi komplikasyonları tahmin etmede

yüksek performans gösterdiğini vurgulamaktadır. (Ganatra, 2025; T. Park et al., 2025) Bu durum, özellikle kritik bakım ünitelerinde hasta güvenliğini önemli ölçüde artırabilir. Ayrıca, KKDS'lerin hemşirelerin klinik karar verme süreçlerini desteklemesi, bakımın standardizasyonuna ve kanıta dayalı uygulamaların yaygınlaşmasına katkı sağlayabilir. (Lu et al., 2021; Zhao et al., 2022)

Bir diğer önemli fırsat, YZ'nin hemşirelerin idari yükünü azaltma ve iş akışını iyileştirme potansiyelidir. (Hassanein et al., 2025; Y. Park et al., 2025) Bu sayede hemşireler, doğrudan hasta bakımına daha fazla zaman ayırabilir ve bu da hasta memnuniyeti ve bakım kalitesinde artışa yol açabilir. Ayrıca, generatif YZ gibi yeni teknolojilerin eğitim materyali geliştirme ve hasta iletişimini destekleme gibi alanlarda sunduğu olanaklar, hemşirelik uygulamalarını zenginleştirme potansiyeli taşımaktadır. (Watson et al., 2025; Williams et al., 2026)

Etik Zorluklar ve Zayıf Yönler

YZ'nin sunduğu bu fırsatlar kadar, derlemede öne çıkan başlıca etik sorunlar da mevcuttur.

Veri Gizliliği ve Güvenliği: Özellikle hassas çocuk ve ergen verilerinin kullanımı, önemli gizlilik endişelerini beraberinde getirmektedir. Sisk ve arkadaşlarının (2020) çalışması, ebeveynlerin bu konudaki endişelerini açıkça ortaya koymaktadır.(Sisk et al., 2020) Sağlık kurumlarının, YZ sistemlerinde kullanılan verilerin anonimleştirilmesi ve güvenliğinin sağlanması için uluslararası standartlara (ör. GDPR, HIPAA) uygun politika ve altyapılar geliştirmesi zorunludur.

Algoritmik Önyargı ve Sağlık Eşitsizliği: Çeşitli ve temsili veri setleriyle eğitilmeyen YZ modelleri, mevcut sağlık eşitsizliklerini derinleştirebilir. Bu derlemede incelenen çalışmaların çoğunun belirli coğrafi bölgelerden (ör. Asya, Kuzey Amerika) gelmesi,(Williams et al., 2026) geliştirilen modellerin diğer popülasyonlara genellenebilirliği konusunda soru işaretleri yaratmaktadır. Algoritmik önyargıların tespiti ve giderilmesi için modellerin farklı demografik gruplar üzerinde test edilmesi ve sürekli izlenmesi gerekmektedir.

Şeffaflık ve Hesap Verebilirlik: Özellikle derin öğrenme gibi "kara kutu" modellerinin klinik kararlardaki rolü, şeffaflık ve hesap verebilirlik sorunlarını gündeme getirmektedir. Verhoeven ve arkadaşları (2026) ile Parsons ve arkadaşlarının (2025) vurguladığı gibi, açıklanabilir YZ (XAI) yaklaşımlarının benimsenmesi, klinisyenlerin ve hastaların sistemlere olan güvenini artırabilir.(Parsons et al., 2025; Verhoeven et al., 2026) Ancak XAI'nin klinik pratikte nasıl etkin bir şekilde kullanılacağına dair standartlar ve eğitimler henüz yeterince gelişmemiştir.

Hasta/Ebeveyn Katılımı: Çocuk ve ergenlerin kendilerinin ve ebeveynlerinin YZ'ye yönelik tutumları, bu teknolojilerin başarısı için kritiktir. Reinhart ve arkadaşlarının (2026) niteliksel çalışması, çocukların YZ'yi destekleyici bir araç olarak gördüğünü ancak nihai kararın sağlık profesyonellerinde olması gerektiğini vurgulamıştır.(Reinhart et al., 2026) Bu bulgu, YZ araçlarının geliştirilmesinde ve uygulanmasında hasta ve ebeveynlerin aktif katılımının önemini bir kez daha ortaya koymaktadır. Kullanıcı odaklı tasarım ve yaşa uygun bilgilendirme süreçleri hayati önem taşır.

Literatürdeki Boşluklar ve Metodolojik Eksiklikler

Derlemenin en önemli bulgularından biri, mevcut araştırmaların çoğunlukla algoritmik performansa (doğruluk, duyarlılık, seçicilik) odaklandığı, klinik uygulanabilirlik, gerçek yaşam doğrulama, maliyet-etkinlik ve uzun vadeli hasta sonuçları gibi konulara yeterince eğilmediğidir. Bir diğer önemli eksiklik, YZ uygulamalarının hemşirelik mesleği üzerindeki etkisini (iş yükü, iş memnuniyeti, tükenmişlik) inceleyen çalışmaların sınırlı olmasıdır. Ayrıca, çocuk ve ergen hemşireliğine özgü gelişimsel ve psikososyal faktörleri dikkate alan YZ modellerinin geliştirilmesi ve test edilmesi gerekmektedir

Gelecek Vizyonu: Klinik Entegrasyon, Etik ve Politika Önerileri

YZ'nin çocuk ve ergen hemşireliğindeki potansiyelinin gerçekleştirilebilmesi için multidisipliner bir yaklaşım ve kapsamlı bir stratejik planlama gerekmektedir.

Multidisipliner İş Birliğinin Optimizasyonu

YZ'nin etkin, güvenli ve adil bir şekilde uygulanabilmesi için disiplinler arası iş birliği şarttır. Suresh ve Misra (2025), klinisyenler ile bilgisayar bilimcilerinin ortak çalışmasının; model geliştirme, doğrulama, eğitim programları oluşturma ve etik standartlar geliştirme açısından kritik olduğunu vurgulamaktadır.(Suresh & Misra, 2025) Bu iş birliği sayesinde YZ araçları klinik yargıyı tamamlayıcı nitelikte geliştirilebilirken; hasta güvenliği, veri kalitesi ve adalet gibi temel ilkeler korunabilir. Multidisipliner ekiplerin oluşturulması ve bu ekipler arasında etkin iletişimin sağlanması için sağlık kurumlarının özel bütçe ve zaman ayırması gerekmektedir.

Etik ve Yasal Çerçevenin Güçlendirilmesi

Çocuk ve ergenlere yönelik YZ uygulamaları için özel etik kurallar ve yasal düzenlemeler geliştirilmelidir. Sezgin ve Boch'un (2026) önerdiği Pediatrik AI Hazırlık Çerçevesi (PAIR) gibi modeller, etik yönetişimden veri temsil yeteneğine kadar yedi alanda hazırlık kriterleri sunarak pediatrik YZ uygulamalarının sorumlu entegrasyonunu desteklemektedir.(Sezgin & Boch, 2026) Bu çerçeveler, şeffaflık, hesap verebilirlik, veri gizliliği ve algoritmik adaleti temel ilkeler olarak benimsemelidir. Ayrıca, YZ kaynaklı olası hatalarda sorumluluğun kimde olacağına dair yasal düzenlemelerin acilen yapılması gerekmektedir.

Eğitim ve Yeterlilik

Hemşirelik müfredatına YZ okuryazarlığı, veri bilimi ve etik konularını kapsayan dersler entegre edilmeli; mevcut sağlık profesyonellerine yönelik sürekli eğitim programları düzenlenmelidir. (Özsezer, 2022) Hemşirelerin YZ'nin yetenekleri, sınırlılıkları, potansiyel önyargıları ve etik sonuçları hakkında bilgi sahibi olması, bu teknolojilerin güvenli kullanımı ve hasta savunuculuğu açısından kritiktir (Ferrara & Topaz, 2026). Eğitimler, teorik bilginin yanı sıra, YZ araçlarının simüle edilmiş veya gerçek klinik ortamlarda kullanımını da içermelidir.(Ferrara & Topaz, 2026)

Sonuç ve Öneriler

Bu derleme, yapay zekâ teknolojilerinin çocuk ve ergen hemşireliğinde bakım kalitesini artırma ve sağlık hizmetlerini dönüştürme konusunda önemli bir potansiyele sahip olduğunu göstermektedir. YZ, klinik karar desteğinden eğitim ve yönetime kadar geniş bir yelpazede

yenilikçi çözümler sunmaktadır. Ancak bu potansiyelin güvenli, adil ve etik bir şekilde gerçekleştirilebilmesi için multidisipliner iş birliğinin güçlendirilmesi, kapsamlı etik ve yasal çerçevelerin oluşturulması ve hemşirelerin YZ okuryazarlığının artırılması zorunludur.

Aşağıdaki öneriler, bu hedeflere ulaşılmasına rehberlik edecektir:

Eğitim ve Yeterlilik: Hemşirelik lisans ve lisansüstü eğitim müfredatlarına YZ okuryazarlığı, veri bilimi, etik ve güvenlik konularını kapsayan zorunlu dersler eklenmelidir. Klinik hemşirelere yönelik uygulamalı ve sürekli eğitim programları geliştirilmelidir.

Etik ve Yasal Çerçeve: Çocuk ve ergenlere yönelik YZ uygulamaları için özel etik kurallar ve yasal düzenlemeler geliştirilmelidir. Bu düzenlemeler, şeffaflık, hesap verebilirlik, veri gizliliği, algoritmik adalet ve bilgilendirilmiş onam gibi temel ilkeleri kapsamalıdır.

Katılımcı Tasarım: YZ araçlarının gereksinim analizi, tasarımı ve değerlendirme süreçlerine çocuklar, ergenler, ebeveynler ve sağlık profesyonelleri aktif olarak dahil edilmelidir. Kullanıcı odaklı ve yaşa uygun tasarımlara öncelik verilmelidir.

Klinik Entegrasyon ve Değerlendirme: YZ sistemleri, mevcut klinik iş akışlarını aksatmayacak ve hemşirenin klinik yargısını destekleyecek şekilde entegre edilmelidir. Sistemlerin etkinliği, güvenliği ve maliyet etkinliği, gerçek klinik ortamlarda yapılan uzun vadeli çalışmalarla kanıtlanmalıdır.

Önyargı ve Eşitsizlik: Algoritmik önyargıların tespiti ve giderilmesi için çeşitli ve temsili veri setleri kullanılmalıdır. YZ uygulamalarının sağlık eşitsizliklerini derinleştirmediğinden emin olmak için sürekli izleme ve değerlendirme mekanizmaları kurulmalıdır.

Disiplinler Arası Araştırma: Algoritmik başarının ötesine geçerek klinik uygulanabilirlik, hasta deneyimi, sağlık eşitliği ve hemşirelik iş gücüne etki gibi konulara odaklanan, bilgisayar bilimcileri, klinisyenler, etik uzmanları ve sosyal bilimcileri bir araya getiren multidisipliner araştırmalar teşvik edilmelidir.

Derlemenin Sınırlılıkları

Bu derleme, belirli anahtar kelimeler ve veri tabanlarıyla sınırlıdır ve bu nedenle bazı ilgili çalışmaların gözden kaçmış olma ihtimali vardır. Geleneksel derleme yöntemi, sistematik derlemelerin aksine, seçim yanlılığına daha açıktır. Çalışmaların metodolojik kalitesi sistematik bir değerlendirme aracıyla (ör. PRISMA) değerlendirilmemiştir. Sadece İngilizce ve Türkçe yayınlarla sınırlı olmak, dil yanlılığına neden olabilir. Ayrıca, YZ alanındaki hızlı gelişmeler, bu derlemede yer alan çalışmaların kısa sürede güncelliğini yitirebileceği anlamına gelmektedir.

KAYNAKLAR

Alowais, S. A., Alghamdi, S. S., Alsuhebany, N., Alqahtani, T., Alshaya, A. I., Almohareb, S. N., Aldairem, A., Alrashed, M., Bin Saleh, K., Badreldin, H. A., Al Yami, M. S., Al Harbi, S., & Albekairy, A. M. (2023). Revolutionizing healthcare: the role of artificial intelligence in clinical practice. *BMC Medical Education*, *23*(1), 689. https://doi.org/10.1186/s12909-023-04698-z

Alqaraleh, M., Almagharbeh, W. T., & Ahmad, M. W. (2025). Exploring the impact of artificial intelligence integration on medication error reduction: A nursing perspective. *Nurse Education in Practice*, *86*, 104438. https://doi.org/10.1016/j.nepr.2025.104438

Can, B., Başer, A., Baktır Altuntaş, S., Özceylan, G., & Kolcu, G. (2021). Artıfıcıal Intellıgence In Health Educatıon. *SDÜ Tıp Fakültesi Dergisi*, *28*(2), 355–359. https://doi.org/10.17343/sdutfd.876439

Cellina, M., Cè, M., Alì, M., Irmici, G., Ibba, S., Caloro, E., Fazzini, D., Oliva, G., & Papa, S. (2023). Digital Twins: The New Frontier for Personalized Medicine? *Applied Sciences*, *13*(13), 7940. https://doi.org/10.3390/app13137940

Clausen, C., Leventhal, B., Nytrø, Ø., Koposov, R., Røst, T. B., Westbye, O. S., Koochakpour, K., Frodl, T., Stien, L., & Skokauskas, N. (2023). Usability of the IDDEAS prototype in child and adolescent mental health services: A qualitative study for clinical decision support system development. *Frontiers in Psychiatry*, *14*. https://doi.org/10.3389/fpsyt.2023.1033724

Cortese, S., Arango, C., Aymerich, C., Catalan, A., Chetouani, M., Cohen, D., Coghill, D., Gabellone, A., Iniesta, R., Kadan, A., Kerbage, H., Kessing, L. V., Margari, L., Matera, E., Marzulli, L., Mezinska, S., Mulraney, M., Nagy, P., Oliver, D., … Fusar-Poli, P. (2026). Child and adolescent psychiatry: challenges, solutions, opportunities, and future directions. *World Psychiatry*, *25*(2), 190–224. https://doi.org/10.1002/wps.70044

Ferrara, S. A., & Topaz, M. (2026). Artificial Intelligence in Pediatric Practice: Controversies, Policy, and Practical Pointers forNurse Practitioners. *Journal of Pediatric Health Care*, *40*(2), 308–317. https://doi.org/10.1016/j.pedhc.2025.12.014

Ganatra, H. A. (2025). Machine Learning in Pediatric Healthcare: Current Trends, Challenges, and Future Directions. *Journal of Clinical Medicine*, *14*(3), 807. https://doi.org/10.3390/jcm14030807

Hassanein, S., El Arab, R. A., Abdrbo, A., Abu-Mahfouz, M. S., Gaballah, M. K. F., Seweid, M. M., Almari, M., & Alzghoul, H. (2025). Artificial intelligence in nursing: an integrative review of clinical and operational impacts. *Frontiers in Digital Health*, *7*. https://doi.org/10.3389/fdgth.2025.1552372

Jang, W., & Lee, B. (2026). From scores to signals: evolution and innovations in pediatric early warning systems. *Acute and Critical Care*, *41*(2), 252–261. https://doi.org/10.4266/acc.004475

Kerth, J.-L., Hagemeister, M., Bischops, A. C., Reinhart, L., Dukart, J., Heinrichs, B., Eickhoff, S. B., & Meissner, T. (2024). Artificial intelligence in the care of children and adolescents with chronic diseases: a systematic review. *European Journal of Pediatrics*, *184*(1), 83. https://doi.org/10.1007/s00431-024-05846-3

Lee, Y., Hong, S., & Yon, D. K. (2025). Artificial intelligence in pediatric healthcare: bridging potential, clinical practice, and ethical considerations. *Clinical and Experimental Pediatrics*, *68*(9), 652–655. https://doi.org/10.3345/cep.2025.01844

Li, S., & Wang, H. (2018). Traditional Literature Review and Research Synthesis. In *The Palgrave Handbook of Applied Linguistics Research Methodology* (pp. 123–144). Palgrave Macmillan UK. https://doi.org/10.1057/978-1-137-59900-1_6

Lu, S.-C., Brown, R. J., & Michalowski, M. (2021). A Clinical Decision Support System Design Framework for Nursing Practice. *ACI Open*, *05*(02), e84–e93. https://doi.org/10.1055/s-0041-1736470

Martinez-Ortigosa, A., Martinez-Granados, A., Gil-Hernández, E., Rodriguez-Arrastia, M., Ropero-Padilla, C., & Roman, P. (2023). Applications of Artificial Intelligence in Nursing Care: A Systematic Review. *Journal of Nursing Management*, *2023*, 1–12. https://doi.org/10.1155/2023/3219127

Mastorci, F., Lazzeri, M. F. L., Vassalle, C., & Pingitore, A. (2024). The Transition from Childhood to Adolescence: Between Health and Vulnerability. *Children*, *11*(8), 989. https://doi.org/10.3390/children11080989

Özsezer, G. (2022). The future of artificial intelligence in nursing. *Journal of Human Sciences*, *19*(2), 285–299. https://doi.org/10.14687/jhs.v19i2.6217

Park, T., Lee, I.-H., Lee, S. W., & Kong, S. W. (2025). Artificial intelligence in pediatric healthcare: current applications, potential, and implementation considerations. *Clinical and Experimental Pediatrics*, *68*(9), 641–651. https://doi.org/10.3345/cep.2025.00962

Park, Y., Chang, S. J., & Kim, E. (2025). Artificial intelligence in critical care nursing: A scoping review. *Australian Critical Care*, *38*(4), 101225. https://doi.org/10.1016/j.aucc.2025.101225

Parsons, C. S., Zuiderwijk, A., Orchard, N. A., Oosterhoff, J. H. F., & de Reuver, M. (2025). Task-Technology Fit of Artificial Intelligence-based clinical decision support systems: a review of qualitative studies. *BMC Medical Informatics and Decision Making*, *25*(1), 397. https://doi.org/10.1186/s12911-025-03237-8

Reddy, S. M., Patel, S., Weyrich, M., Fenton, J., & Viswanathan, M. (2020). Comparison of a traditional systematic review approach with review-of-reviews and semi-automation as strategies to update the evidence. *Systematic Reviews*, *9*(1), 243. https://doi.org/10.1186/s13643-020-01450-2

Reinhart, L., Kerth, J.-L., Bischops, A. C., Hagemeister, M., Krassuski, L., Mayatepek, E., & Meissner, T. (2026). Children and adolescents‘ views on artificial intelligence in pediatric healthcare: a qualitative focus group study. *BMC Pediatrics*, *26*(1), 563. https://doi.org/10.1186/s12887-026-07121-w

Sedgwick, R., Bittar, A., Kalsi, H., Barack, T., Downs, J., & Dutta, R. (2023). Investigating online activity in UK adolescent mental health patients: a feasibility study using a natural language processing approach for electronic health records. *BMJ Open*, *13*(5), e061640. https://doi.org/10.1136/bmjopen-2022-061640

Sezgin, E., & Boch, S. (2026). The pediatric AI readiness framework: bridging evidence to practice in pediatric artificial intelligence. *Frontiers in Artificial Intelligence*, *9*. https://doi.org/10.3389/frai.2026.1800047

Shen, M. (2026). Autistic Children’s Emotional Behavior Warning System Based on Smart Wearable Devices. *Engineering Reports*, *8*(3). https://doi.org/10.1002/eng2.70645

Sisk, B. A., Antes, A. L., Burrous, S., & DuBois, J. M. (2020). Parental Attitudes toward Artificial Intelligence-Driven Precision Medicine Technologies in Pediatric Healthcare. *Children*, *7*(9), 145. https://doi.org/10.3390/children7090145

Sultan, D., Omarov, B., Kozhamkulova, Z., Kazbekova, G., Alimzhanova, L., Dautbayeva, A., Zholdassov, Y., & Abdrakhmanov, R. (2023). A Review of Machine Learning Techniques in Cyberbullying Detection. *Computers, Materials & Continua*, *74*(3), 5625–5640. https://doi.org/10.32604/cmc.2023.033682

Suresh, S., & Misra, S. M. (2025). A Review of the Role of Artificial Intelligence in Pediatric Clinical Care, Education, and Research. *Current Pediatrics Reports*, *13*(1), 32. https://doi.org/10.1007/s40124-025-00362-w

Verhoeven, R., Bouisaghouane, W., & Hulscher, J. B. (2026). Explainable AI: Ethical Frameworks, Bias, and the Necessity for Benchmarks. *European Journal of Pediatric Surgery*, *36*(03), 168–173. https://doi.org/10.1055/a-2702-1843

Watson, A. L., Bond, C., Aveyard, H., Smith, G. D., & Jackson, D. (2025). Generative <scp>AI</scp> at the Bedside: An Integrative Review of Applications and Implications in Clinical Nursing Practice. *Journal of Clinical Nursing*. https://doi.org/10.1111/jocn.70151

Williams, S., Dabkowski, E., & Webb, C. (2026). Generative artificial intelligence in undergraduate paediatric nursing education: A scoping review of current applications and ethical considerations. *Nurse Education Today*, *164*, 107157. https://doi.org/10.1016/j.nedt.2026.107157

Yan, A. P., Guo, L. L., Inoue, J., Arciniegas, S. E., Vettese, E., Wolochacz, A., Crellin-Parsons, N., Purves, B., Wallace, S., Patel, A., Roshdi, M., Jessa, K., Cardiff, B., & Sung, L. (2025). A roadmap to implementing machine learning in healthcare: from concept to practice. *Frontiers in Digital Health*, *7*. https://doi.org/10.3389/fdgth.2025.1462751

Yang, X., Li, T., Su, Q., Liu, Y., Kang, C., Lyu, Y., Zhao, L., Nie, Y., & Pan, Y. (2025). Application of large language models in disease diagnosis and treatment. *Chinese Medical Journal*, *138*(2), 130–142. https://doi.org/10.1097/CM9.0000000000003456

Zhao, Y., Hu, J., Gu, Y., Wan, Y., Liu, F., Ye, C., & Zhang, X. (2022). Development and Implementation of a Pediatric Nursing-Clinical Decision Support System for Hyperthermia. *CIN: Computers, Informatics, Nursing*, *40*(2), 131–137. https://doi.org/10.1097/CIN.0000000000000812

KUR'ÂN'DA SIRAT KAVRAMININ ANLAM KATMANLARI
LAYERS OF MEANİNG OF THE CONCEPT OF SİRAT İN THE QUR'AN

Büşra ERDEN
Master's student Büşra Erden, Institute of Social Sciences, Hacı Bektaş Veli University, Nevşehir, Turkey

ÖZET

Kur'ân-ı Kerîm'in temel kavramlarından biri olan sırat, insanı dünya ve ahiret saadetine ulaştıran ilâhî rehberliği ifade eden önemli bir kavramdır. Kur'ân'da farklı bağlamlarda kullanılan bu kavram, yalnızca "yol" anlamına gelmemekte; hidayet, istikamet, kulluk, iman ve ahlâk ekseninde şekillenen kapsamlı bir anlam dünyasına işaret etmektedir. Bu çalışma, Kur'ân'da geçen sırat kavramının anlam katmanlarını ortaya koymayı ve kavramın semantik zenginliğini tespit etmeyi amaçlamaktadır.

Araştırmada öncelikle sırat kavramının sözlük ve terim anlamları incelenmiş, ardından Kur'ân'da geçtiği ayetler bağlam merkezli bir yaklaşımla değerlendirilmiştir. Bu çerçevede sıratın en sık kullanıldığı şekil olan "sırât-ı müstakîm" ifadesi üzerinde durulmuş; kavramın doğruluk, istikamet, denge ve ilâhî rehberlik gibi temel anlam boyutları ele alınmıştır. Ayrıca sırat kavramının sebîl, tarîk, hidâyet ve dalâlet gibi yakın anlamlı kavramlarla ilişkisi incelenerek aralarındaki benzerlik ve farklılıklar ortaya konulmuştur.

Çalışmada, Kur'ân'da sırat kavramının daima tekil olarak kullanılmasının, hakikate ulaştıran yolun özde bir ve tek olduğuna işaret ettiği değerlendirilmiştir. Bunun yanında sıratın yalnızca ahiret hayatında karşılaşılacak bir köprü olarak değil, insanın inanç, ibadet, ahlâk ve sosyal hayatını düzenleyen ilâhî bir yaşam modeli olarak sunulduğu görülmüştür. Bu yönüyle sırat, müminin hayatını Kur'ân ve sünnet doğrultusunda şekillendiren temel bir ilke olarak karşımıza çıkmaktadır.

Sonuç olarak çalışma, sırat kavramının Kur'ân'da çok katmanlı bir anlam yapısına sahip olduğunu ve bu kavramın ilâhî hidayetin, istikametin ve kurtuluşun sembolü olarak merkezi bir konumda bulunduğunu ortaya koymaktadır.

Anahtar Kelimeler: Tefsir, Kur'ân-ı Kerîm, Sırat, Sırât-ı Müstakîm, Hidâyet, Semantik.

ABSTRACT

One of the fundamental concepts in the Qur'an, ṣirāṭ (the straight path), represents the divine guidance that leads humanity to prosperity in both this world and the Hereafter. Used in various contexts throughout the Qur'an, this concept denotes not merely a "path" but also a comprehensive semantic framework centered on guidance, righteousness, servitude to God, faith, and morality. This study aims to examine the semantic dimensions of the concept of ṣirāṭ in the Qur'an and to reveal its rich conceptual structure.

The study first examines the lexical and terminological meanings of ṣirāṭ, followed by a contextual analysis of the Qur'anic verses in which the concept appears. Particular attention is given to the expression al-Ṣirāṭ al-Mustaqīm (the Straight Path), the most frequent form of the concept in the Qur'an, with emphasis on its principal semantic dimensions, including righteousness, steadfastness, balance, and divine guidance. Furthermore, the relationship between ṣirāṭ and related Qur'anic concepts such as sabīl (way), ṭarīq (path), hidāyah (guidance), and ḍalālah (misguidance) is examined in order to identify their similarities and differences.

The study concludes that the consistent use of ṣirāṭ in the singular form throughout the Qur'an signifies the essential unity of the path leading to the ultimate truth. It also demonstrates that ṣirāṭ is presented not merely as a bridge to be encountered in the Hereafter but as a comprehensive divine model of life that governs one's faith, worship, morality, and social conduct. In this respect, ṣirāṭ functions as a fundamental principle guiding the believer's life in accordance with the Qur'an and the Sunnah.

In conclusion, this study demonstrates that the concept of ṣirāṭ possesses a multilayered semantic structure in the Qur'an and occupies a central position as a symbol of divine guidance, righteousness, and salvation.

Keywords: Tafsir, Qur'an, Ṣirāṭ, al-Ṣirāṭ al-Mustaqīm, Guidance, Semantics.

INTRODUCTION

The Qur'an includes many subjects, ranging from matters of belief to matters of the unseen, from acts of worship to moral rules, and to the stories of the prophets from which a person should take advice and lessons. Although the Qur'an was revealed gradually over twenty-three years, it does not contain any contradictions, and it is remarkable that the surahs revealed at different times and in different places maintain a sense of unity.

The examination of these concepts mentioned in the Qur'an will help in a better understanding of the Qur'an. This is because a concept is used multiple times across various verses in the Qur'an. If the semantic differences between these concepts are not examined, that verse will not be properly understood.

The concept of *sirat* has many meanings such as path and passage. The concept of *sirat* is used together with the adjective *mustaqim*. It is used in the sense of the religion of Tawhid and the religion of Islam, which are in accordance with the consent of Allah and lead to Him: *"And whoever holds firmly to Allah has indeed been guided to a straight path (Sirat al-mustaqim)." "Indeed, Allah is my Lord and your Lord, so worship Him. That is the straight path (Sirat al-mustaqim)."*

The word *sirat* is also used with different adjectives in some verses. *"And they had been guided to good speech and guided to the path of the Praiseworthy (Hamid)."* In this verse, the word *sirat* is used with the attribute *Hamid*. *"Say, 'All are waiting; so wait. For you will know who the companions of the sound path (sawiyy) are and who is guided.'"* In this verse, however, the word *sirat* is characterized by *sawiyy*.

1. CONCEPTUAL FRAMEWORK

1.1. The Lexical Meaning of the Concept of Sirat

In the dictionary, the word *sirat* is derived from the root *s-r-t*, which means "to swallow," and it signifies "path" or "road." Its infinitive is *seret* or *seretan*. The origin of this word comes from the expression: "I swallowed the food." A path is called *sirat* based on the conception that it swallows the traveler, or that the traveler swallows it.

1.2. The Terminological Meaning of the Concept of Sirat

When the terminological meaning of the concept of *sirat* is mentioned, what comes to mind is the *sirat* in the Hereafter. Particularly in the sciences of creed (*akaid*) and kalam, it has gained the meaning of "a bridge erected over Hell, across which the people of Paradise will pass easily, while the people of Hell will slip and fall off."

The concept of *sirat* is used in the Holy Qur'an in the sense of a path. What is meant by this path is understood from the adjectives, words, or contexts used alongside the concept. This concept is used in the singular form in the Holy Qur'an; it has no plural form, meaning there is no usage like "*sirats*." This indicates that it is an unchanging and fixed path. It carries both a worldly and an otherworldly meaning: just as it refers to Allah's straight path in worldly life, it also refers to the "Bridge of Sirat" in the Hereafter.

In the Qur'an, the concept of *sirat* appears thirty-three times together with the adjective *Mustaqim*. *"Indeed, Allah is my Lord and your Lord, so worship Him. That is the straight path."* In this verse, the expression *sirat al-mustaqim* is used in the sense of the straight path shown by Allah, the religion of Islam.

In two verses, it is used in the sense of the "right path" together with the adjective *Sawiy*, meaning "straight" or "even." *"O my father! Indeed, there has come to me of knowledge that which has not come to you, so follow me; I will guide you to an even path."* The other verse is as follows: Say, *"All are waiting; so wait. For you will know who the companions of the sound path are and who is guided!"*

1.3. Concepts Related to Sirat

1.3.1. Sabil

The word *sabil* is derived from the root *s-b-l* and literally means path or route. Isfahani defines this word as an "unobstructed, straight path," and states that its plural form is *subul*. He has provided examples of where the word appears in the Qur'an and expressed that it means "the path of truth, the sound path."

The concept of *sabil* is used for anything that serves as a means to reach a goal, whether it is good or evil. It carries a more general meaning compared to the concept of *sirat* and simultaneously expresses all kinds of paths. It is a concept that encompasses both favorable and wicked meanings. For example, just as there is *"Sabilallah"* (the path of Allah), it can also be used in the sense of *"Sabil al-shaytan"* (the path of Satan). At the same time, the word *sabil* is a concept used to express individual paths.

"I will deprive of My signs those who unjustly puff themselves up on the earth; and [even] if they see every sign, they will not believe in it. And if they see the way of righteousness, they will not adopt it as a way; but if they see the way of error, they will adopt it as a way. That is because they have denied Our signs and have been heedless of them." The "way of righteousness" (*sabil al-rushd*) mentioned in the verse encompasses all virtues and beauties; the "way of error" (*sabil al-ghayy*), on the other hand, embraces all forms of deviation and evil, such as disbelief, idolatry, and hypocrisy.

"And He has cast into the earth firmly set mountains, lest it shift with you, and rivers and roads, that you may be guided." In this verse, the concept of *sabil* is used in its lexical meaning—that is, in the sense of a known path, whether long or short, walked upon by humans or animals.

"It is He who made the earth for you a bed and inserted therein for you roads and sent down from the sky, water, rain." We see that the concept of *sabil* is used in the sense of a known path in this verse as well. Likewise, Muqatil ibn Sulayman interpreted this verse as meaning roads on earth.

1.3.2. Tarik

It is an Arabic word; it is a noun derived from the verb *tereka*, and its plural is *turuk*. The word *tarik* has various meanings, which we can summarize as follows:

- a) A star rising, anything that comes at night, the morning star, the star with the sharpest light that pierces through the darkness.
- b) A path beaten by feet.
- c) Drinking murky water, the weakening of one's knees, a decline in intellect, a tendency toward play and amusement, layers of darkness descending, greed and avarice, setting traps and nets.

The word *tarik* is a concept that encompasses both material and spiritual paths. Additionally, the word *tarik* has both singular and plural forms, which implies the idea of multiple paths.

The word *tarik* appears 11 times in the Holy Qur'an.

"They said, 'O our people, indeed we have heard a Book revealed after Moses confirming what was before it, guiding to the truth and to a straight path [tarik].'" The word *tarik* in this verse is used in the sense of Allah's religion as the right path.

"They said, 'Indeed, these two are magicians who intend to drive you out of your land with their magic and to do away with your exemplary way [tarik].'" It is used in the sense of the path that is imitated and taken as an example.

"Indeed, those who disbelieve and commit wrong—never will Allah forgive them, Nor will He guide them to a path [tarik]—Except the path [tarik] of Hell; they will abide therein forever. And that, for Allah, is easy." In verse 168, the word *tarik* is used in the sense of the path of faith, while in verse 169, it is used in the sense of the path leading to Hell.

1.3.3. Hidayat

The word *hidayat* (*huda, hedy*) is used as a verbal noun (*masdar*) in the meanings of "going to the right path" and "pointing to the right path," and as a noun meaning "right path" and "guidance."

The concept of *hidayat* means a path, much like the concept of *sirat*. However, when the concept of *hidayat* is used as a noun, it means the "guide" or "leader" who shows the path. It demonstrates that while a human can invite others to the straight path, granting guidance belongs solely to Allah: *"Indeed, [O Muhammad], you do not guide whom you like, but Allah guides whom He wills. And He is most knowing of the [rightly] guided."*

1.3.4. Dalalat

Lexically, *dalalat* means "the absence of that which leads to the desired objective and entering a path that does not lead to the objective."

Dalalat means straying or deviating from the straight path and is used as the antonym of *hidayat*. In other words, *dalalat* is losing the path that leads to the desired thing.

The concept of *dalalat* is used in various meanings in the Holy Qur'an, which can be summarized as follows:

1. *"And did He not find you lost and guide you?"* In this verse, the word *dalalat* is used in the sense of making a mistake or being bewildered.
2. *"O David, indeed We have made you a successor upon the earth. So judge between the people in truth and do not follow your own desire, as it will lead you astray from the way of Allah. Indeed, those who go astray from the way of Allah will have a severe punishment for their forgetting the Day of Account."* The word *dal

2. The Moral and Practical Dimension of the Sirat (Straight Path)

Allah Almighty has always commanded His servants to remain steadfast, that is, to be upright as He has instructed. For this reason, He sent prophets as the finest examples for humanity to follow. For a person to remain on the straight path (istikāmah) means living upon the path of truth while avoiding all forms of excess and deviation. Allah Almighty says in the Qur'an:

"Indeed, those who say, 'Our Lord is Allah,' and then remain steadfast—the angels descend upon them, saying, 'Do not fear, nor grieve, but rejoice in the Paradise that you have been promised."

Being steadfast means turning away from everything other than Allah and directing oneself solely toward Him. When faced with hardships and calamities, a believer should not rebel against Allah but should instead remain patient, just as the prophets did.

For believers, Ṣirāṭ al-Mustaqīm (the Straight Path) means distancing oneself from all beliefs outside Islam and from everything that contradicts the Qur'an and the Sunnah, while walking only on the path of Allah. A believer should fulfill whatever Allah commands, even if it is difficult for the soul, seeking only His pleasure. One must adhere solely to the Qur'an and the Sunnah and continue steadfastly upon this path. Alongside acts of worship prescribed by Allah, moral virtues such as good character and patience should also occupy a central place in one's life.

A person should live with the awareness that every moment of life is under Allah's observation and that, ultimately, everyone will be held accountable. A believer should regard worldly life as a test. With every step taken, one should ask whether it is in accordance with Allah's pleasure and live with this consciousness. Rather than living randomly or carelessly, one should live a balanced and upright life.

Having the consciousness of the Straight Path does not mean allowing others' opinions or changing social trends to determine one's direction. A person should not abandon the right path

merely because it conflicts with personal interests. One should not display different identities at work, at home, or while performing acts of worship. Allah Almighty says:

"So remain steadfast as you have been commanded, together with those who have turned to Allah with you, and do not transgress. Indeed, He is All-Seeing of what you do."

As emphasized in this verse, the command to "remain steadfast as you have been commanded" should become the guiding principle of one's life.

Since **Ṣirāṭ al-Mustaqīm** is the path of the believers, the first essential requirement is faith. Allah Almighty says:

"As for those who believe in Allah and hold firmly to Him, He will admit them into His mercy and bounty and will guide them to Himself on a straight path."

In social life, the "straight path" refers to individuals living together in harmony. The principles that define this path include the following:

First, justice and fairness. The straight path requires a person to defend their own rights without violating the rights of others.

Second, empathy and tolerance. Diversity is an inevitable part of society. We should strive to understand those who think differently or have different lifestyles from our own. Even if we cannot fully understand them, we should respect their right to live and their way of life.

Third, honesty and trustworthiness. A society without honesty and trust cannot enjoy peace and stability. People naturally wish to distance themselves from such an environment.

Fourth, solidarity and mutual assistance. Human beings are inherently social. At times, no one can accomplish everything alone. In social life, the straight path means protecting the vulnerable, extending a helping hand to those in need, and sharing in both the joys and sorrows of others.

Fifth, a sense of responsibility. A person's responsibility is not limited to the home. Parks, streets, and traffic are also part of our shared living environment. Obeying rules, keeping the environment clean, and avoiding behavior that disturbs others in public spaces are all aspects of individual responsibility.

CONCLUSION

In conclusion, this study has examined the concept of *sirat*, which holds a significant place in the Holy Quran, within the framework of its lexical and terminological meanings, its usages in the Quran, its connections with related concepts, its reflections in the hadiths, and its theological dimensions. As a result of the evaluations made, it has been understood that the concept of *sirat* does not merely express the meaning of a "path," but rather represents a comprehensive divine guidance that encompasses both human worldly and hereafter lives.

Indeed, when the concept of *sirat* is reflected in daily life, it is directly related to fundamental values such as justice, honesty, trust, responsibility, security, cooperation, tolerance, mercy, and respect. Because walking on the straight path shown by Allah is not limited only to performing acts of worship, but also requires adhering to moral principles in human relations.

Therefore, *sirat* must be evaluated as a comprehensive lifestyle model that encompasses both individual and social life.

Ultimately, the concept of *sirat* is one of the most fundamental symbols of the path of guidance that the Quran offers to humanity. This concept signifies that a person organizes their faith, worship, morality, and social life in accordance with divine criteria. It is hoped that believers who live under the guidance of the Quran and the Sunnah, stay away from extremes, and strive to sustain their lives upon the *Sirat al-Mustaqim* will attain peace in worldly life and the mercy and pleasure of Allah in the hereafter. In this aspect, *sirat* is not merely a concept, but a divine direction and a path of salvation that guides a believer's life.

BIBLIOGRAPHY

el-İsfahânî Rağıb. *el-Müfredât Kur'ân Kavramları Sözlüğü.* İstanbul: Çıra Yayınları, Ağustos 2010.

Ağırakça Ahmed. *Şamil İslam Ansiklopedisi*. İstanbul: Şâmil Yayınevi, Temmuz 2000.

Süleymân Mukâtil bin. *Tefsîr-i Kebîr.* İstanbul: İşaret Yayınları, 2006.

Mustafa İbrahim. Hasan Ahmet. Abdulkadir Hamit. Neccar Muhammed Ali. *Mu'cemul-vasit* İstanbul: Çağrı Yayınları, 1989. tarik md.

Manzûr İbn. *Lisânü'l-ʿArab.* Beyrut: Dersaadet. tarik md.

Fîrûzâbâdî. *Kâmûsu'l-Muhît-Tercümesi*. İstanbul: Türkiye Yazma Eserler Kurumu Başkanlığı Yayınları, 2014.

Cürcani Ali İbn Muhammed es-Seyyid eş-Şerif. *Tarifat. Arapça- Türkçe Terimler Sözlüğü.* İstanbul: Bahar Yayınları, 1997.

Tehânevî. *Keşşâfü Istılâhâti'l-Fünun ve'l-Ulûm.* Lübnan: Mektebetu Lübnan Naşirun, 1996.

YABANCILARA TÜRKÇE ÖĞRETİMİNDE EDEBÎ METİN UYARLAMASI: AHMET HAMDİ TANPINAR'IN "FAL" HİKÂYESİNİN B1 DÜZEYİNE UYARLANMASI

LITERARY TEXT ADAPTATION IN TEACHING TURKISH TO FOREIGNERS: ADAPTING AHMET HAMDİ TANPINAR'S SHORT STORY "FAL" TO THE B1 LEVEL

Öğr. Gör. Havana KAYNAK
Türkçe Öğretimi Uygulama ve Araştırma Merkezi, Gümüşhane Üniversitesi, Gümüşhane, TÜRKİYE
ORCID ID: https://orcid.org/0009-0000-8607-7620

Yağmur ÜLKMEN
ORCID ID: https://orcid.org/0000-0001-7264-8031

ÖZET

Türkçenin yabancı dil olarak öğretiminde özgün edebî metinlerin kullanımı, öğrencilerin dil becerilerinin yanı sıra hedef kültürü tanımalarına da katkı sağlamaktadır. Özgün edebi metinlerin kullanımı, hedef kitlenin kültür düzeyini artıracağı gibi hedef dile ait düşünme ve algılama sürecini şekillendirmede de fayda sağlayacaktır. Ancak özgün metinlerin özellikle de klasik edebî metinlerin dil ve anlatım özellikleri, özellikle başlangıç ve orta düzey öğreniciler için çeşitli güçlükler oluşturabilmektedir. Bu bağlamda, özgün yazınsal metinlerin ders kitapları ile yardımcı kaynaklarda yer alabilmesi, bu metinlerin belirli ölçütler doğrultusunda hedef dil düzeylerine uygun şekilde uyarlanmasını gerekli kılmaktadır. Türk edebiyatının önemli eserleri, Türkçeyi yabancı dil olarak öğrenen bireylerin yararlanabilmesi amacıyla farklı dil yeterlik düzeylerine uyarlanarak yayımlanmaktadır. Bu eserler sayesinde öğreniciler yalnızca Türkçenin dilsel özelliklerini geliştirmekle kalmamakta, aynı zamanda Türk toplumunun kültürel değerlerini, yaşam biçimini ve düşünce dünyasını da tanıma fırsatı bulmaktadır. Bu bağlamda, *Avrupa Ortak Başvuru Metni (CEFR)* doğrultusunda gerçekleştirilen metin uyarlamaları, edebî eserlerin yabancı dil olarak Türkçe öğretiminde daha etkili ve işlevsel biçimde kullanılmasına katkı sağlamaktadır. Bu çalışmada, ders dışı okuma materyallerine duyulan gereksinime katkı sağlamak amacıyla, Türk edebiyatının önemli yazarlarından Ahmet Hamdi Tanpınar'ın *Fal* adlı hikâyesi, B1 düzeyindeki Türkçe öğrenicilerine yönelik olarak metin değiştirim ölçütleri çerçevesinde uyarlanmıştır. Çalışmanın temel amacı, uyarlanan metni Türkçeyi yabancı dil olarak öğrenen bireylerin okuma becerilerini geliştirmede kullanılabilecek yardımcı bir kaynak hâline getirmektir. Uyarlama sürecinde, B1 düzeyindeki öğrenicilerin kazanmış olması beklenen dil yapıları, *İstanbul Üniversitesi Yeni İstanbul Yabancılar İçin Türkçe B1 Ders Kitabı* ile *Diller İçin Avrupa Ortak Başvuru Metni (CEFR)* esas alınarak belirlenmiş; metin, bu ölçütler doğrultusunda yeniden düzenlenmiştir. Çalışmada nitel araştırma yöntemlerinden doküman incelemesi kullanılmış; metin, sözcük seçimi, cümle yapısı, metin bütünlüğü ve kültürel ögelerin sunumu bakımından incelenmiştir. Uyarlama sürecinde B1 düzeyi öğrenicilerinin dil yeterlikleri dikkate alınarak uzun ve karmaşık cümleler sadeleştirilmiş, kullanım sıklığı düşük

sözcükler daha yaygın karşılıklarıyla değiştirilmiş, anlamayı güçleştiren anlatım özellikleri yeniden yapılandırılmıştır. Bununla birlikte, hikâyenin edebî niteliğini ve kültürel dokusunu oluşturan temel unsurlar korunmaya özen gösterilmiştir. Çalışmanın sonucunda, *Fal* hikâyesinin uygun uyarlama ölçütleri doğrultusunda B1 düzeyinde kullanılabilecek bir okuma metnine dönüştürülebileceği öngörülmektedir. Araştırmanın, yabancılara Türkçe öğretiminde klasik Türk edebiyatı ürünlerinin ders materyali olarak kullanımına ilişkin çalışmalara katkı sağlaması ve edebî metin uyarlamalarına yönelik yeni araştırmalara örnek oluşturması beklenmektedir.

Anahtar Kelimeler: Yabancılara Türkçe öğretimi, metin uyarlama, Ahmet Hamdi Tanpınar, *Fal*.

ABSTRACT

The use of authentic literary texts in teaching Turkish as a foreign language contributes to students' understanding of the target culture alongside the development of their language skills. Utilizing authentic literary texts not only enhances the target audience's cultural awareness but also helps shape the thought processes and perceptions associated with the target language. However, the linguistic and stylistic characteristics of authentic texts—particularly classical literary works—can pose various challenges, especially for beginner and intermediate learners. Consequently, incorporating authentic literary texts into textbooks and supplementary materials necessitates adapting them to suit specific target language proficiency levels in accordance with established criteria. Significant works of Turkish literature are published in adapted versions tailored to different proficiency levels for individuals learning Turkish as a foreign language. Through these works, learners not only improve their command of the Turkish language but also gain insight into the cultural values, lifestyle, and worldview of Turkish society. In this context, text adaptations aligned with the Common European Framework of Reference for Languages (CEFR) facilitate a more effective and functional use of literary works in teaching Turkish as a foreign language. To address the need for supplementary reading materials, this study adapts the short story Fal (The Fortune) by Ahmet Hamdi Tanpınar—a prominent figure in Turkish literature—for B1-level Turkish learners, adhering to specific text adaptation criteria. The primary objective of the study is to transform the adapted text into a supplementary resource that can be used to enhance the reading skills of individuals learning Turkish as a foreign language. During the adaptation process, the language structures expected to be acquired by B1-level learners were determined based on the Istanbul University New Istanbul Turkish for Foreigners B1 Coursebook and the Common European Framework of Reference for Languages (CEFR), and the text was reorganized in accordance with these criteria. The study employed document analysis—a qualitative research method—to examine the text in terms of word choice, sentence structure, textual coherence, and the presentation of cultural elements. During the adaptation process, the language proficiency of B1-level learners was taken into account; consequently, long and complex sentences were simplified, low-frequency words were

replaced with more common equivalents, and stylistic features that hindered comprehension were restructured. At the same time, care was taken to preserve the fundamental elements that constitute the story's literary quality and cultural fabric. The study concludes that the story "Fal" can be transformed into a reading text suitable for the B1 level by adhering to appropriate adaptation criteria. It is anticipated that this research will contribute to the body of work regarding the use of classical Turkish literary pieces as instructional materials in teaching Turkish to foreigners and serve as a model for future research on literary text adaptation.

Keywords: Teaching Turkish to foreigners, text adaptation, Ahmet Hamdi Tanpınar, Fal.

GİRİŞ

Yabancılara Türkçe öğretiminde öğretici ile öğrenci arasındaki iletişimin sağlıklı ve etkili bir biçimde kurulmasında temel unsur dildir. Dil, yalnızca bireylerin duygu, düşünce ve ihtiyaçlarını ifade etmelerini sağlayan bir iletişim aracı değil, aynı zamanda toplumların sahip olduğu kültürel birikimin korunmasına ve kuşaktan kuşağa aktarılmasına imkân tanıyan önemli bir yapıdır. Bu nedenle dil, bireyler arasındaki iletişimin sağlanmasının yanı sıra toplumsal ve kültürel sürekliliğin oluşturulmasında da belirleyici bir role sahiptir. TDK[1] tarafından dil; insanlar arasında anlaşmayı ve iletişimi sağlayan, uzlaşmaya dayalı sesli veya görsel işaretler sistemi ile duygu ve düşüncelerin aktarılmasına yarayan bir anlatım aracı olarak tanımlanmaktadır. Bu tanım, dilin öncelikle iletişim kurma ve anlam aktarma işlevini öne çıkarmaktadır. Bununla birlikte Karataş (2001), dili yalnızca iletişim aracı olarak değil, aynı zamanda toplumu millet hâline getiren temel unsurlardan biri ve edebiyatın başlıca malzemesi olarak değerlendirmektedir. Karataş'a göre dil; bireylerin duygu ve düşüncelerini sözlü ya da yazılı biçimde başkalarına aktarmalarını sağlayan, aynı zamanda kültürel ve estetik değerlerin korunmasına aracılık eden canlı bir varlıktır. Bu tanımlardan hareketle dilin, iletişim kurma işlevinin ötesinde toplumsal kimliğin oluşumu, kültürel değerlerin aktarımı ve edebî ürünlerin ortaya konulması açısından da önemli bir işlev üstlendiği söylenebilir. Özellikle yabancılara Türkçe öğretimi bağlamında dil, öğrencilerin yalnızca dilsel bilgi ve beceriler kazanmalarını değil, aynı zamanda hedef dilin ait olduğu kültürel dünyayı tanımalarını da sağlamaktadır. Bu yönüyle yabancılara Türkçe öğretiminde dil öğretimi ve kültür aktarımı birbirini tamamlayan iki temel unsur olarak değerlendirilebilir.

İnsanlar tarih boyunca göç, ticaret, eğitim, turizm, diplomatik ilişkiler ve kültürel etkileşim gibi çeşitli nedenlerle yabancı dil öğrenme ihtiyacı duymuştur. Bu bağlamda Türkçe de geçmişten günümüze farklı toplumlar tarafından çeşitli amaçlarla öğrenilen diller arasında yer almıştır. Türkçenin yabancı dil olarak öğretimine yönelik çalışmaların tarihî geçmişinin oldukça eski dönemlere uzandığı görülmektedir. Nitekim Dîvânu Lugâti't-Türk, Muhâkemetü'l-Lugateyn ve Codex Cumanicus gibi eserler, Türkçenin farklı topluluklara tanıtılması ve öğretilmesi bakımından önemli kaynaklar arasında değerlendirilmektedir. Bu eserler, Türkçenin yabancı dil olarak öğretiminin tarihsel süreç içerisinde çeşitli ihtiyaçlara

[1] TDK (t.y.), https://sozluk.gov.tr

bağlı olarak geliştiğini göstermesi açısından önem taşımaktadır. Günümüzde ise Türkçenin yabancı dil olarak öğretimi, hem yurt içinde hem de yurt dışında kurumsal bir nitelik kazanarak yaygınlaşmıştır. Türkiye'de Türkçe öğretimi amacıyla faaliyet gösteren TÖMER'ler, yurt dışında ise Yunus Emre Enstitüsü başta olmak üzere farklı kurum ve kuruluşlar, Türkçenin yabancılara öğretimine yönelik çalışmalar yürütmektedir. Bu gelişmeler, Türkçeye yönelik ilginin arttığını ve Türkçenin yabancı dil olarak öğretiminin günümüzde giderek daha geniş bir alana yayıldığını göstermektedir.

Günümüzde yabancı dil öğretimi, yalnızca bireysel tercih ve uygulamalara dayalı bir süreç olmaktan çıkmış; uluslararası ölçekte kabul gören ortak standartlar ve ölçütler doğrultusunda yapılandırılmaya başlanmıştır. Bu bağlamda yabancı dil öğretimi, öğrenimi ve değerlendirilmesi süreçlerinde temel başvuru kaynaklarından biri olan Diller İçin Avrupa Ortak Başvuru Metni (Common European Framework of Reference for Languages), dil öğretim süreçlerinin planlanmasına, uygulanmasına ve değerlendirilmesine yönelik ortak bir çerçeve sunmaktadır. Diller İçin Avrupa Ortak Başvuru Metni, yabancı dil öğrenen bireylerin hedef dili etkili bir iletişim aracı olarak kullanabilmeleri için sahip olmaları gereken bilgi, beceri ve yeterlikleri belirli ölçütler çerçevesinde tanımlamaktadır. Bunun yanı sıra öğrenenlerin dil kullanımına ilişkin yeterliklerini farklı düzeylere ayırarak, dil öğrenme süreçlerinde kaydedilen ilerlemenin izlenmesine ve değerlendirilmesine imkân sağlamaktadır. Böylece dil öğretiminde ulaşılması hedeflenen kazanımların belirlenmesi ve öğretim sürecinin bu hedefler doğrultusunda yapılandırılması mümkün hâle gelmektedir. Bu yönüyle Diller İçin Avrupa Ortak Başvuru Metni, yabancı dil öğretiminde öğretim hedeflerinin belirlenmesi, öğrenme sürecinin düzenlenmesi ve öğrenenlerin dil yeterliklerinin ölçülüp değerlendirilmesi bakımından önemli bir başvuru kaynağı niteliğindedir. Ayrıca dil yeterliklerinin ortak ve uluslararası ölçütler doğrultusunda tanımlanması, farklı eğitim kurumları ve öğretim programları arasında belirli bir standartlaşmanın sağlanmasına da katkıda bulunmaktadır (CEF, 2001: 1).

Yabancılara Türkçe öğretiminde temel dil becerileri, genel olarak anlamaya ve anlatmaya dayalı beceriler olmak üzere iki grupta ele alınmaktadır. Anlamaya dayalı beceriler okuma ve dinleme; anlatmaya dayalı beceriler ise konuşma ve yazmadan oluşmaktadır. Bu dört temel dil becerisinin yanı sıra dil bilgisi öğretimi de yabancı dil öğretim sürecinin önemli bileşenlerinden biridir. Öğrencilerin öğrendikleri dil bilgisi yapılarını yalnızca kuramsal düzeyde bilmeleri değil, bu yapıları dört temel dil becerisi aracılığıyla kullanarak farklı iletişimsel bağlamlarda işlevsel hâle getirebilmeleri amaçlanmaktadır.

Yabancılara Türkçe öğretiminde dil yeterlikleri, Avrupa Ortak Başvuru Metni doğrultusunda A1, A2, B1, B2 ve C1 düzeyleri çerçevesinde yapılandırılmaktadır. Öğretim sürecinde dil bilgisi yapıları, kelime bilgisi ve dört temel dil becerisine yönelik kazanımlar, seviyelerin özellikleri dikkate alınarak basitten karmaşığa ve kolaydan zora doğru aşamalı bir biçimde sunulmaktadır. Her düzeyde öğrencilerin okuma, dinleme, konuşma ve yazma becerilerine ilişkin belirli kazanımlara ulaşmaları hedeflenmektedir. Öğrencilerin bulundukları düzeye ilişkin bilgi, beceri ve kazanımları edinmeleri, bir sonraki düzeye geçebilmeleri açısından

önem taşımaktadır. Dil öğretim sürecinde öğrencilerin dört temel dil becerisini geliştirmeye yönelik farklı yöntem, teknik ve uygulamalardan yararlanılmaktadır. Konuşma becerisinde öğrencilerin öğrendikleri kelime ve dil bilgisi yapılarını uygun bağlamlarda kullanarak sözlü iletişim kurmaları amaçlanırken, dinleme becerisinde seviyelerine uygun metinleri anlayarak dinlediklerine ilişkin çeşitli görevleri yerine getirmeleri beklenmektedir. Yazma becerisinde öğrencilerin, öğrendikleri kelime ve dil bilgisi yapılarını kullanarak belirli bir konu doğrultusunda anlamlı, tutarlı ve seviyelerine uygun metinler oluşturmaları hedeflenmektedir. Okuma becerisinde ise öğrencilerin seviyelerine uygun metinleri anlamaları, metinde yer alan açık ve örtük bilgileri değerlendirmeleri ve metne ilişkin çeşitli soruları yanıtlamaları yoluyla okuduğunu anlama becerilerinin geliştirilmesi amaçlanmaktadır.

Bu doğrultuda dört temel dil becerisi ile dil bilgisi öğretimi, birbirinden bağımsız alanlar olarak değil, öğrencinin hedef dili etkili, doğru ve işlevsel bir biçimde kullanabilmesini sağlayan bütüncül bir yapının unsurları olarak değerlendirilmelidir. Öğrencinin öğrendiği kelime ve dil bilgisi yapılarını okuma, dinleme, konuşma ve yazma süreçlerinde kullanabilmesi, dilsel yeterliğinin gelişmesi açısından büyük önem taşımaktadır. Bu nedenle yabancılara Türkçe öğretiminde öğretim materyallerinin ve öğrenme etkinliklerinin, öğrencilerin seviyelerine uygun biçimde hazırlanması ve dil becerilerinin birbirini destekleyecek şekilde geliştirilmesi gerekmektedir. Yabancılara Türkçe öğretiminde temel dil becerileri arasında yer alan okuma, öğrenicilerin hedef dili anlamlandırmalarına, dil bilgisel yapı ve sözcük dağarcıklarını geliştirmelerine önemli ölçüde katkı sunmaktadır. Okuma becerisinin etkili bir biçimde geliştirilebilmesi için öğrenicilerin dil düzeylerine uygun, anlaşılır ve aynı zamanda ilgilerini canlı tutabilecek metinlerle düzenli olarak çalışmalar yapmaları gerekmektedir. Bir metnin hangi amaçla kaleme alındığı, onun türsel özelliklerinin belirlenmesinde önemli bir ölçüttür. Yazar, iletmek istediği olay, duygu ya da düşünceyi, seçtiği metin türünün kendine özgü niteliklerinden yararlanarak okuyucuya aktarır (Erkul, 2004: 178).

Okuma etkinliklerinin gerçekleştirilmesinde kullanılan metinler, öğrenme sürecinin temel kaynakları arasında yer almaktadır. Bununla birlikte yabancı dil olarak Türkçe öğretiminde farklı dil düzeylerine hitap eden okuma kitaplarının yanı sıra edebî eserler, şiirler, atasözleri ve benzeri yardımcı materyallerin yeterli çeşitlilikte bulunmaması, okuma becerisinin geliştirilmesinde birtakım sınırlılıkların ve güçlüklerin ortaya çıkmasına yol açmaktadır. Bu nedenle, öğrenicilerin düzeylerine ve öğrenme ihtiyaçlarına uygun nitelikli metinlerin hazırlanması ve öğretim sürecine dâhil edilmesi önem taşımaktadır. Metin, başlangıç ve bitiş sınırları bulunan ve dilsel göstergelerin birbirleriyle ilişkili biçimde bir araya gelmesiyle oluşan anlamlı bir bütündür (Günay, 2007: 44). Her metin, belirli bir amaç doğrultusunda oluşturulmakta ve belirli bir konu etrafında şekillenmektedir. Bununla birlikte özellikle özgün edebî metinlerde yer alan kelime ve kelime gruplarının çeşitliliği, kullanılan dil bilgisi yapılarının karmaşıklığı, cümlelerin uzunluğu ve kültürel ögelerin yoğunluğu, bu metinlerin yabancı dil öğrenicilerinin mevcut dil düzeyleriyle her zaman örtüşmemesine neden olabilmektedir. Bu durum, yabancılara Türkçe öğretiminde öğrencilerin seviyelerine uygun

okuma materyallerinin hazırlanmasını gerekli kılmaktadır. Bu doğrultuda, özgün metinlerin farklı dil düzeylerindeki öğreniciler tarafından kullanılabilmesini sağlamak amacıyla metin uyarlama çalışmalarından yararlanılmaktadır. Metin uyarlama, özgün metnin temel içeriği ve anlam bütünlüğü mümkün olduğunca korunarak metinde yer alan dilsel unsurların hedef öğrenici grubunun seviyesine göre yeniden düzenlenmesini ifade etmektedir. Bu süreçte öğrencilerin dil düzeylerinin üzerinde bulunan sözcükler, dil bilgisi yapıları, cümle kuruluşları ve söz dizimsel özellikler daha anlaşılır bir biçimde yeniden yapılandırılabilmektedir. Böylelikle özgün edebî metinlerin yabancı dil olarak Türkçe öğrenenler için erişilebilir hâle getirilmesi ve bu metinlerden okuma becerisinin geliştirilmesine yönelik daha etkin bir şekilde yararlanılması hedeflenmektedir.

Bu çalışmada, yabancılara Türkçe öğretiminde B1 düzeyindeki öğrencilerin okuma becerilerinin geliştirilmesine katkı sağlamak amacıyla bir edebî metin uyarlama çalışması gerçekleştirilmiştir. Çalışmanın inceleme ve uyarlama materyalini Ahmet Hamdi Tanpınar'ın Fal adlı hikâyesi oluşturmaktadır. Özgün hikâye; söz varlığı, dil bilgisi yapıları, cümle kuruluşları ve söz dizimsel özellikleri bakımından incelenmiş; metnin B1 düzeyindeki öğrencilerin dilsel yeterliklerine uygun hâle getirilmesi amaçlanmıştır. Uyarlama sürecinde B1 düzeyinde öğrenilmesi ve kullanılması beklenen dil bilgisi yapıları, söz varlığı ve dil becerilerine ilişkin kazanımlar dikkate alınmıştır. Bu doğrultuda öğrencilerin mevcut dil yeterliklerini aşabilecek kelime ve ifadeler, karmaşık dil bilgisi yapıları ve söz dizimsel özellikler belirlenerek hedef seviyeye uygun biçimde yeniden düzenlenmiştir. Bununla birlikte uyarlama sırasında hikâyenin temel konusu, ana düşüncesi, olay örgüsü ve anlatım bütünlüğünün mümkün olduğunca korunmasına özen gösterilmiştir. Böylece Tanpınar'ın özgün anlatımının ve edebî niteliğinin korunması ile B1 düzeyindeki öğrencilerin metni anlayabilmeleri arasında bir denge kurulmaya çalışılmıştır. Çalışmanın temel amacı, özgün bir Türk edebiyatı metninin B1 düzeyindeki yabancı öğrencilere yönelik olarak nasıl uyarlanabileceğini ortaya koymak ve edebî metinlerin yabancılara Türkçe öğretiminde kullanılabilirliğine katkı sağlamaktır. Bu doğrultuda Fal hikâyesinin B1 düzeyine uyarlanmasıyla öğrencilerin hem okuma becerilerinin geliştirilmesine hem de Türk dili ve edebiyatının kültürel unsurlarıyla karşılaşmalarına olanak sağlayacak bir materyal ortaya koymak hedeflenmiştir.

Araştırmanın Amacı Ve Önemi

Bu araştırmanın amacı, Türkçenin yabancı dil olarak öğretiminde kullanılabilecek özgün edebî metinlerin dil düzeylerine uygun biçimde uyarlanmasına yönelik uygulamalı bir örnek ortaya koymak ve metin uyarlama sürecine ilişkin sistematik bir yaklaşım geliştirilmesine katkı sağlamaktır. Bu doğrultuda Ahmet Hamdi Tanpınar'ın Fal adlı hikâyesi, Avrupa Ortak Başvuru Metni (CEFR) esas alınarak B1 düzeyindeki Türkçe öğrenicilerinin dil yeterliklerine uygun biçimde uyarlanmıştır. Çalışmada, uyarlama sürecinde gerçekleştirilen sözcüksel, sözdizimsel, metinsel ve kültürel düzenlemelerin gerekçeleri ortaya konularak, özgün edebî metinlerin yabancı dil olarak Türkçe öğretiminde kullanılabilirliğinin değerlendirilmesi

amaçlanmaktadır. Bunun yanı sıra araştırma, Türk edebiyatının önemli eserlerinin yabancı dil olarak Türkçe öğretiminde daha etkin ve yaygın biçimde kullanılmasına katkı sunmayı hedeflemektedir.

Araştırmanın önemi, yabancı dil olarak Türkçe öğretiminde özgün edebî metinlerin hem dil becerilerinin geliştirilmesine hem de hedef kültürün tanıtılmasına önemli katkılar sağlamasına dayanmaktadır. Bununla birlikte, bu metinler sahip oldukları zengin söz varlığı, karmaşık cümle yapıları ve kültürel göndermeler nedeniyle özellikle orta düzey öğreniciler için doğrudan kullanılmaya her zaman uygun olmayabilmektedir. Bu nedenle, öğrenicilerin dil düzeylerine uygun biçimde gerçekleştirilen metin uyarlamaları, edebî eserlerin öğretim materyali olarak kullanılabilirliğini artırmaktadır.

Bu araştırma, Ahmet Hamdi Tanpınar'ın Fal adlı hikâyesini B1 düzeyine uyarlayarak, özgün bir edebî metnin yabancı dil olarak Türkçe öğretiminde nasıl öğretim materyaline dönüştürülebileceğine ilişkin uygulamalı bir model sunmaktadır. Çalışmanın, metin uyarlama sürecinde izlenen ölçütleri sistematik bir çerçevede ortaya koyması bakımından alana katkı sağlaması; öğreticilere, materyal geliştiricilere ve araştırmacılara yol gösterici olması beklenmektedir. Ayrıca araştırmanın, Türk edebiyatına ait özgün eserlerin yabancı dil olarak Türkçe öğretiminde kullanımını teşvik etmesi ve metin uyarlama alanında gerçekleştirilecek yeni araştırmalara örnek oluşturması hedeflenmektedir.

Araştırmanın Sınırlılıkları

Bu araştırma, Ahmet Hamdi Tanpınar'ın Fal adlı hikâyesinin yabancılara Türkçe öğretiminde B1 düzeyine uyarlanmasıyla sınırlandırılmıştır. Araştırma kapsamında yalnızca bu hikâye ele alınmış olup elde edilen bulgular farklı yazarların eserleri ya da farklı metin türleri için genellenemez. Uyarlama sürecinde, B1 düzeyindeki öğrenicilerin sahip olması beklenen dil yeterlikleri esas alınmıştır. Bu doğrultuda sözcük seçimi ve dil bilgisi yapılarının belirlenmesinde İstanbul Üniversitesi Yeni İstanbul Yabancılar İçin Türkçe ders kitaplarının A1, A2 ve B1 düzeyleri ile Diller İçin Avrupa Ortak Başvuru Metni (CEFR) temel alınmıştır.

Araştırma, metin uyarlama sürecinin ölçütleri doğrultusunda gerçekleştirilen doküman incelemesiyle sınırlıdır. Bu kapsamda hazırlanan uyarlanmış metnin B1 düzeyindeki Türkçe öğrenicileri üzerindeki etkililiği, okunabilirliği veya dil becerilerine katkısı uygulamalı olarak test edilmemiştir. Dolayısıyla çalışma, uyarlanan metnin öğretim ortamındaki etkililiğine ilişkin deneysel bulgular sunmamaktadır. Bununla birlikte, araştırmada belirlenen uyarlama ölçütlerinin ve izlenen sürecin, yabancılara Türkçe öğretiminde kullanılacak farklı edebî metinlerin uyarlanmasına yönelik benzer çalışmalara yöntemsel açıdan katkı sağlayacağı düşünülmektedir.

Ahmet Hamdi Tanpınar ve Hikâyeciliği

Ahmet Hamdi Tanpınar (1901–1962), Cumhuriyet dönemi Türk edebiyatının önde gelen şair, romancı, hikâyeci, denemeci ve edebiyat tarihçilerinden biridir. İstanbul'da doğan Tanpınar, çocukluk ve gençlik yıllarını babasının görevi nedeniyle Anadolu'nun farklı şehirlerinde

geçirmiştir. Bu deneyimler, eserlerinde önemli yer tutan zaman, bellek, mekân, yalnızlık ve insan psikolojisi gibi temaların şekillenmesinde etkili olmuştur. İstanbul Darülfununu Edebiyat Fakültesinde Yahya Kemal Beyatlı'nın öğrencisi olan Tanpınar, mezuniyetinin ardından çeşitli şehirlerde öğretmenlik yapmış; Ankara Gazi Terbiye Enstitüsü ile İstanbul Üniversitesi Edebiyat Fakültesinde görev almıştır. 1943–1946 yılları arasında milletvekilliği yapan yazar, daha sonra yeniden akademik çalışmalarına dönmüş ve yaşamının sonuna kadar edebiyat alanındaki çalışmalarını sürdürmüştür (Balcı, 2019).

Tanpınar, Doğu ile Batı kültürünü estetik bir anlayışla buluşturan özgün sanatçı kimliğiyle Türk edebiyatında önemli bir yere sahiptir. Eserlerinde bireyin iç dünyasını merkeze almış; zaman, rüya, bellek, yalnızlık, ölüm, aşk ve geçmişe özlem gibi temaları şiirsel bir anlatımla işlemiştir. Henri Bergson'un zaman anlayışından, Yahya Kemal Beyatlı'nın estetik görüşlerinden ve Fransız sembolist şairlerinden etkilenen yazar, modern anlatım tekniklerini geleneksel Türk kültürüyle birleştirmiştir. Şiirsel dili, güçlü tasvirleri ve psikolojik çözümlemeleri, onun edebî üslubunun belirgin özellikleri arasında yer almaktadır (Tanpınar, 1977; Okay, 2010).

Tanpınar, şiir, roman, hikâye, deneme, makale ve edebiyat tarihi alanlarında önemli eserler vermiştir:

Tevfik Fikret: Hayati, Şahsiyeti, Şiirleri ve Eserlerinden Seçmeler (1937); Namık Kemal Antolojisi (1942); Abdullah Efendi'nin Rüyaları (1943); Beş Şehir (1946); On Dokuzuncu Asır Türk Edebiyatı Tarihi (1949); Huzur (1949); Yaz Yağmuru (1955); Şiirler (1961); Saatleri Ayarlama Enstitüsü (1961); Yahya Kemal (1962); Edebiyat Üzerine Makaleler (1969); Yaşadığım Gibi (1970); Sahnenin Dışındakiler (1973); Tanpınar'ın Mektupları (1974); Mahur Beste (1975); Aydaki Kadın (1987); İki Ateş Arasında (1998); Edebiyat Dersleri (2002); Tanpınar'dan Ders Notları (2004); Günlüklerin Işığında Tanpınar'la Baş Başa (2007); Tanpınar'dan Notlar (2015); Hep Aynı Boşluk (2016); Tanpınar'dan Çeviriler (2017); Hüsrev ü Şirin (2017); Suat'ın Mektubu (2018); Alketis-Medeia-Elektra (Euripides'ten, 2018) (Tanpınar, 2022: 4).

Tanpınar'ın hikâyeciliği, modern Türk hikâyeciliğinin önemli örnekleri arasında değerlendirilmektedir. Hikâyelerinde olay örgüsünden çok karakterlerin iç dünyası, duygu ve düşünceleri ön plana çıkmaktadır. Bilinç akışı, iç konuşma, geriye dönüş ve sembolik anlatım gibi modern anlatım tekniklerinden yararlanan yazar, zaman ve mekânı karakterlerin psikolojik dünyasını destekleyen estetik unsurlar olarak kullanmıştır. Modernleşme, yalnızlık, kimlik arayışı, zaman, rüya, musiki ve bireyin varoluşsal sorunları hikâyelerinde sıkça işlediği temalar arasında yer almaktadır.

Tanpınar, hikâyelerini ilk olarak *Abdullah Efendi'nin Rüyaları* (1943) ve *Yaz Yağmuru* (1955) adlı iki kitapta yayımlamıştır. Bununla birlikte "yazarın sağlığında ayrı ayrı neşrettiği hikâyeleri uzun yıllardır, *Hikâyeler* adıyla tek kitap halinde yayımlanmasına rağmen *Emirgân'da Akşam Saati, Fal, Son Meclis* gibi bazı hikâyeler, Akün çalışmasında gösterildiği halde çok yakın zamanda kitaba dâhil edilmiştir" (Samsakçı, 2014:132). Dergâh Yayınları tarafından yayımlanan *Hikâyeler* adlı toplu basımda yer alan on dört hikâyede Tanpınar;

felsefe, tarih, musiki ve psikoloji gibi farklı alanlardan beslenen kültürel birikimini edebî ve estetik bir anlatım aracılığıyla yansıtmaktadır. Yazar, söz konusu hikâyelerinde bireyin iç dünyasına, psikolojik durumlarına ve gerçeklikle kurduğu ilişkiye odaklanırken, sahip olduğu çok yönlü kültürel birikimi anlatının çeşitli unsurlarıyla bütünleştirmektedir.
Bu araştırmanın inceleme konusu olan *Fal* adlı hikâye ise Tanpınar'ın bireyin iç dünyasını ve psikolojik çözümlemelerini ön plana çıkaran metinlerden biridir. Hikâyede merak, belirsizlik, kader, sezgi ve gelecek kaygısı gibi temalar, sembolik ve şiirsel bir anlatımla ele alınmaktadır. Zengin söz varlığı, kültürel unsurları ve estetik anlatımıyla dikkat çeken *Fal*, yabancı dil olarak Türkçe öğretiminde metin uyarlamasına elverişli nitelikler taşıması bakımından çalışmanın inceleme metni olarak tercih edilmiştir.

YÖNTEM

Araştırma Modeli

Bu araştırma, nitel araştırma yaklaşımı doğrultusunda şekillenmiştir. Araştırmada veri toplama yöntemi olarak doküman incelemesi, veri analizinde ise metin çözümleme tekniği kullanılmıştır. Doküman incelemesi, araştırma konusu ile ilgili yazılı kaynakların sistematik olarak incelenmesini ve yorumlanmasını esas alan nitel bir araştırma yöntemidir. Bu kapsamda araştırmanın temel veri kaynağını Ahmet Hamdi Tanpınar'ın Fal adlı hikâyesi oluşturmaktadır. Araştırma sürecinde öncelikle hikâye, sözcük seçimi, söz varlığı, cümle yapıları, metinsel bütünlük ve kültürel ögeler bakımından ayrıntılı olarak incelenmiştir. Ardından metin, yabancı dil olarak Türkçe öğretiminde B1 düzeyindeki öğrenicilerin dil yeterlikleri dikkate alınarak uyarlanmıştır. Uyarlama sürecinde, öğrenicilerin düzeyine uygun olmayan sözcükler daha yaygın kullanıma sahip karşılıklarıyla değiştirilmiş; uzun ve karmaşık cümle yapıları sadeleştirilmiş, metnin anlam bütünlüğünü bozmayacak şekilde gerekli düzenlemeler yapılmıştır. Bununla birlikte, metnin edebî niteliğini ve kültürel özelliklerini yansıtan temel unsurların korunmasına özen gösterilmiştir. Uyarlama sürecinde, yabancı dil olarak Türkçe öğretiminde yaygın olarak kullanılan İstanbul Üniversitesi Yeni İstanbul Yabancılar İçin Türkçe öğretim setlerinin A1, A2 ve B1 düzeylerindeki söz varlığı ve dil bilgisi kazanımları ile Diller İçin Avrupa Ortak Başvuru Metni (CEFR)'nde yer alan B1 düzeyi yeterlikleri ölçüt olarak esas alınmıştır. Bu doğrultuda gerçekleştirilen metin çözümlemesi sonucunda Fal adlı hikâye, B1 düzeyindeki Türkçe öğrenicilerinin dil yeterliklerine uygun biçimde yeniden düzenlenmiştir.

Evren ve Örneklem

Araştırmanın evrenini Ahmet Hamdi Tanpınar'ın hikâyeleri oluşturmaktadır. Araştırmanın örneklemi ise yazarın Hikâyeler adlı eserinde yer alan Fal adlı hikâyedir. Çalışmada örneklem olarak Fal hikâyesinin seçilmesinde, metnin Türk kültürüne özgü ögeler içermesi, zengin söz varlığına sahip olması ve yabancı dil olarak Türkçe öğretiminde metin uyarlamasına elverişli bir yapı sunması etkili olmuştur. Araştırmada incelenen Fal hikâyesi, Dergâh Yayınları tarafından yayımlanan Hikâyeler adlı eserden alınmıştır. Uyarlama süreci, söz konusu metin

esas alınarak gerçekleştirilmiş ve tüm çözümleme ile düzenlemeler bu baskı üzerinden yürütülmüştür.

Veri Toplama Araçları

Bu araştırmada veri toplama aracı olarak Ahmet Hamdi Tanpınar'ın Fal adlı hikâyesi ile metin uyarlama sürecine rehberlik eden kaynaklar kullanılmıştır. Araştırmanın temel veri kaynağını Fal adlı hikâye oluştururken, uyarlama ölçütlerinin belirlenmesinde Diller İçin Avrupa Ortak Başvuru Metni (CEFR)'nde yer alan B1 düzeyi dil yeterlikleri ve kazanımları esas alınmıştır. Bunun yanı sıra, sözcük seçimi, dil bilgisi yapıları ve dil düzeyinin belirlenmesinde İstanbul Üniversitesi Yeni İstanbul Yabancılar İçin Türkçe B1 Ders Kitabı temel başvuru kaynağı olarak kullanılmıştır. Bu kaynaklar doğrultusunda hikâye; söz varlığı, dil bilgisi yapıları, cümle uzunluğu, metinsel bütünlük ve kültürel ögeler bakımından incelenmiş, B1 düzeyindeki Türkçe öğrenicilerinin dil yeterliklerine uygun olacak şekilde uyarlanmıştır.

Verilerin Analizi

Bu araştırmada verilerin analizi beş aşamada gerçekleştirilmiştir. İlk aşamada Ahmet Hamdi Tanpınar'ın Fal adlı hikâyesi; söz varlığı, dil bilgisi yapıları, cümle uzunlukları, metinsel bütünlük ve kültürel ögeler bakımından ayrıntılı olarak incelenmiştir. Bu inceleme sonucunda metnin B1 düzeyindeki Türkçe öğrenicileri açısından güçlük oluşturabilecek dilsel ve yapısal özellikleri belirlenmiştir. İkinci aşamada, metnin uyarlanmasında esas alınacak ölçütler belirlenmiştir. Bu doğrultuda Diller İçin Avrupa Ortak Başvuru Metni (CEFR)'nde yer alan B1 düzeyi dil yeterlikleri ile İstanbul Üniversitesi Yeni İstanbul Yabancılar İçin Türkçe B1 Ders Kitabındaki söz varlığı ve dil bilgisi kazanımları incelenmiştir. Üçüncü aşamada, belirlenen uyarlama ölçütleri doğrultusunda metin; sözcüksel, sözdizimsel, metinsel ve kültürel düzeylerde yeniden düzenlenmiştir. Bu süreçte metnin anlam bütünlüğü ve edebî niteliğinin korunmasına özen gösterilmiş, yalnızca B1 düzeyindeki öğrenicilerin anlamasını güçleştiren unsurlar üzerinde gerekli uyarlamalar yapılmıştır. Dördüncü aşamada, uyarlanan metin yeniden incelenerek gerçekleştirilen düzenlemelerin B1 düzeyi kazanımlarına uygunluğu değerlendirilmiştir. Böylece uyarlama sürecinin hedef dil düzeyini karşılayıp karşılamadığı kontrol edilmiştir. Son aşamada ise uyarlanan metnin yabancı dil olarak Türkçe öğretiminde etkin bir biçimde kullanılabilmesini desteklemek amacıyla B1 düzeyine uygun okuma öncesi, okuma sırası ve okuma sonrası etkinlikler hazırlanmıştır. Bu etkinliklerle öğrenicilerin okuduğunu anlama, söz varlığını geliştirme, yorumlama ve üretim becerilerinin desteklenmesi amaçlanmıştır.

BULGULAR

Özgün Metnin (*Fal* Hikâyesi) Özellikleri

Edebî Özellikler

Ahmet Hamdi Tanpınar'ın hikâyeleri, *Abdullah Efendi'nin Rüyaları* (1943) ve *Yaz Yağmuru* (1955) adlı iki kitapta toplanmıştır. Tanpınar'ın şiirlerinde olduğu gibi hikâyelerinin sayısı da

oldukça sınırlıdır. *Abdullah Efendi'nin Rüyaları* adlı kitapta "Abdullah Efendi'nin Rüyaları", "Geçmiş Zaman Elbiseleri", "Bir Yol", "Erzurumlu Tahsin" ve "Evin Sahibi" olmak üzere beş hikâye yer almaktadır. 1955 yılında yayımlanan *Yaz Yağmuru* ise "Yaz Yağmuru", "Teslim", "Acıbadem'deki Köşk", "Rüyalar", "Âdem'le Havva", "Bir Tren Yolculuğu" ve "Yaz Gecesi" başlıklı yedi hikâyeden oluşmaktadır. Tanpınar'ın kitaplarına dâhil etmediği "Fal", "Emirgân'da Akşam Saati", "Son Meclis" ve "Bu Her Akşam Böyledir" adlı hikâyeler de dikkate alındığında, yazarın hikâye külliyatının toplam on altı metinden oluştuğu görülmektedir (Uçman, 2006:495). Dergâh Yayınları tarafından yayımlanan *Hikâyeler*[2] adlı toplu basımda *Fal* adlı hikâye de diğer metinlerle birlikte okura sunulmuştur. Bu eserde Tanpınar, felsefe, tarih, musiki ve psikolojiye dayanan zengin kültürel birikimini estetik bir anlatımla yansıtmıştır. *Fal,* Ahmet Hamdi Tanpınar'ın bireyin iç dünyasını ve psikolojik çözümlemelerini merkeze alan hikâyelerinden biridir. Hikâyede, gündelik yaşamın sıradan bir kesiti üzerinden bireyin merak duygusu, geleceği öğrenme isteği, belirsizlik karşısındaki psikolojik durumu ve insanın kaderle kurduğu ilişki ele alınmaktadır. Fal olgusu, hikâyede yalnızca geleceği haber veren geleneksel bir uygulama olarak değil; insanın bilinmezlik karşısındaki beklentilerini, korkularını ve umutlarını yansıtan sembolik bir unsur olarak kullanılmıştır. Tanpınar, olay örgüsünden çok karakterlerin ruh hâllerine ve iç dünyalarına odaklanarak psikolojik derinliği ön plana çıkarmıştır.

Dil ve anlatım bakımından *Fal,* Tanpınar'ın estetik anlayışını yansıtan özellikler taşımaktadır. Şiirsel anlatımı, zengin söz varlığı, ayrıntılı betimlemeleri ve sembolik anlatım unsurlarıyla dikkat çeken hikâyede uzun ve iç içe geçmiş cümleler, soyut ifadeler ve kültürel göndermeler önemli bir yer tutmaktadır. Bu özellikler metnin edebî değerini artırmakla birlikte, yabancı dil olarak Türkçe öğrenen orta düzey öğreniciler açısından anlaşılmasını güçleştirebilmektedir. Bu nedenle *Fal* hikâyesi, hem Türk kültürüne özgü unsurlar içermesi hem de söz varlığı, anlatım özellikleri ve kültürel zenginliği bakımından yabancı dil olarak Türkçe öğretiminde metin uyarlamasına uygun nitelikler taşımaktadır. Hikâyenin B1 düzeyine uyarlanmasıyla, metnin estetik ve kültürel özelliklerinin korunması hedeflenirken, öğrenicilerin dil düzeyine uygun, anlaşılabilir ve öğretim sürecinde kullanılabilir bir okuma metni oluşturulması amaçlanmıştır.

Teması: Fal hikâyesinin temel teması, insanın geleceği öğrenme arzusu ve belirsizlik karşısındaki psikolojik tutumudur. Tanpınar, fal olgusunu yalnızca geleneksel bir inanış olarak değil, insanın umutlarını, korkularını, beklentilerini ve bilinmeyene duyduğu ilgiyi yansıtan sembolik bir unsur olarak kullanır. Hikâyede ayrıca merak, kader, rastlantı, insan psikolojisi ve hayatın sıradan ayrıntılarının anlamlandırılması gibi temalar da işlenmektedir. Hikâye, insanların belirsizlik karşısında en sıradan ve tutarsız söylemleri bile kendi yaşamlarıyla ilişkilendirme eğilimini eleştirel ve ironik bir bakışla ortaya koymaktadır.

Olay Örgüsü: Hikâye, anlatıcının bir arkadaşıyla birlikte bir falcı kadının evine gitmesiyle başlar. Yoksul ve bakımsız bir evde geçen olaylarda, kahveler içildikten sonra falcı kadın

[2] Bu çalışmada incelenen "Fal" adlı hikâyeye ilişkin değerlendirmeler, Ahmet Hamdi Tanpınar'ın *Hikâyeler* (2011) adlı eserinde yer alan metin esas alınarak gerçekleştirilmiştir.

fincanları yorumlamaya başlar. Fal sırasında sürekli isimler, meslekler, hastalıklar, miras, aşk, para ve yolculuk gibi birbirinden farklı tahminlerde bulunur. Söylediklerinin çoğu belirsiz ve geneldir; buna rağmen konuşmasını sürekli yeni ayrıntılar ekleyerek sürdürür. Anlatıcı ve arkadaşı falcıyı sessizce dinler ancak falcı onların tepkilerinden hareketle tahminlerini değiştirmeye ve genişletmeye devam eder. Fal bittikten sonra iki arkadaş evden ayrılır. Hikâyenin sonunda arkadaşının, "Sana on dakikada herkesin romanını anlattı." sözüyle, falcının aslında insanların ortak beklenti, korku ve hayallerinden hareket ederek herkese uyabilecek bir hayat hikâyesi kurduğu vurgulanır. Böylece Tanpınar, fal üzerinden insan psikolojisini ve insanların bilinmeyene duyduğu ilgiyi ironik bir bakış açısıyla ele alır.

Karakterler: "Fal" hikâyesinde karakter sayısı sınırlıdır. Hikâyenin temel kişileri anlatıcı, onun arkadaşı ve falcı kadındır. Özellikle falcı kadın, sıra dışı görünüşü, davranışları ve konuşmalarıyla hikâyenin en dikkat çekici karakteridir. Fal sırasında adı geçen diğer kişiler ise gerçek birer karakterden çok, falcının hayal dünyasının ve belirsiz tahminlerinin bir parçası olarak karşımıza çıkar.

Anlatıcı: Hikâye, birinci tekil kişi anlatıcı tarafından aktarılmaktadır. Anlatıcı, arkadaşıyla birlikte falcı kadını ziyaret eder. Olayları gözlemleyen, falcı kadının davranışlarını ve içinde bulunulan ortamı ayrıntılı biçimde betimleyen kişidir. Anlatım boyunca eleştirel ve sorgulayıcı bir tutum sergiler.

Zaman: Hikâyede belirli bir tarih ya da zaman dilimi belirtilmemektedir. Olaylar, anlatıcının arkadaşıyla birlikte falcıya yaptığı kısa bir ziyaret süresince gerçekleşir. Zaman, kronolojik olarak ilerler ve birkaç saatlik bir zaman dilimini kapsar. Bu nedenle hikâyede belirsiz ancak dar zamanlı bir kurgu hâkimdir.

Mekân: Hikâyenin temel mekânı falcı kadının evidir. Tanpınar, bu mekânı eski, yoksul, bakımsız ve kasvetli bir atmosfer içinde ayrıntılı betimlemelerle anlatır. Odanın eski eşyaları, mangal, yıpranmış koltuklar, tozlu pencereler ve doldurulmuş papağan gibi ayrıntılar, hikâyenin atmosferini güçlendirir. Mekân, yalnızca olayların geçtiği bir yer değil; aynı zamanda falcı kadının yaşamını ve hikâyedeki gizemli havayı yansıtan estetik bir unsur olarak kullanılmıştır.

Anlatım tekniği: *Fal* hikâyesinde birinci tekil kişi (ben) anlatıcı kullanılmıştır. Anlatıcı, olayları kendi gözlem ve deneyimleri doğrultusunda aktarmaktadır. Hikâyede betimleme, diyalog ve gözlem teknikleri ön plandadır. Özellikle falcı kadın ile anlatıcı arasında geçen konuşmalar olay örgüsünü oluştururken, ayrıntılı mekân tasvirleri ve psikolojik gözlemler hikâyenin atmosferini güçlendirmektedir. Tanpınar, sembolik anlatım ve çağrışıma dayalı ifadelerle karakterlerin iç dünyasını yansıtarak estetik ve şiirsel bir anlatım ortaya koymuştur.

Dilbilgisel Özellikler

Özgün "Fal" hikâyesi, dil bilgisel özellikleri açısından analiz edildiğinde şu bulgular elde edilmiştir:

Kelime ve Cümle Sayısı: Özgün hikâye 149 cümle ve 1658 kelimeden oluşmaktadır. En uzun cümle 74 kelime, en kısa cümle ise 1 kelime içermektedir. Bulgular, hikâyenin cümle

uzunluğu açısından değişken ve bazı cümlelerin B1 seviyesi için fazla uzun olduğunu göstermektedir.

Kelime türleri dağılımı: Metinde yer alan kelimeler, isim soylu ve fiil soylu kelimeler olmak üzere iki temel kategori altında incelenmiştir. Yapılan inceleme sonucunda, çalışmaya konu olan hikâyede toplam 1.185 isim soylu ve 473 fiil soylu kelime bulunduğu belirlenmiştir.

Zaman kullanımı: Çalışmamıza konu olan metinde: belirli geçmiş zaman (62), geniş zaman (11), belirsiz geçmiş zamanın hikâyesi (10), belirsiz geçmiş zaman (5), şimdiki zamanın hikâyesi (19), geniş zamanın hikâyesi (2), şimdiki zaman (6), gelecek zaman(13),emir kipi (4), istek kipi (1), yeterlilik fiili (5) ve gereklilik kipi(1) kullanılmıştır.

B1 Seviyesine uygun olmayan yapılar: Çalışmamıza konu olan metinde B1 düzeyine uygun olmayan yapılar şunlardır:

- Geniş zamanın hikâyesine ait kullanımlar (gezdirdi, sarardı vb.)
- Ettirgen çatı kullanımları (oturttu vb.)
- Şimdiki zamanın rivayeti (biliyormuş)
- Geniş zamanın hikâyesi (bekleyemezdi, devam edebilirdi, olabilirdi)
- Sıfat-fiil yapıları (söndürülmüş, unutturan, sarkan, tıkırdayan, olabilecek vb.)
- Belirsiz geçmiş zamanın hikâyesine ait kullanımlar (pişmişti, yapışmıştı, olmuştu vb.),
- Ettirgen çatı kullanımları (gezdirdi, sarardı)
- Dönüşlü çatı kullanımları (mırıldandı, kımıldanmadık)
- İşteş çatı kullanımları (çarpıştı, tanıştın, buluştunuz, barışırsınız)
- Bağlaç kullanımları (yahut)
- Mecaz anlamlı kelimeler ve deyimler (savuştu, göz var, damar bulmuş, kaderin kapısında idi, karanlığa hücum etmek vb.)
- Doğrudan anlatım kullanımları (dedi, diyordu vb.)

UYARLAMA SÜRECİ

Metinlerin öğrenici düzeyine göre yeniden düzenlenmesi, yabancı dil öğretimi alanında farklı araştırmacılar tarafından çeşitli ölçütler doğrultusunda ele alınan bir konudur. Özellikle Nation (1993), Yano Y., Long M. H. ve Ross S. (1994), Alderson (2000) ve Chang (2008) gibi araştırmacılar, yabancı dil öğretiminde metinlerin öğrenici düzeyine uygun hâle getirilmesine yönelik farklı sadeleştirme çalışmaları yapmışlardır. Söz konusu araştırmalarda uyarlama işlemleri; metnin mevcut unsurlarının korunması, bazı bölümlerin değiştirilmesi, metnin özetlenmesi ve öğrenici düzeyi açısından gerekli görülmeyen unsurların metinden çıkarılması gibi temel uygulamalar çerçevesinde değerlendirilmiştir. Bunun yanı sıra, metinlerin anlaşılabilirliğini artırmak amacıyla sözcük seçimi ve söz dizimi üzerinde gerçekleştirilen düzenlemelere de yer verilmiştir. Bu alandaki çalışmalardan hareketle Bölükbaş (2015), "Yabancı Dil Olarak Türkçe Öğretiminde Okuma Metinlerinin Dil

Düzeylerine Göre Sadeleştirilmesi" başlıklı bildirisinde, yabancı dil olarak Türkçe öğretiminde kullanılan okuma metinlerinin öğrencilerin dil düzeylerine uygun biçimde sadeleştirilmesini sistematik bir çerçevede ele almıştır. Bölükbaş, , yabancı dil olarak Türkçe öğretiminde okuma metinlerinin sadeleştirilmesine yönelik uyarlama işlemlerinin koruma, değiştirme, özetleme ve silme olmak üzere dört temel strateji altında sınıflandırıldığını ifade etmiştir. Değiştirme stratejisi ise metnin öğrenici düzeyine uygun hâle getirilmesi amacıyla dilsel ve biçimsel unsurlar üzerinde gerçekleştirilen düzenlemeleri kapsamaktadır. Alan yazında bu strateji; sözcüksel, söz dizimsel, sözcüksel-söz dizimsel ve biçimsel değiştirme olmak üzere farklı alt kategoriler altında incelenmektedir (Alderson, 2000; Chang, 2008; Nation, 1993). Alan yazında yer alan bu sınıflandırmalar ve yapılan incelemeler doğrultusunda, bu çalışmada da metin uyarlama sürecinde değiştirme stratejisinden yararlanılmıştır. Bununla birlikte, mevcut sınıflandırmalarda yer alan alt kategoriler çalışmanın amacı ve incelenen metnin özellikleri doğrultusunda yeniden düzenlenmiştir. Bu kapsamda sözcüksel, söz dizimsel, sözcüksel-söz dizimsel ve biçimsel değiştirme kategorileri, sözcüksel değişiklikler ve söz dizimsel değişiklikler olmak üzere iki temel başlık altında değerlendirilmiştir. Ayrıca, B1 düzeyindeki öğrenicilerin dil bilgisel yeterlikleri dikkate alınarak dil bilgisel değişiklikler ile anlamlandırılması güç ve yapısal açıdan karmaşık cümlelerin daha anlaşılır hâle getirilmesi amacıyla karmaşık cümlelerin genişletilerek açıklanması başlıkları oluşturulmuştur. Belirlenen bu uyarlama ölçütleri doğrultusunda Ahmet Hamdi Tanpınar'ın Fal adlı hikâyesi, B1 dil düzeyindeki öğrenicilerin anlayabileceği ve takip edebileceği bir yapıya dönüştürülmüştür.

Sözcüksel Değişimler

Sözcüksel değişimler kapsamında, özgün metinde yer alan ve B1 seviyesine uygun olmayan sözcükler daha sık kullanılan ve öğrenicilerin düzeyine uygun karşılıklarıyla değiştirilmiştir. Ayrıca metindeki mecazlı ifadeler ve deyimler, B1 düzeyindeki öğrenicilerin anlayabileceği biçimde düzenlenmiş ve gerekli görülen durumlarda bu ifadelerin gerçek anlamlı kullanımları tercih edilmiştir. Bu tercihlere ait birkaç örnek aşağıdaki tabloda verilmiştir.

Sözcük Değişimleri	
Orijinal Metindeki Kullanım	Uyarlanmış Metindeki Kullanım
İskemle	Sandalye
Muzaffer	Başarılı
Manasız	Anlamsız
Hakiki	Gerçek
teşerrüf ettim	Memnun oldum

Mecaz ve Deyim Değişimleri	
Orijinal Metindeki Kullanım	Uyarlanmış Metindeki Kullanım
Gözleri parladı	Umutlandı/sevindi
Göz var	Nazar değiyor
Romanını anlattı	Hayat hikâyesini anlattı
Kaderin kapısında idi	Geleceğini bildiğini sanıyordu
Damar bulmuş	Uygun konu bulmuş

Söz Dizimsel Değişiklikler

Söz dizimsel değişiklikler kapsamında, özgün metindeki uzun ve karmaşık cümleler anlaşılırlığı artırmak amacıyla bölünerek B1 seviyesine uygun hâle getirilmiş, devrik ve kuralsız cümle yapıları ise kurallı cümlelere dönüştürülmüştür. Bu uyarlamalara ait birkaç örnek aşağıdaki gibidir.

Orijinal Cümle 1: Yerde bir bohça gibi katlanmış büyükçe kilim, yanı başında, kenarında birkaç mevsim evvelin sigaraları söndürülmüş, üzerindeki tencere kapağı ile yarı örtülü bir mangal, kenarda son derecede ince ayaklı ve arkalıklı narin, yaldızları baştan başa dökülmüş, yırtık astarının sarkan parçaları kumaşın yokluğunu unutturan iki iskemle, bir de bizi buyur ettikleri, bir ayağı kısa olduğu için her hareketimizde tıkırdayan, kendi aksaklığından gelen küçük imkânlarla bir kayık veya tahtarevalli gibi sallanan genişçe kavisli, arkalıklı bir kanepe vardı.

Uyarlanmış Cümle 1: Yerde büyük bir kilim vardı. Kilim katlı duruyordu. Yanında, üstü bir tencere kapağıyla yarı kapalı bir mangal duruyordu. Kenarda eski ve kumaşları yırtık iki sandalye vardı. Sandalyelerin boyaları da yer yer dökülmüştü. Bizi eski bir kanepeye oturttular. Kanepenin bir ayağı kısaydı. Bu yüzden kanepe hareket ettikçe ses çıkarıyor ve sallanıyordu

Orijinal Cümle 2: Pencereler dar, camlar sokağın biçareliğini göstermemeğe çalışır gibi tozlu ve çok ihtiyar yüzlerdeki çizgilere benzeyen ve acayip, derin kazılmış yağmur izleri ile örtülü idi.

Uyarlanmış Cümle 2: Pencereler dar ve camlar tozluydu. Üzerlerinde eski yağmur izleri vardı. Bu yüzden dışarıyı görmek zordu.

Orijinal Cümle 3: Bütün bu şeyler, odaya sanki bir sefalet tam barınabileceği yeri bulmuş ve hemen şu anda oracığa sığınmış hissini veriyordu.

Uyarlanmış Cümle 3: Bütün bunlar odayı çok bakımsız ve kötü gösteriyordu.

Dil Bilgisel Değişiklikler

Dil bilgisel değişiklikler kapsamında, B1 seviyesinin kazanımları dışında kalan dil bilgisi yapıları sadeleştirilmiş ve bunların yerine B1 düzeyine uygun yapılar kullanılmıştır.

Sıfat-fiil yapılarının uyarlanması: B1 düzeyine uygun olmayan sıfat-fiil yapıları, düzeye uygun yapılarla değiştirilmiştir.

Orijinal Cümle 1: Sadece kadının tepesinde, yarım tutam mısır püskülünü, kan kırmızı bir fiyango ile öbürlerinden ayıran ve kıpkırmızı satveti içinde eski Roma cengaverlerinin miğferlerini hatırlatan bez parçasının gidip gelişine bakıyordu.
Uyarlanmış Cümle 1: Sadece kadının başındaki kırmızı bez parçasına ve bu bez parçasının gidip gelişine bakıyordu.
Orijinal Cümle 2: O oturduğu zaman küt topuklarını daha iyi gördüm, ve elini sıkarken avucumda duyduğum terli yapışkanlıktan biraz daha iğrendim.
Uyarlanmış Cümle 2: O oturdu. Ben de kalın topuklarını daha iyi gördüm. Elini sıktım. Avucu terli ve yapış yapıştı. Bu yüzden kendimi daha kötü hissettim.

Birleşik zamanlı fiillerin uyarlanması: B1 düzeyine uygun olmayan birleşik zamanlı fiiller (geniş zamanın hikâyesi, gelecek zamanın hikâyesi ve belirsiz geçmiş zamanın hikâyesi) düzeye uygun yapılarla değiştirilmiştir.
Orijinal Cümle 1: On, on beş dakikadır, kapalı duran fincan, tabağına iyice yapışmıştı.
Uyarlanmış Cümle 1: Fincan on, on beş dakikadır kapalıydı. Bu nedenle tabağına iyice yapıştı.
Orijinal Cümle 2: Heyhat ki fincanın sahibi baba mirasının acılığını otuz sene evvel tanımıştı.
Uyarlanmış Cümle 2: Ne yazık ki fincanın sahibi, babasından kalan mirasın acısını otuz yıl önce tanıdı.
Fiilde çatı yapılarının uyarlanması: : B1 düzeyine uygun olmayan fiilde çatı yapıları (ettirgen çatı, dönüşlü çatı ve edilgen çatı) düzeye uygun yapılarla değiştirilmiştir.
Orijinal Cümle 1: Arkadaşının yüzü ister istemez sarardı; fakat kadın derhal tashih etti, bir yerden atlama.
Uyarlanmış Cümle 1: Arkadaşımın yüzünün rengi birden değişti. Ama kadın hemen sözünü düzeltti: Bir yerden atlama.
Orijinal Cümle 2: Bir handa buluştunuz, sonra gazinoya gittiniz, işte orada sizi gördüler.
Uyarlanmış Cümle 2: Bir otelde bir araya geldiniz. Sonra bir eğlence yerine gittiniz. İşte orada sizi gördüler.
Bağlaç kullanımları: B1 düzeyine uygun olmayan bağlaçlar, düzeye uygun yapılarla değiştirilmiştir.
Orijinal Cümle 1: Yahut ki terzidir?
Uyarlanmış Cümle 1: Yoksa terzi mi?
Doğrudan anlatım kullanımları: B1 düzeyine uygun olmayan doğrudan anlatım kullanımları, düzeye uygun dolaylı anlatım kullanımlarıyla değiştirilmiştir.
Orijinal Cümle 1: Bizi buraya sokan ihtiyar adam: "Madamın biraz işi var" dedi.
Uyarlanmış Cümle 1: Yaşlı adam bizi içeri aldı. Sonra, madamın biraz işi var diye belirtti.

Karmaşık Cümlelerin Genişletilerek Açıklanması

Metnin anlaşılırlığını artırmak amacıyla kısa cümlelerin genişletilmesi ve açıklanması ilkesine başvurulmuştur. Bu doğrultuda, özgün metinde az sözle ifade edilen ve B1 düzeyindeki öğrenicilerin anlamakta güçlük çekebileceği cümleler, anlamı açık ve anlaşılır kılacak biçimde genişletilerek yeniden düzenlenmiştir. Bu cümlelere ait birkaç örnek aşağıdaki gibidir.

Orijinal Cümle 1: Şurası muhakkak ki, sağ omuzuna asılı çok küçük kaplumbağa kabukları, çok kısa etekleri, mavi ceketiyle, acayip tavırlarıyla ilk görene bu zannını veriyordu.

Uyarlanmış Cümle 1: Sağ omzunda çok küçük kaplumbağa kabukları vardı. Kısa bir etek ve mavi bir ceket giyiyordu. Garip davranışları vardı. Bu yüzden ilk bakışta herkes onu deli sanıyor olabilirdi.

Orijinal Cümle 2: Tekrar mavi bakışlar üzerimize çevrildi; bizi bir yığın ıslak, kirli paçavraya sardı:

- Sende göz var oğul. ..

Uyarlanmış Cümle 2: Tekrar bize baktı. Bu bakışlar bizi rahatsız etti. Sende nazar var oğul diyerek düşündü.

Uyarlanmış Metnin Özellikleri

Kelime ve Cümle Sayısı: Fal hikâyesinin uyarlanmış metninde 249 cümle ve 1365 kelime yer almaktadır. Bu durum göz önünde bulundurulduğunda özgün metne oranla cümle sayısının ve kelime sayısının arttığı görülmektedir. Bu artış, özgün metnin B1 düzeyine uyarlanması sürecinde anlaşılması güç yapıların açıklayıcı ifadelerle genişletilmesi, deyim, atasözü ve kalıp ifadelerin daha anlaşılır biçimlere dönüştürülmesi, devrik ve eksiltili cümlelerin düzenlenmesi ve gerekli görülen yerlerde metne açıklayıcı ifadelerin eklenmesinden kaynaklanmaktadır.

Kelime türleri dağılımı: Uyarlanmış metinde 200 fiil soylu sözcük ve 1165 isim soylu sözcük bulunmaktadır. Bu durum göz önünde bulundurulduğunda özgün metne oranla fiil soylu ve isim soylu sözcüklerin arttığı görülmektedir.

Dilbilgisi Yapıları: Özgün metinde B1 düzeyinin üzerinde yer alan dilbilgisi yapıları ve sözcükler belirlenerek, bunların yerine Avrupa Dilleri Ortak Çerçeve Metni (CEFR) kapsamında A1, A2 ve B1 düzeyleri için belirlenen dil yeterlilikleri ve kazanımları ile İstanbul Yabancılar İçin Türkçe setindeki ilgili düzeylerin dilbilgisi yapıları ve sözcükleri esas alınmıştır.

Zaman kullanımı: Belirli geçmiş zaman (95), şimdiki zamanın hikâyesi (37), geniş zaman (20), belirsiz geçmiş zaman (9), gelecek zaman (13), şimdiki zaman (15), yeterlilik fiili (2), gereklilik kipi(1), istek kipi (1) ve emir kipi (4) kullanılmıştır.

SONUÇ

Bu çalışmada, Ahmet Hamdi Tanpınar'ın Fal adlı hikâyesinin yabancılara Türkçe öğretiminde B1 düzeyindeki öğrenicilerin okuma becerilerini destekleyecek biçimde uyarlanması amaçlanmıştır. Bu doğrultuda özgün metin; söz varlığı, dil bilgisi, kalıp ifadeler, deyimler, atasözleri ve söz dizimi bakımından incelenmiş ve B1 düzeyindeki öğrenicilerin dil yeterlikleri dikkate alınarak çeşitli uyarlama işlemlerine tabi tutulmuştur. Uyarlama sürecinde İstanbul Yabancılar İçin Türkçe A1, A2 ve B1 ders kitaplarından ve Avrupa Dilleri Ortak Çerçeve Metni'nde (CEFR) belirtilen B1 düzeyine ilişkin dil yeterliklerinden yararlanılmıştır. Böylece özgün metnin temel anlamı, anlatı bütünlüğü ve kültürel özellikleri mümkün olduğunca korunurken öğrenicilerin metni daha kolay anlamlandırabilmeleri hedeflenmiştir.

Çalışmada gerçekleştirilen uyarlamaların yalnızca kelime ve dil bilgisi düzeyinde gerçekleşmediği, aynı zamanda söz dizimsel ve anlamsal düzenlemeleri de kapsadığı görülmüştür. B1 düzeyindeki öğrenicilerin anlamlandırmakta zorlanabileceği uzun, karmaşık ve devrik cümleler yeniden yapılandırılmış; gerekli görülen durumlarda uzun cümleler daha kısa ve bağımsız cümlelere ayrılmıştır. Bunun yanı sıra bağlam içerisinde anlamlandırılması güç olan bazı ifadeler, açıklayıcı unsurlar eklenerek daha anlaşılır hâle getirilmiştir. Bu durum, edebî metin uyarlamasının yalnızca zor kelime ve dil bilgisi yapılarını daha basit karşılıklarla değiştirmekten ibaret olmadığını; metnin söz dizimsel ve anlamsal yapısının da öğrenici düzeyi doğrultusunda yeniden düzenlenmesini gerektirdiğini ortaya koymaktadır. Özgün metin ile uyarlanmış metnin karşılaştırılması sonucunda cümle ve sözcük sayılarında farklılıklar meydana geldiği tespit edilmiştir. Uyarlanmış metinde cümle sayısının yanı sıra isim soylu ve fiil soylu sözcüklerin sayısında da artış görülmüştür. Söz konusu artışın temel nedeninin, anlamın daha açık ve anlaşılır biçimde aktarılabilmesi amacıyla bazı ifadelerin açıklanması ve uzun cümlelerin birden fazla kısa cümleye dönüştürülmesi olduğu düşünülmektedir. Dolayısıyla metin uyarlama sürecinde metnin kısaltılmasının her durumda anlaşılabilirliği artırmadığı; aksine bazı durumlarda anlamın açık biçimde aktarılabilmesi için açıklayıcı unsurların eklenmesinin gerekli olabileceği sonucuna ulaşılmıştır. Çalışmanın önemli sonuçlarından biri de metinde yer alan kültürel ögelerin uyarlama sürecinde mümkün olduğunca korunmuş olmasıdır. Tanpınar'ın hikâyesinde yer alan fal, nazar, geleneksel inanışlar, toplumsal yaşam biçimleri ve dönemin kültürel özelliklerini yansıtan unsurlar, metnin kültürel bağlamını oluşturan temel ögeler olarak değerlendirilmiştir. Bu unsurların metinden tamamen çıkarılması yerine, B1 düzeyindeki öğrenicilerin anlayabileceği biçimde açıklanması veya bağlam içerisinde daha anlaşılır hâle getirilmesi tercih edilmiştir. Böylece metnin dilsel açıdan sadeleştirilmesi sırasında kültürel ve edebî değerlerinin kaybolmasının önüne geçilmeye çalışılmıştır. Bu yaklaşım, yabancılara Türkçe öğretiminde edebî metinlerin yalnızca dil öğretimine yönelik materyaller olarak değil, aynı zamanda hedef dilin kültürünü tanıtan araçlar olarak da değerlendirilmesi gerektiğini göstermektedir.

Sonuç olarak, özgün edebî metinlerin yabancılara Türkçe öğretiminde kullanılabilmesi için öğrenicilerin dil düzeyleri dikkate alınarak sistemli ve ölçütlere dayalı bir biçimde

uyarlanması gerekmektedir. Özellikle B1 düzeyindeki öğreniciler açısından söz varlığı, dil bilgisi ve söz dizimi bakımından gerçekleştirilen düzenlemelerin metnin anlaşılabilirliğini artırmada önemli olduğu görülmüştür. Bununla birlikte uyarlama sürecinde metnin edebî niteliğinin, temel anlamının, olay örgüsünün, anlatı bütünlüğünün ve kültürel unsurlarının mümkün olduğunca korunması gerektiği sonucuna ulaşılmıştır. Bu bağlamda edebî metin uyarlaması, yabancılara Türkçe öğretiminde hem dil becerilerinin geliştirilmesine hem de öğrenicilerin Türk edebiyatı ve kültürüyle tanışmasına katkı sağlayabilecek önemli bir uygulamadır. Fal hikâyesi üzerinde gerçekleştirilen bu çalışma, özgün bir edebî metnin B1 düzeyindeki öğrenicilerin dilsel ihtiyaçlarına uygun hâle getirilirken metnin kültürel ve edebî değerlerinin de korunabileceğini ortaya koymaktadır. Gelecekte yapılacak çalışmalarda farklı tür ve yazarlara ait edebî metinlerin farklı dil düzeylerine göre uyarlanması ve hazırlanan metinlerin öğreniciler üzerindeki etkisinin uygulamalı çalışmalarla değerlendirilmesi, yabancılara Türkçe öğretiminde kullanılabilecek nitelikli ve kültürel açıdan zengin materyallerin geliştirilmesine katkı sağlayacaktır.

KAYNAKÇA

Alderson, J.C. (2000). Assessing reading. Cambridge: Cambridge University Press.

Balcı, Y. (2019, 31 Aralık). Türk Edebiyatı İsimler Sözlüğü: Ahmet Hamdi Tanpınar maddesi. Erişim adresi: https://teis.yesevi.edu.tr. Erişim tarihi: 10.07.2026

Bölükbaş, F. (2015, Mayıs). Yabancı Dil Olarak Türkçe Öğretiminde Okuma Metinlerinin Sadeleştirilmesi. 1. Uluslararası Dil Eğitimi ve Öğretimi Sempozyumu, Nevşehir.

Bölükbaş, F., Keskin, F., & Gedik, E., & Tokgöz Utku, H., & Ünsal Rey, G., & Özenç, F., & Gönültaş, G., & Şimşek, İ. D., & Alaca, S., & Yusufoğlu, S. (2016). İstanbul Yabancılar İçin Türkçe Ders Kitabı A1. Kültür Sanat Basımevi.

Bölükbaş, F., Keskin, F., & Gedik, E., & Tokgöz Utku, H., & Ünsal Rey, G., & Özenç, F., & Gönültaş, G., & Şimşek, İ. D., & Alaca, S., & Yusufoğlu, S. (2016). İstanbul Yabancılar İçin Türkçe Ders Kitabı A2. Kültür Sanat Basımevi.

Bölükbaş, F., Keskin, F., & Gedik, E., & Tokgöz Utku, H., & Ünsal Rey, G., & Özenç, F., & Gönültaş, G., Göçen Özdemirel, G., & Yavuz, S., & Şimşek, İ. D., & Alaca, S., & Yusufoğlu, S. (2016). İstanbul Yabancılar İçin Türkçe Ders Kitabı B1. Kültür Sanat Basımevi.

Chang, S. (2008). A preliminary categorization of techniques of simplification in ESL/EFL graded readers. (Ed. Martin Edwardes) Proceedings of the BAAL Annual Conference. London: Scitsiugnil Press.

Common European Framework Of Reference For Language, Learning, Teaching, Assesment. (2001). Cambridge: Cambridge University Press.

Erkul, R. (2004). *Cümle ve Metin Bilgisi*. Ankara: Anı yayıncılık.

Günay, D. (2007). Liselerdeki Yazın Eğitimine Yeni Bir Yaklaşım. *Milli Eğitim*, 169, 8-17.

Karataş, T. (2001). Ansiklopedik Edebiyat Terimleri Sözlüğü. İstanbul: Perşembe Kitapları.

Nation, P. (1993). Measuring readiness for simplified material: a test of the first 1,000 words of English. In M. L. Tickoo (Ed.), Simplification: Theory and Application (pp. 193-203). Singapore: SEAMEO Regional Language Center.

Okay, M. O. (2010). Ahmet Hamdi Tanpınar. *İslâm Ansiklopedisi*. Türkiye Diyanet Vakfı Yay. 567- 570. (https://islamansiklopedisi.org.tr/tanpinar-ahmet-hamdi).

Samsakçı M. (2014). Tanpınar'ın Eşiğinde; Ahmet Hamdi Tanpınar ve Eserleri Üzerine Düşünceler. İstanbul: Kitabevi Yayınları.
Tanpınar A. H. (1977). Edebiyat Üzerine Makaleler. Haz. Zeynep Kerman. İstanbul: Dergâh Yay.
Tanpınar, A. H. (2011). *Hikâyeler*. Dergâh Yayınları
Tanpınar, Ahmet Hamdi (2022). Yaşadığım Gibi. İstanbul: Dergâh Yayınları.
TDK (t.y.), https://sozluk.gov.tr/ (Erişim tarihi: 10.07. 2026)
Uçman, A. (2006). Değişen Değerler Karşısında Ahmet Hamdi Tanpınar. Türkiye Araştırmaları Literatür Dergisi, (7), 479-509.

Ek: Uyarlanmış Metin *Fal*

Fal

Oda gerçekten çok kötü ve bakımsızdı. Yerde büyük bir kilim vardı. Kilim katlı duruyordu. Yanında, üstü bir tencere kapağıyla yarı kapalı bir mangal duruyordu. Kenarda eski ve kumaşları yırtık iki sandalye vardı. Sandalyelerin boyaları da yer yer dökülmüş. Eski bir kanepeye oturduk. Kanepenin bir ayağı kısaydı. Bu yüzden kanepe hareket ettikçe ses çıkarıyor ve sallanıyordu. Karşımızda, duvarda, bir papağan vardı. Bu papağan süs eşyasıydı. İki kollu bir dalın üzerindeydi. Bu eve ait değilmiş gibi görünüyordu. Pencereler dar ve camlar tozluydu. Üzerlerinde eski yağmur izleri vardı. Bu yüzden dışarıyı görmek zordu. Üstelik eski ve kırık perde çubukları vardı. Üzerlerinde eski perdeler ve tüller asılıydı. Bütün bunlar odayı çok bakımsız ve kötü gösteriyordu.

Yaşlı adam bizi içeri aldı. Sonra, madamın biraz işi var diye belirtti. "Ben kahvenizi yapacağım, o da şimdi gelir." diyerek yanımızdan ayrıldı. Gerçekten kadın gelmeden önce elinde temiz bir kahve tepsisi ile odaya girdi. İkimiz de fincanlara şüpheyle baktık. Fakat fincanlar temizdi. Zaten kahve de iyi pişmiş. Belli ki bu, işinin bir parçasıydı. Bu evde kahve çok önemliydi. Her zaman temiz ve güzel hazırlanmasına dikkat ediliyordu.

Biraz sonra kapı açıldı. Önce içeriye büyük ve zayıf bir kedi girdi. Tüylerinin yarısı siyah, yarısı beyazdı. Kedi sıcak olduğu için mangalın yanına gitti ve oraya kıvrıldı. Ondan biraz sonra falcı kadın odaya girdi. Ama bu gerçek ve başarılıydı. Çok hayali bir tiyatro oyununa veya komik bir oyundaki evin delisi karakterinin sahneye gelmesine benziyordu. Hiç acele etmiyordu. Zıplaya zıplaya yürüyordu. Ama amacı bu etki değildi. Çünkü aynı hareketlerde ve yüzünün ifadesinde iki ayrı yöne gitmeye çalışır gibi garip bir zıtlık vardı. İlk anda sakin ve sert görünüyordu. Durumu kontrol etmek istiyordu. Ama bunu nasıl yapacağını bilmiyordu. Bu yüzden böyle davranıyordu. Bize kısa bir hoş geldin dedikten sonra bizi bir süre ayakta inceledi. Adlarımızı sordu. Kısa bir duraklamadan sonra adlarımızı söyledik. O zaman “Memnun oldum" diye ellerimizi sıktı. Sonra kedisinin yanına geldi. Tam karşımıza, yerdeki katlı kilimin üzerine konan üçayaklı sandalyeye oturdu. Çok ciddi görünüyordu. Ne yaşlıydı ne de çirkindi. Saçlarının yarıdan fazlası dökülmüş. Saçları sarıya boyalı ama dipleri beyazdı. Yüzündeki makyajın bazı yerleri akmış. Büyük, açık mavi gözleri vardı. Boyu kısaydı. Ayakları çıplak ve kirliydi. Farklı görünüyordu. Biraz da korkutucuydu. Bu duygu çok güçlüydü. İnsan, yoksulluktan daha ağır bir durum vardı diye düşünüyordu.

Delilik her zaman üzücüdür. Ama gerçekten deli miydi? Sağ omzunda çok küçük kaplumbağa kabukları vardı. Kısa bir etek ve mavi bir ceket giyiyordu. Garip davranışları vardı. Bu yüzden ilk bakışta herkes onu deli sanıyor olabilir. O oturdu. Ben de kalın topuklarını daha iyi gördüm. Elini sıktım. Avucu terli ve yapış yapıştı. Bu yüzden kendimi daha kötü hissettim. O bunları bilmiyordu. Biraz sonra bize bir doktor gibi dikkatle baktı. Sonra "Bismillah" diyerek elini sağdaki fincana uzattı. Sessizce dua etti.

Sonra tekrar bize baktı ve bu kimin diye sordu. Fincan on, on beş dakikadır kapalıydı. Bu nedenle tabağına iyice yapıştı. Tekrar bize baktı. Bu bakışlar bizi rahatsız etti. Sende nazar var oğul diyerek düşündü. Tabağın kenarındaki telvelere dikkatle bakıyordu. Yüzü üzgündü. Belki de bizim de üzülmemizi istiyordu. İkimiz de hareket etmedik. Belirsiz ve bilinmeyen şeyler hakkında konuşmak kolaydır. Herkes bunu bilir. Ama falcı kadın haklıydı. Fala inandıktan sonra, nazara ne diye inanmamalı? Bunlar, birbirine çok benziyordu. Cemal adında birini tanıyor musun? Soruyu sorduktan sonra kadının yüzü gittikçe sertleşiyor ve gizemli görünüyordu.

Fincan hâlâ ters duruyordu. Ama biz bu soruya tepki vermedik. Cemal adında yüzlerce kişi var. Ama kadın önemli bir ipucu buldu. Sorular sormaya devam etti. Cemal, yoksa Cemile, Cevat, Cevdet, Cahit… C harfi aklımıza birçok kişi getirdi. Ama hangi kişiden söz ettiğini anlayamadık. Bu ilk denemenin boşa gitmesi kadını üzmedi. O, geçmişin ve geleceğin bilgisine sahipti. Bu yüzden kendinden emin konuşuyordu. Her şeyi söyleyebilir, her tahmini yapabilirdi. Hayatımızı çok iyi biliyor gibi bakıyordu. Biz bile bazı şeyleri bilmiyorduk. Ama o biliyor gibiydi.

- Bir dükkân... Mirastan bir dükkân var...

Ne yazık ki fincanın sahibi, babasından kalan mirasın acısını otuz yıl önce tanıdı. Ölüden hiçbir şey bekleyemez. Bu yüzden hiç sevinmedi. Sadece kadının başındaki kırmızı bez parçasına ve bu bez parçasının hareketine bakıyordu. Hayır, bu falcı kadın sadece zavallıydı. Ama kadın bunun farkında değildi. Başını tabağa eğdi. Fincandaki telveleri dikkatle inceliyordu. Birden omuzlarını oynattı. Bismillah diyerek fincanı tabaktan ayırdı. Yerinden biraz kalktı. Bize doğru eğildi. Sonra, Camiye mi yakın, yoksa kiliseye mi diye sordu.

C harfinden sonra sanki odaya gerçek bir insan geldi. Bunu herkes hissetti. Adı F harfiyle başlıyor. Fahamet, Fahriye yoksa Fahire... Fanni? Sesi bu sefer üzüntülü idi. Birden aklına son bir isim geldi: Fecriye... Hayır, hiçbiri değildi. Ama oldukça kararsız cevap verdik. Bu yüzden falcı kadın böyle bir kişinin var olduğuna inandı. Onun için devam etti:

- Ferran'dır, Ferran ...

Halinde bebeği için telaşlı loğusa bir annenin hali vardı. Ne yazık ki F harfi doğru adı bulmasına yardım etmedi. Tekrar ilk şekle döndü: Ferhan, Furuzan...

Ama hemen vazgeçmedi. Gözleri yeniden parladı. Sağ kolunu havaya kaldırdı. Sonra, yoksa bu babasının adı mı ya da F harfiyle başlayan biriyle mi yaşıyor diye sordu. En iyisi bu isim konusunu burada bitirelim.

Karanlığa yeni baştan ve başka bir taraftan saldırmak lazımdı. Bir iş yerinde mi çalışıyor? Yoksa terzi mi? Yetenekli ve nazik bir kızdır. Ayrıca şanslıdır. Oğul bu kadını üç kişi arasında görüyorum. Anasıdır o... Yoksa ablası mı?

İki arkadaş bu kadın hakkında hiçbir şey bilmiyordu. Falcı kadın bu kadını hayal etmiş.

-Bir başka kadın daha var, Margrit, Melek, M harfi. Ama hastadır biliyorsun. İkisi de Margrit ya da Melek adlı bu kadını tanımıyordu. Ama bu ikinci kadının hastalığına epeyce üzüldüler. Kadın mahkemeden resmî bir belge almış. Belgede mühür ve imza vardı. Hep seni düşünüyor. Kızı üzmüşsün. Ama gideceksin. Düpedüz değil, köprübaşında tut ki Kadıköy iskelesinde göreceksin... Ondan sonra.

Sonra bir adam askerden geliyor. Uzun ve yakışıklı. Asker mi yoksa pilot mu?.. Ama kadın güzeldir. Burası özellikle önemliydi. Yavaş yavaş portre çizilmeğe başladı. Esmer, uzun boylu ve tatlı gülüşlü genç kadını gözümüzde canlandırdık.

Sonra kadın:

Bir şoför var, çok para yemiş… diye devam etti. Falcı kadın kısa bir süre düşünerek para miktarını hesapladı: Yaklaşık üç, dört bin lira... Ama bu kadın da çok zengin... Kibar kadındır, Maçka'da oturur, yoksa Şişli'de, ama öbür kadın onu çekemez... Rakip şoför artık yoktu. Bu sebeple memnunduk. - Sakın şüphelenme... Birinci kadınla kimse yaşamaz, o senindir. Yanımdaki arkadaşım gözlerini kapattı. İlk kadını yeniden hayal etti. Sonra, M, Melahat, o da kocasından boşanmış... Ama sizden bir kadın daha tanır.

Fincan gittikçe genişliyordu. Hemen hemen bir mahalle dolusu olduk. Bir yolculuğa çıkacak mısın? Çocuğun hastalığı geçecek ama ayak kazası var. Zavallı çocuk... Arkadaşımın yüzünün rengi birden değişti. Ama kadın hemen sözünü düzeltti: Bir yerden atlama. Gece bir hediye verdin. Herkes bu hediyeyi beğendi. O kadın anasıdır. Elindeki her şeyi satar: o olmasa ölür senin için o kadın?1 - Kimin anası, çocuğun anası mı? Ama kadının başındaki süsler, soruya cevap verilmesine fırsat vermedi. O kadar telaş ve korku saldılar ki, biz de geri çekildik. Sonra falcı kadın son tahminini söyledi:

– Ah, zavallı ölecek...

Farkında olmadan sorduk:

-Kim?

Kadın kısa bir süre düşündü. Galiba zararsız bir cevap arıyordu.

-Askerden geldi… Hem bir kaza ile... Ama sende de çok para var, ha... Valisin sen? Yoksa... Avukat, avukat dedim ama mahkemeyle ilgili biri var. Yoksa hastahane mi? doktor odası gibi bir yer görünüyor. Bütün hatası sürekli yüzümüze bakmasıydı. Biz sakin duruyorduk. Bu durum onu şaşırtıyordu. Sonra başka açıklamalar bulmaya çalışıyordu.

O kadın hasta. Ama ilk kadın değil. O kadın onu kıskanıyor. O da ölecek… Sonra Ankara'da bir milletvekili var. O, büyük bir yardım edecek. Bir otelde bir araya geldiniz. Sonra bir eğlence yerine gittiniz. İşte orada sizi gördüler. Bir süre daha düşündü: Yattığın yerde bir düşmanın var ama sana bir şey yapmayacak... diye söyledi. Bir ziyafet sofrası... Aman Allah'ım sakın kavga etme, bu ay şansın iyi değil. Bir mektup alacaksın... Yoksa bir haber

gelir. Sonra telefon çalar. Gece seni çağırırlar. Ama bayramı bekle. Ancak o zaman sorunlarınız biter.

Yüzü yine değişti. Bir süre düşündü. Sonra Kiracı çıkacak. Ama sen orada oturmayacaksın… diye belirtti.

Oğlun ne zaman Amerika'daysa o zaman onu gördün. Adı Zeynep'tir, yoksa Zehra, hayır ikisi de değil. Fakat kadın söylemeğe vakit bırakmıyordu ki. Yarın sabah vapurla gelecek. Kadıköy'den. Daktilon çok kıskançtır değil mi. o da rahat durmaz ya...

Falcı kadının kafasındaki kırmızı süs, insan halinin çaresizliğine bir daha acıdı. Bu durum saatlerce devam edebilir. Ben dostuma, dostum bana baktı ve yavaşça kalktık. Parasını vererek çıktık, falcı kadın arkamızdan:

-İşin olursa bir kutu şeker isterim, diye söyleniyordu.

Kapıdan çıkarken ben gülmek istedim, fakat arkadaşım bırakmadı:

-Ne istiyorsun?.. diye sordu. Sana on dakikada herkesin romanını anlattı...

Yetişmez mi? Gerçekten de böyle idi; beş on dakika içinde herkesin romanını anlattı. İnsanların hayatı çoğu zaman küçük ve önemsiz şeylerin etrafında geçmez mi?

YABANCILARA TÜRKÇE ÖĞRETİMİNDE B1 DÜZEYİNE YÖNELİK METİN UYARLAMA: SAİT FAİK ABASIYANIK'IN *İPEKLİ MENDİL* HİKÂYESİ ÖRNEĞİ

TEXT ADAPTATION FOR THE B1 LEVEL IN TEACHING TURKISH TO FOREIGNERS: A CASE STUDY OF SAIT FAIK ABASIYANIK'S STORY "*İPEKLI MENDIL*"

Öğr. Gör. Havana KAYNAK
Türkçe Öğretimi Uygulama ve Araştırma Merkezi, Gümüşhane Üniversitesi, Gümüşhane, TÜRKİYE
ORCID ID: 0009-0000-8607-7620

ÖZET

Türkçenin yabancı dil olarak öğretiminde özgün edebî metinlerin kullanımı, öğrencilerin dil becerilerinin yanı sıra hedef kültürü tanımalarına da katkı sağlamaktadır. Özgün edebi metinlerin kullanımı, hedef kitlenin kültür düzeyini artıracağı gibi hedef dile ait düşünme ve algılama sürecini şekillendirmede de fayda sağlayacaktır. Ancak özgün metinlerin özellikle de klasik edebî metinlerin dil ve anlatım özellikleri, başlangıç ve orta düzey öğreniciler için çeşitli güçlükler oluşturabilmektedir. Bu bağlamda, özgün yazınsal metinlerin ders kitapları ile yardımcı kaynaklarda yer alabilmesi, bu metinlerin belirli ölçütler doğrultusunda hedef dil düzeylerine uygun şekilde uyarlanmasını gerekli kılmaktadır. Türk edebiyatının önemli eserleri, Türkçeyi yabancı dil olarak öğrenen bireylerin yararlanabilmesi amacıyla farklı dil yeterlik düzeylerine uyarlanarak yayımlanmaktadır. Bu eserler sayesinde öğreniciler yalnızca Türkçenin dilsel özelliklerini geliştirmekle kalmamakta, aynı zamanda Türk toplumunun kültürel değerlerini, yaşam biçimini ve düşünce dünyasını da tanıma fırsatı bulmaktadır. Bu bağlamda, Avrupa Dilleri Ortak Çerçeve Programı (CEFR) doğrultusunda gerçekleştirilen metin uyarlamaları, edebî eserlerin yabancı dil olarak Türkçe öğretiminde daha etkili ve işlevsel biçimde kullanılmasına katkı sağlamaktadır. Bu çalışmada, ders dışı okuma materyallerine duyulan gereksinime katkı sağlamak amacıyla, Türk edebiyatının önemli yazarlarından Sait Faik Abasıyanık'ın *İpekli Mendil* adlı hikâyesi, B1 düzeyindeki Türkçe öğrenicilerine yönelik olarak metin değiştirim ölçütleri çerçevesinde uyarlanmıştır. Çalışmanın temel amacı, uyarlanan metni Türkçeyi yabancı dil olarak öğrenen bireylerin okuma becerilerini geliştirmede kullanılabilecek yardımcı bir kaynak hâline getirmektir. Uyarlama sürecinde, B1 düzeyindeki öğrenicilerin kazanmış olması beklenen dil yapıları, *Yeni İstanbul Uluslararası Öğrenciler İçin Türkçe Ders Kitabı -A1, A2 ve B1-* ile Avrupa Dilleri Ortak Çerçeve Programı (CEFR) esas alınarak belirlenmiş; metin, bu ölçütler doğrultusunda yeniden düzenlenmiştir. Çalışmada nitel araştırma yöntemlerinden doküman analizi kullanılmış; metin, sözcük seçimi, cümle yapısı, metin bütünlüğü ve kültürel ögelerin sunumu bakımından incelenmiştir. Uyarlama sürecinde B1 düzeyi öğrenicilerinin dil yeterlikleri dikkate alınarak uzun ve karmaşık cümleler sadeleştirilmiş, kullanım sıklığı düşük sözcükler daha yaygın karşılıklarıyla değiştirilmiş, anlamayı güçleştiren anlatım özellikleri

yeniden yapılandırılmıştır. Bununla birlikte, hikâyenin edebî niteliğini ve kültürel dokusunu oluşturan temel unsurlar korunmaya özen gösterilmiştir. Çalışmanın sonucunda, *İpekli Mendil* hikâyesinin uygun uyarlama ölçütleri doğrultusunda B1 düzeyinde kullanılabilecek bir okuma metnine dönüştürülebileceği öngörülmektedir. Araştırmanın, yabancılara Türkçe öğretiminde klasik Türk edebiyatı ürünlerinin ders materyali olarak kullanımına ilişkin çalışmalara katkı sağlaması ve edebî metin uyarlamalarına yönelik yeni araştırmalara örnek oluşturması beklenmektedir.

Anahtar Kelimeler: Yabancılara Türkçe öğretimi, metin uyarlama, Sait Faik Abasıyanık, *İpekli Mendil*.

ABSTRACT

The use of authentic literary texts in teaching Turkish as a foreign language contributes not only to the development of students' language skills but also to their understanding of the target culture. Incorporating such texts enhances the cultural awareness of the target audience and helps shape the thought processes and perceptions associated with the target language. However, the linguistic and stylistic characteristics of authentic texts-particularly classical literary works-can pose various challenges for beginner and intermediate learners. Consequently, including authentic literary texts in textbooks and supplementary materials necessitates adapting them to suit specific target language proficiency levels based on established criteria. Significant works of Turkish literature are published in adapted versions tailored to different proficiency levels for individuals learning Turkish as a foreign language. Through these works, learners not only improve their command of the language but also gain insight into the cultural values, lifestyle, and worldview of Turkish society. In this context, text adaptations carried out in alignment with the Common European Framework of Reference for Languages (CEFR) contribute to the more effective and functional use of literary works in teaching Turkish as a foreign language. To address the need for supplementary reading materials, this study adapts the short story *İpekli Mendil* (The Silk Handkerchief)-written by Sait Faik Abasıyanık, a prominent figure in Turkish literature-for B1-level Turkish learners, adhering to established text adaptation criteria. The primary objective is to transform the adapted text into a supplementary resource that can be used to enhance the reading skills of individuals learning Turkish as a foreign language. During the adaptation process, the linguistic structures expected of B1-level learners were identified based on the *New Istanbul Turkish for International Students* textbook series (levels A1, A2, and B1) and the CEFR, and the text was restructured in accordance with these criteria. The study employed document analysis-a qualitative research method-to examine the text in terms of word choice, sentence structure, textual coherence, and the presentation of cultural elements. During the adaptation process, the language proficiency of B1-level learners was taken into account; consequently, long and complex sentences were simplified, low-frequency words were replaced with more common equivalents, and stylistic features that hindered comprehension were restructured. At the same time, care was taken to preserve the

fundamental elements that constitute the story's literary quality and cultural fabric. The study concludes that the story *İpekli Mendil* can be transformed into a reading text suitable for the B1 level by adhering to appropriate adaptation criteria. It is anticipated that this research will contribute to the body of work regarding the use of classic Turkish literary texts as instructional materials in teaching Turkish to foreigners and serve as a model for future research on literary text adaptation.

Keywords: Teaching Turkish to foreigners, text adaptation, Sait Faik Abasıyanık, *İpekli Mendil*.

GİRİŞ

Yabancılara Türkçe öğretiminde alıcı -öğrenci- ve verici -öğretici- arasındaki anlaşmayı kolaylaştıran en önemli unsur dildir. Dil, geçmişten günümüze insanlar arasındaki iletişimin sağlanmasında ve kültürel birikimin nesilden nesile aktarılmasında önemli bir rol üstlenmiştir. Bu yönüyle dil, insanlık tarihi açısından vazgeçilmez bir unsur olarak değerlendirilmektedir. Nitekim dil kavramı için Türk Dil Kurumu (t.y.) tarafından; "dil geçmişten günümüze kadar insanlar arasındaki iletişimi ve anlaşmayı sağlayan, uzlaşmaya dayalı olarak oluşturulmuş sesli veya görsel işaretler sistemi; lisan. Düşünce ve duyguları bildirmeye yarayan herhangi bir anlatım aracı" şeklinde farklı tanımlamalar yapılırken, Turan Karataş'ın (2001) *Ansiklopedik Edebiyat Terimleri Sözlüğü* adlı kitabında ise dil için;

> Toplumu "millet" yapan unsurların başında geldiği için "milli" bir mahiyet arz eder. Edebiyatın biricik/ana malzemesi dildir. İnsanlar arasında anlaşma, iletişim kurma vasıtası olmasının yanında duygu ve düşüncelerin söz veya yazı aracılığıyla başkalarına aktarılmasına, iletilmesine; kıymet ifade eden sözün korunmasına yarayan çok mühim bir canlı unsur/varlık. (s.104)

tanımlamaları yer almaktadır. Bu tanımlamalar dikkate alındığında dilin iletişim ve kültür aktarımı işlevlerini öne çıkaran farklı tanımlarla açıklanmaya çalışıldığını söylemek mümkündür.

İnsanlar göç, ticaret, eğitim, turizm vb. gibi çeşitli nedenlerle yabancı dil öğrenmektedir. Bu sebeple Türkçe de geçmişten günümüze farklı topluluklar tarafından çeşitli amaçlarla öğrenilen diller arasında yer almaktadır. Türkçenin yabancılara öğretilmesine yönelik çalışmaların tarihî geçmişi oldukça eskiye dayanmaktadır. Nitekim Dîvânu Lugâti't-Türk, Muhâkemetü'l-Lugateyn ve Codex Cumanicus gibi eserlerin Türkçenin farklı topluluklara öğretilmesi amacıyla kaleme alınmış olması, Türkçenin yabancı dil olarak öğretiminin tarihsel gelişimini göstermesi bakımından önem taşımaktadır. Günümüzde ise yurt içinde Türkçe öğretimi amacıyla faaliyet gösteren TÖMER'lerin, yurt dışında ise başta Yunus Emre Enstitüsü olmak üzere çeşitli kurum ve kuruluşların faaliyet göstermesi, Türkçeye yönelik ilginin ve Türkçenin yabancı dil olarak öğretiminin yaygınlaştığını ortaya koymaktadır.

Günümüzde yabancı dil öğretimi, bireysel tercihler doğrultusunda belirlenen ölçütlerden ziyade, uluslararası kabul gören belirli standartlar çerçevesinde gerçekleştirilmektedir. Bu doğrultuda, yabancı dil öğretiminde temel başvuru kaynaklarından biri olan Avrupa Dilleri

Ortak Çerçeve Programı (CEFR), dil öğretim ve öğrenme süreçlerinin planlanması, uygulanması ve değerlendirilmesine yönelik ortak bir çerçeve sunmaktadır. Metin, yabancı dil öğrenen bireylerin hedef dili iletişim amacıyla kullanabilmeleri için sahip olmaları gereken bilgi, beceri ve yeterlikleri açıklamakta; dil kullanımına ilişkin yeterlikleri belirli düzeyler içerisinde tanımlamaktadır. Bunun yanı sıra öğrenenlerin dil öğrenme süreçlerindeki ilerlemelerinin izlenmesine ve değerlendirilmesine yönelik ölçütler ortaya koymaktadır. Bu yönüyle Avrupa Dilleri Ortak Çerçeve Programı (CEFR), yabancı dil öğretiminde hedeflerin belirlenmesi, öğretim sürecinin yapılandırılması ve öğrenenlerin dil yeterliklerinin ölçülüp değerlendirilmesi açısından önemli bir başvuru kaynağı niteliğindedir (CEF 2001:1).

Yabancılara Türkçe öğretiminde dil becerileri, genel olarak anlamaya dayalı beceriler ve anlatmaya dayalı beceriler olmak üzere iki temel grupta ele alınmaktadır. Anlamaya dayalı beceriler okuma ve dinleme, anlatmaya dayalı beceriler ise konuşma ve yazma becerilerinden oluşmaktadır. Bu dört temel dil becerisine ek olarak dil bilgisi öğretimi de yabancı dil öğretim sürecinin önemli bir bileşenini oluşturmaktadır. Öğrencilerin öğrendikleri dil bilgisi yapılarını dört temel dil becerisi aracılığıyla kullanmaları ve bu yapıları iletişimsel bağlamlarda işlevsel hâle getirmeleri beklenmektedir.

Yabancılara Türkçe öğretiminde dil yeterlikleri A1, A2, B1, B2, C1 ve C2 olmak üzere altı temel düzeyden oluşmaktadır. A1, A2, B1, B2 ve C1 düzeyleri hemen hemen her dil kursunda yer alırken C2 dil düzeyi sadece bazı dil kurslarında mevcuttur. Öğretim sürecinde dil bilgisi yapıları ve dil becerilerine ilişkin kazanımlar, düzeylere göre basitten karmaşığa ve kolaydan zora doğru aşamalı bir biçimde sunulmaktadır. Her düzeyin sonunda öğrencilerin okuma, dinleme, konuşma ve yazma becerileri açısından belirli kazanımlara ulaşmaları beklenmektedir. Öğrencinin ilgili düzeye yönelik kazanımları edinmesi, bir sonraki düzeye geçebilmesinin temel koşullarından biri olarak değerlendirilmektedir. Tüm düzeylerin başarıyla tamamlanmasının ardından öğrencinin dil yeterliğini gösteren bir sertifika alması mümkün olmaktadır.

Öğrencilerin dört temel dil becerisini geliştirmeye yönelik uygulamalarda farklı yöntem ve tekniklerden yararlanılmaktadır. Konuşma becerisinde öğrencilerden, öğrendikleri kelime ve dil bilgisi yapılarını doğru ve uygun bir biçimde kullanarak sözlü iletişim kurmaları beklenmektedir. Dinleme becerisinde ise öğrencilerin düzeylerine uygun, hedef kelime ve dil bilgisi yapılarını içeren metinleri dinlemeleri ve dinledikleri metni anlamaya yönelik çeşitli görevleri yerine getirmeleri istenmektedir. Yazma becerisinde öğrencilerden düzeylerine uygun bir konu veya soruya, öğrendikleri kelime ve dil bilgisi yapılarını kullanarak anlamlı, doğru ve tutarlı bir metin oluşturmaları beklenmektedir. Okuma becerisinde ise öğrencilerin düzeylerine uygun metinleri anlamaları ve metne ilişkin hazırlanan açık uçlu, çoktan seçmeli, doğru-yanlış vb. soruları yanıtlamaları vasıtasıyla okuduğunu anlama becerilerinin geliştirilmesi ve değerlendirilmesi amaçlanmaktadır. Bu bağlamda dört temel dil becerisi ve dil bilgisi öğretimi birbirinden bağımsız süreçler olarak değil, öğrencinin hedef dili anlamlı ve işlevsel bir biçimde kullanmasını sağlayan bir bütünün parçaları olarak değerlendirilmektedir.

Öğrencinin öğrendiği dil bilgisi yapılarını ve kelimeleri farklı beceri alanlarında kullanabilmesi, dil yeterliğinin gelişimi açısından önem taşımaktadır.

Yabancılara Türkçe öğretiminde temel dil becerilerinden biri olan okuma, öğrencilerin hedef dili anlamlandırmalarına ve dilsel yeterliklerini geliştirmelerine önemli katkılar sağlamaktadır. Öğrencilerin okuma becerilerini geliştirebilmeleri için seviyelerine uygun ve ilgi çekici okuma metinleri ile pratik yapmaları gerekmektedir.

Metnin yazılma amacı, metnin türünü belirleyen temel unsurlardan biridir. Yazar, aktarmak istediği olay, duygu veya düşünceyi, belirlediği metin türünün özelliklerine uygun olarak okuyucuya sunar (Erkul 2004:178). Nitekim kitapları oluşturan yazılar her türlü okuma etkinliği için ana kaynaklardır. Ancak yabancılara Türkçe öğretiminde kullanılabilecek, farklı dil düzeylerine uygun okuma kitapları, edebî metinler, şiirler, atasözleri vb. materyallerin sayısının sınırlı olması, bu becerinin geliştirilmesi sürecinde çeşitli güçlüklerin ortaya çıkmasına neden olmaktadır.

Metin, başı ve sonu ile kapalı bir yapı oluşturan dilsel göstergelerin art arda geldiği anlamlı yapıdır (Günay 2007:44). Her yazı mutlaka bir amaç için yazılmıştır ve konusu vardır. Fakat özellikle özgün edebî metinlerin içerdiği söz varlığı, dil bilgisi yapıları, cümle uzunlukları ve kültürel unsurlar, metinlerin öğrencilerin dil düzeylerine doğrudan uygunluğunu engelleyebilmektedir. Bu nedenle yabancılara Türkçe öğretiminde öğrencilerin dil düzeylerine uygun nitelikte okuma materyalleri oluşturmak amacıyla metin uyarlama çalışmalarından yararlanılmaktadır. Metin uyarlama sürecinde özgün metnin temel anlamı ve anlatım bütünlüğü mümkün olduğunca korunurken öğrencilerin düzeyini aşan kelime, dil bilgisi yapıları ve söz dizimsel özellikler hedef düzeye uygun hâle getirilmektedir. Böylece özgün edebî metinlerin yabancı dil olarak Türkçe öğrenen öğrenciler tarafından daha anlaşılır ve işlevsel bir biçimde kullanılabilmesi amaçlanmaktadır.

Bu çalışmada da yabancılara Türkçe öğretiminde B1 düzeyindeki öğrencilerin okuma becerilerini geliştirmeye yönelik bir metin uyarlama çalışması gerçekleştirilmiştir. Çalışmanın metnini Sait Faik Abasıyanık'ın *İpekli Mendil* adlı hikâyesi oluşturmaktadır. Özgün metin; kelime hazinesi, dil bilgisi yapıları, cümle kuruluşları ve söz dizimsel özellikleri bakımından incelenmiş ve B1 düzeyindeki öğrencilerin sahip olması gereken dil kazanımları dikkate alınarak uyarlanmıştır. Uyarlama sürecinde metnin özgün anlatımının ve temel anlamının korunmasına özen gösterilmiş; öğrencilerin B1 düzeyinde karşılaşmaları beklenen kelime ve dil bilgisi yapıları esas alınarak metnin anlaşılabilirliğinin artırılması amaçlanmıştır.

Araştırmanın Amacı ve Önemi

Araştırmanın amacı, Türkçenin yabancı dil olarak öğretiminde kaynaklık edebilecek özgün metinlerin dil düzeylerine uygun biçimde uyarlanmasına yönelik bir örnek ortaya koymak ve metin uyarlama sürecine ilişkin sistematik bir yaklaşım geliştirilmesine katkı sağlamaktır. Ayrıca bu çalışma ile Türk edebiyatına ait özgün metinlerin yabancı dil olarak Türkçe öğretiminde kullanımına dikkat çekilmek ve bu metinlerin öğretim sürecinde daha yaygın

biçimde kullanılmasına katkı sunulmak hedeflenmektedir. Bu doğrultuda, Sait Faik Abasıyanık'ın *İpekli Mendil* adlı hikâyesinin B1 düzeyine uyarlaması yapılmıştır.
Araştırmanın önemi, yabancı dil olarak Türkçe öğretiminde özgün edebî metinlerin kullanımı, öğrenicilerin hem dil becerilerini geliştirmelerine hem de Türk kültürünü tanımalarına katkı sağlamaktadır. Ancak bu metinler, dilsel ve yapısal özellikleri nedeniyle başlangıç ve orta düzey öğreniciler tarafından doğrudan kullanılmaya her zaman elverişli değildir. Bu durum, hedef dil düzeyine uygun metin uyarlamalarını gerekli kılmaktadır. Bu çalışmada, Sait Faik Abasıyanık'ın *İpekli Mendil* adlı hikâyesi B1 düzeyine uyarlanarak yabancı dil olarak Türkçe öğretiminde özgün edebî metinlerin nasıl öğretim materyaline dönüştürülebileceğine ilişkin uygulamalı bir örnek sunulmaktadır. Araştırmanın, metin uyarlama sürecinde izlenen ölçütlerin sistematik biçimde ortaya konulması bakımından alana katkı sağlaması; öğreticilere, materyal geliştiricilere ve araştırmacılara yol gösterici olması beklenmektedir. Ayrıca çalışma, Türk edebiyatına ait özgün eserlerin yabancı dil olarak Türkçe öğretiminde daha etkin kullanılmasına katkı sunması ve metin uyarlama alanında yapılacak yeni araştırmalara örnek teşkil etmesi bakımından önem taşımaktadır.

Araştırmanın Sınırlılıkları

Araştırma Sait Faik Abasıyanık'ın *İpekli Mendil* adlı hikâyesinin B1 düzeyine uyarlanması ile sınırlandırılmıştır. Uyarlama yapılırken yabancılara Türkçe öğretimi alanında kullanılan *Yeni İstanbul Uluslararası Öğrenciler İçin Türkçe Ders Kitabı* setinin A1, A2 ve B1 düzeylerindeki sözcükler, dil bilgisi kuralları ve Avrupa Dilleri Ortak Çerçeve Programı (CEFR) esas alınmıştır. Bu çalışma kapsamında uyarlanan metin, öğrenciler üzerinde test edilmemiştir.

Sait Faik Abasıyanık ve Hikâyeciliği

18 Kasım 1906'da Adapazarı'nda dünyaya gelen Sait Faik Abasıyanık'ın asıl adı Mehmet Sait'tir (Çelik 2020). Hiç evlenmeyen Abasıyanık, 11 Mayıs 1954 yılında alkol düşkünlüğü nedeniyle yakalandığı siroz hastalığı sonucunda vefat etmiştir (Okay 2008: 583).
Varlıklı bir ailede büyüyen Abasıyanık, Adapazarı başta olmak üzere İstanbul, İsviçre ve Fransa'da da eğitim görmüştür. Sait Faik, eğitimini tamamlamak amacıyla gittiği Fransa'da, Fransızcasını geliştirerek Batı edebiyatını da daha yakından tanıma fırsatı bulmuştur (Okay 2008: 583). Çalışmak için girdiği birçok yerde düzen tutturamayan Abasıyanık, liseye gittiği dönemlerde önceleri şiir yazmış sonradan ise hikâyeler yazarak edebiyata yönelmiştir (Çelik 2020). İlk hikâyesi *Uçurtmalar*, Milliyet gazetesinde yayımlanmış; ilk şiiri *Hamal* ise Mektep dergisinde çıkmıştır (Çelik 2020). 1936 yılında da ilk kitabı olan *Semaver*'i yayımlamıştır. Önceleri Maupassant tarzında hikâyeler yazan ve kendisine Refik Halit Karay, Ömer Seyfettin, Reşat Nuri Güntekin gibi önemli hikâyecileri örnek alan Abasıyanık, sonraları kendine özgü bir dil geliştirerek hikâyeler yazmaya başlamıştır (Çelik 2020). İlk dönemler yazdığı hikâyelere bakıldığında, gözlem ve acıma duygusunun ağırlıklı olarak kullanıldığı dikkat çekmektedir (Çelik 2020). *Lüzumsuz Adam* adlı eserinden itibaren kendine özgü bir dil

geliştiren Abasıyanık, hikâyelerinde insanın basit hayatını daha içten, samimi ve sevgi temasını önceler bir pencereden ben diliyle okura sunmaya başlamıştır (Çelik 2020). *Alemdağda Var Bir Yılan* adlı eseriyle Abasıyanık, geleneksel hikâye formunun dışına çıkarak bireyin iç dünyasını ve bilinçaltını ele alır (Çelik 2020). Abasıyanık'ın hikâyelerine genel olarak bakıldığında onun, hayvanlar, çocuklar, Anadolu'dan İstanbul'a göç edenler, tabiat, gecekondu ve farklı semtlerde yaşayanlar, özellikle deniz ve balıkçılıkla ilgilenen insanlar başta olmak üzere her türden canlı ve cansıza yer verdiği söylenilebilir (Okay 2008: 583-584). Ömer Seyfettin'den sonra Türk hikâyeciliğinin ikinci safhası olarak kabul edilen Abasıyanık (Okay 2008: 584), 1953 yılında Mark Twain Cemiyeti Şeref Üyeliği'ne seçilen ve ödül alan ilk Türk yazar olma özelliğine de sahiptir (Çelik 2020).

YÖNTEM

Araştırma Modeli

Bu çalışma, nitel araştırma yöntemleri çerçevesinde yürütülmüştür. Çalışmanın veri toplama ve analiz etme sürecinde doküman analizi ve metin çözümleme tekniklerinden yararlanılmıştır. Araştırmanın esas veri kaynağını Sait Faik Abasıyanık'ın *İpekli Mendil* adlı hikâyesi oluşturmaktadır. Metin analizi aşamasında ise metin uyarlama sürecinde kaynak kabul edilen Türkçe öğretim seti (*Yeni İstanbul Uluslararası Öğrenciler İçin Türkçe Ders Kitabı*) ve Avrupa Dilleri Ortak Çerçeve Programı (CEFR) esas alınmıştır. Bu kaynaklar doğrultusunda Sait Faik Abasıyanık'ın *İpekli Mendil* adlı hikâyesi sistematik biçimde incelenmiş ve metin çözümleme tekniği ile hikâyenin dil, söz varlığı ve cümle yapıları değerlendirilerek söz konusu metin B1 düzeyine uygun hâle getirilmiştir.

Evren ve Örneklem

Çalışmanın evrenini Sait Faik Abasıyanık'ın hikâyeleri oluşturmaktadır. Araştırmanın örneklemini ise yazarın *Seçme Hikâyeler* (1990) adlı kitabında yer alan *İpekli Mendil* hikâyesi oluşturmaktadır. Çalışmamızın evrenini oluşturan hikâye kitabı, Milli Eğitim Bakanlığı Yayınları tarafından çıkarılmıştır.

Veri Toplama Araçları

Bu çalışmada Sait Faik Abasıyanık'ın *İpekli Mendil* adlı hikâyesi, Avrupa Dilleri Ortak Çerçeve Programı'nda yer alan (CEFR) A1, A2 ve B1 düzeyi dil yeterlilikleri ve kazanımları esas alınarak uyarlanmıştır. Buna ilave olarak Türkçe öğretiminde kullanılan *Yeni İstanbul Uluslararası Öğrenciler İçin Türkçe Ders Kitabı*'nın (A1, A2 ve B1) söz varlıklarıyla dil bilgisi yapıları da veri toplama aracı olarak kullanılmıştır.

Veri Analizi

Bu çalışmanın veri analiz süreci dört aşamada yürütülmüştür. İlk aşamada özgün metin dil, içerik ve söz varlığı bakımından çözümlenmiştir. İkinci aşamada metnin hedef dil düzeyi

belirlenerek çalışmada kullanılacak veri toplama araçları incelenmiştir. Üçüncü aşamada veri toplama araçlarından elde edilen ölçütler doğrultusunda metin uyarlama işlemi gerçekleştirilmiştir. Son aşamada ise uyarlanan metin yeniden analiz edilerek metnin hedef düzeye uygunluğu değerlendirilmiştir.

BULGULAR

Özgün Metnin (*İpekli Mendil* Hikâyesi) Özellikleri

Edebî Özellikler

İpekli Mendil, ilk olarak 15 Nisan 1934 tarihinde *Varlık* dergisinde yayımlanmış, daha sonra 1936 yılında ilk baskısı yapılan ve Abasıyanık'ın ilk öykü kitabı olan *Semaver'*de on dokuzuncu hikâye olarak yer almıştır. Bu hikâyeyi Abasıyanık, Bursa Erkek Lisesi'nde öğrenim gördüğü yıllarda, bir ödev olarak kaleme almıştır.

Tema: Hikâyenin teması, küçük bir çocuğun sevdiği kıza hediye etmek istediği ipek mendili satın alacak maddi imkâna sahip olmadığı için bir ipek fabrikasından mendil çalmaya çalışırken hayatını kaybetmesidir. Olay örgüsü tek bir ana olay etrafında şekillenmekte olup karmaşık bir kurgu içermemektedir. Bu yönüyle hikâye, B1 düzeyindeki öğrencilerin olay akışını takip edebileceği ve genel anlamını kavrayabileceği niteliktedir.

Olay Örgüsü: Hikâye anlatıcısı -isimsiz- Bursa'da yer alan bir ipek fabrikasında, ipekli mendil çalmak amacıyla fabrikaya giren fakir bir çocukla karşılaşır. Çocuğun sevdiği kız için mendili çalmaya çalıştığını öğrenen anlatıcı onu affeder. Ancak çocuk daha sonra tekrardan fabrikaya ipekli mendil çalmak için ağaçtan tırmanarak camdan girer ve camdan çıkıp ağaçtan aşağı inerken düşerek hayatını kaybeder. Hikâyenin sonunda çocuğun avucunda sevdiği kız için çaldığı ipekli mendil bulunur.

Karakterler: Hikâyede isimsiz başkişi (aynı zamanda anlatıcı), hırsız, kapıcı ve cambaz olmak üzere dört karakter yer almaktadır. Hikâyeyi birinci kişi -ben- ağzından aktaran anlatıcı, aynı zamanda başkişidir. Ancak bu karakterin kimliği ve kişisel özellikleri hakkında ayrıntılı bilgi verilmemektedir. Hırsız da isimsiz bir karakter olup sevdiği kıza hediye etmek amacıyla bir ipek mendil satın alacak maddi imkâna sahip olmadığı için ipek fabrikasına hırsızlık yapmak üzere gelen on beş yaşındaki bir çocuktur. Kapıcı, ipek fabrikasında görev yapan kişidir. Onun hakkında da ayrıntılı bilgi bulunmamaktadır. Cambaz ise Bursa'ya gelen isimsiz bir gösteri sanatçısı olarak hikâyede yer almaktadır. Genel olarak hikâyedeki karakterlerin tamamı isimsizdir ve yazar, karakterlerini ayrıntılı biçimde tanıtmak yerine olay örgüsü içerisindeki işlevleri doğrultusunda ele almıştır.

Zaman ve Mekân: Hikâyede iki farklı zaman dilimi bulunmaktadır. Bunlardan ilki, başkişi ve hırsızın ilk kez karşılaştığı akşam vaktidir. İkinci zaman dilimi ise hırsızın ağaçtan düşerek hayatını kaybettiği sabaha karşıdır. Bunun yanı sıra hikâyede, başkişinin geçmişe ilişkin anılarını aktardığı geri dönüş (flashback) tekniğine de yer verilmiştir. Mekân bakımından hikâyede Bursa başlıca çevreyi oluşturmaktadır. Olayların büyük bölümü İpek Fabrikası'nda geçmektedir. Bunun yanında başkişi ile hırsızın ilk karşılaşmalarında sohbet ettikleri kapıcı

odası, başkişinin odası ile anlatıcının geri dönüş tekniği aracılığıyla hatırladığı Emir Sultan ve Bursa'da yer alan Gökdere de hikâyedeki diğer mekânlardır. Ancak bu mekânlar ayrıntılı tasvirlerle verilmemiş, olay örgüsünün ilerleyişine hizmet edecek ölçüde işlenmiştir.

Anlatım Tekniği: Hikâyede kahraman bakış açısı kullanılmıştır. Anlatıcı, aynı zamanda hikâyenin başkişisi olduğundan olayları yalnızca kendi gördüğü, yaşadığı ve bildiği kadarıyla aktarmaktadır. Bu durum, anlatımın öznel bir nitelik kazanmasını sağlarken duygu ve düşüncelerin samimi ve içten bir üslupla okuyucuya yansıtılmasına da katkıda bulunmaktadır. Anlatım teknikleri bakımından ise hikâyede özetleme tekniği ağırlıklı olarak kullanılmıştır. Olaylar ayrıntılı betimlemelere yer verilmeden, kronolojik akış içerisinde kısa ve öz bir biçimde aktarılmıştır. Böylece anlatımın akıcılığı korunmuş ve olay örgüsünün ilerleyişi ön planda tutulmuştur.

Dil Bilgisel Özellikler

Kelime ve Cümle Sayısı: Çalışmamıza konu olan hikâye 96 cümle (%12) ve 686 kelimeden (%88) oluşmaktadır. Metindeki en uzun cümle 25 kelime, en kısa cümle ise 1 kelimeden meydana gelmektedir. Cümle uzunlukları incelendiğinde metinde belirgin bir çeşitlilik olduğu görülmektedir. Özellikle uzun cümlelerin, B1 düzeyindeki öğrenicilerin okuduğunu anlama becerisini olumsuz etkileyebileceği düşünülmektedir. Bu nedenle uyarlama sürecinde söz konusu cümlelerin daha kısa, açık ve anlaşılır biçimde yeniden düzenlenmesi uygun görülmüştür.

Kelime Türlerindeki Dağılım: Metindeki kelime türleri, isim soylu ve fiil soylu kelimeler olmak üzere iki kategoride değerlendirilmiştir. Bu değerlendirme sonucunda çalışmamıza konu olan hikâyede 521 isim soylu (%76) ve 165 fiil soylu (%24) kelime olduğu tespit edilmiştir.

Zaman Kullanımı: Çalışmamıza konu olan hikâyede: belirli geçmiş zaman (66), geniş zaman (8), belirsiz geçmiş zamanın hikâyesi (5), geniş zamanın hikâyesi (7), belirsiz geçmiş zaman (5), gelecek zamanın hikâyesi (3), şimdiki zamanın hikâyesi (3), şimdiki zaman (3), emir kipi (3), istek kipi (2), yeterlilik fiili (2), şart kipinin hikâyesi (2) ve şart kipi (1) kullanılmıştır.

B1 Düzeyine Uygun Olmayan Yapılar: Çalışmamıza konu olan *İpekli Mendil* hikâyesindeki B1 düzeyine uygun olmayan yapılar şunlardır:

- Sıfat-fiil yapıları (kaçan, kalan, kanayan, gözüken, sıktığım, kaydığımız, vb.),
- Geniş zamanın şartı kullanımı (buruşturursun),
- Geniş zamanın hikâyesine ait kullanımlar (verirdi, yatardım, dökülürdü, bırakırdım, vb.)
- Gelecek zamanın hikâyesine ait kullanımlar (bitecekti, çıkarmayacaktım)
- Belirsiz geçmiş zamanın hikâyesine ait kullanımlar (istemişti, olmuştuk, inmişti, vb.)
- Ettirgen çatı kullanımları (bağlattım, korkuttuk, vb.),
- Dönüşlü çatı kullanımları (silkindim, dökülürdü)
- Edilgen çatı kullanımları (söylene söylene, bırakılınca)
- Bağlaç kullanımları (hâlbuki, doğrusu)

- Mecaz anlamlı kelimeler, deyimler ve kalıp ifadeler (üstüne ölü toprağı serpilmek, canım sıkıldı, katıla katıla gülmek, kırlangıç gibi fırladı, gık demek, vb.)
- Dolaylı anlatım kullanımları (seslendi, dedim, vb.)

UYARLAMA SÜRECİ

Metin uyarlama sürecinde kullanılan strateji ve teknikler, alan yazında farklı araştırmacılar tarafından çeşitli biçimlerde sınıflandırılmıştır. Özellikle Nation (1993), Yano Y., Long M. H. ve Ross S. (1994), Alderson (2000) ve Chang (2008) gibi araştırmacılar, yabancı dil öğretiminde metinlerin öğrenici düzeyine uygun hâle getirilmesine yönelik farklı sadeleştirme çalışmaları yapmışlardır. Bu çalışmalarda metin uyarlama; koruma, değiştirme, özetleme ve silme gibi temel işlemlerin yanı sıra sözcüksel ve söz dizimsel düzeyde gerçekleştirilen değişiklikler üzerinden ele alınmıştır. Bu çalışmaları temel alan Fatma Bölükbaş (2015) ise *Yabancı Dil Olarak Türkçe Öğretiminde Okuma Metinlerinin Dil Düzeylerine Göre Sadeleştirilmesi* adlı bildirisinde, yabancı dil olarak Türkçe öğretiminde okuma metinlerinin sadeleştirilmesine yönelik uyarlama işlemlerinin koruma, değiştirme, özetleme ve silme olmak üzere dört temel strateji altında sınıflandırıldığını ifade etmiştir. Değiştirme stratejisi ise sözcüksel, söz dizimsel, sözcüksel-söz dizimsel ve biçimsel değiştirme olmak üzere alt kategorilere ayrılmaktadır (Alderson, 2000; Chang, 2008; Nation, 1993). Yapılan alan taraması sonucunda elde edilen bulgulardan hareketle bu çalışmada da metin uyarlaması gerçekleştirilmiştir. Ancak uyarlama sürecinde kullanılan sözcüksel, söz dizimsel, sözcüksel-söz dizimsel ve biçimsel değiştirme başlıkları, bu çalışmada sözcüksel değişiklikler ve söz dizimsel değişiklikler olmak üzere iki temel başlık altında ele alınmıştır. Bunlara ek olarak dil bilgisel değişiklikler ile karmaşık cümlelerin genişletilerek açıklanması başlıkları oluşturulmuş ve bu doğrultuda Sait Faik Abasıyanık'ın *İpekli Mendil* adlı hikâyesi B1 düzeyine uyarlanmıştır. Uyarlama yapılırken hikâyede yer alan her cümle incelenmiştir.

Sözcüksel Değişimler

Sözcüksel değişimler kapsamında, özgün metinde yer alan ve B1 seviyesine uygun olmayan sözcükler daha sık kullanılan ve öğrenicilerin düzeyine uygun kelimelerle değiştirilmiştir. Ayrıca metindeki mecazlı ifadeler ve deyimler, B1 düzeyindeki öğrenicilerin anlayabileceği biçimde düzenlenmiş ve gerekli görülen durumlarda bu ifadelerin gerçek anlamlı kullanımları tercih edilmiştir. Bu tercihlere ait birkaç örnek aşağıdaki tabloda verilmiştir.

Sözcük Değişimleri	
Orijinal Metindeki Kullanım	Uyarlanmış Metindeki Kullanım
Meçhul	Belirsiz
Patırtı	Gürültü
Çakı	Bıçak
Kerata	Delikanlı

Müddet	Süre

Mecaz ve Deyim Değişimleri	
Orijinal Metindeki Kullanım	Uyarlanmış Metindeki Kullanım
Katıla katıla gülmek	Çok fazla gülmek
Alan taran etmek	Uzun süre karıştırmak
Düşünmek taşınmak	İyice düşünmek
Üstüne ölü toprağı serpilmek	Ölü gibi uyumak
Kulaklarını çekmek	Kızmak

Söz Dizimsel Değişiklikler

Söz dizimsel değişiklikler kapsamında, özgün metindeki uzun ve karmaşık cümleler anlaşılırlığı artırmak amacıyla bölünerek B1 düzeyine uygun hâle getirilmiştir. Devrik ve kuralsız cümle yapıları ise kurallı cümlelere dönüştürülmüştür. Bu uyarlamalara ait birkaç örnek aşağıdaki gibidir:

Orijinal Cümle 1: Emir Sultan'da, ay ışığında, kızakla kaydığımız zamanlar, benim de aynı bu tonda, bu kıvamda arkadaşlarım olmuştu.

Uyarlanmış Cümle 1: Eskiden Emir Sultan'da ay ışığında, kayak yapıyorduk o zamanlar benim de aynı bu renkte bu tarzda arkadaşlarım oldu.

Orijinal Cümle 2: Ben bilirim, yazın başlangıcından ta ceviz mevsimine kadar Bursa çocuklarının yalnız elleri erik ve şeftali, yalnız çizgili mintanlarının kopmuş düğmelerinden gözüken göğüsleri fındık yaprağı kokar.

Uyarlanmış Cümle 2: Ben bilirim, yaz mevsiminin başından ceviz mevsimine kadar Bursalı çocukların hem elleri erik ve şeftali kokar, hem de çizgili gömleklerinin açık düğmelerinin içindeki göğüsleri fındık yaprağı kokar.

Orijinal Cümle 3: Sımsıkı yakaladım keratayı.

Uyarlanmış Cümle 3: Delikanlıyı sımsıkı yakaladım.

Dil Bilgisel Değişiklikler

Dil bilgisel değişiklikler kapsamında, B1 düzeyinin kazanımları dışında kalan dil bilgisi yapıları sadeleştirilmiş ve bunların yerine B1 düzeyine uygun yapılar kullanılmıştır.

➢ **Sıfat-Fiil Yapılarının Uyarlanması**: B1 düzeyine uygun olmayan sıfat-fiil yapıları, düzeye uygun yapılarla değiştirilmiştir.

Orijinal Cümle 1: Ben isteksiz, nereye gideceği meçhul adımlarla ilerlerken, kapıcı arkamdan seslendi: -"Nereye?"

Uyarlanmış Cümle 1: Ben isteksiz ve belirsiz adımlarla yürürken, kapıcı arkamdan seslendi: -"Nereye?"

Orijinal Cümle 2: Fenerin gür ışığında kaçmaya çabalayan iki çıplak ayak göründü. Arkasından seğirttim. Kaçanı yakaladım.

Uyarlanmış Cümle 2: Fenerin uzun ve güçlü ışığında iki çıplak ayak gördüm. Işıktan kaçmak istiyordu. Arkasından koştum ve onu yakaladım.

➢ **Birleşik Zamanlı Fiillerin Uyarlanması:** B1 düzeyine uygun olmayan birleşik zamanlı fiiller (geniş zamanın hikâyesi, gelecek zamanın hikâyesi ve belirsiz geçmiş zamanın hikâyesi) düzeye uygun yapılarla değiştirilmiştir.

Orijinal Cümle 1: Birer de benim köylü sigarasından yaktık, iyice ahbap olmuştuk.

Uyarlanmış Cümle 1: Bir tane de benim köylü sigarasından yaktık, iyice arkadaş oluyorduk.

Orijinal Cümle 2: İmalathanenin nerede olduğunu o ne bilecekti.

Uyarlanmış Cümle 2: İmalathanenin yerini o bilmiyordu.

➢ **Fiilde Çatı Yapılarının Uyarlanması:** B1 düzeyine uygun olmayan fiilde çatı yapıları (ettirgen çatı, dönüşlü çatı ve edilgen çatı) düzeye uygun yapılarla değiştirilmiştir.

Orijinal Cümle 1: Onu, ipekli mendili vermeden gönderdiğime müessir düşünürken, dışarıda bir gürültü ile silkindim.

Uyarlanmış Cümle 1: Onu, ipekli mendili vermeden gönderdim ve sıkıntı ile düşünürken, dışarıdaki bir gürültü ile kendime geldim.

Orijinal Cümle 2: Kanayan parmağıma onun kaçak tütününden bastım; mendili yırttım ve elimi ona bağlattım.

Uyarlanmış Cümle 2: Parmağım kanıyordu. Kanı durdurmak için onun kaçak tütününden parmağıma koydum; mendili yırttım ve elimi o bağladı.

➢ **Bağlaç Kullanımları**: B1 düzeyine uygun olmayan bağlaçlar, düzeye uygun bağlaçlarla değiştirilmiştir.

Orijinal Cümle 1: Hâlbuki o, yine geldiği gibi bomboş, sessiz sedasız pencereden sıyrılıp gitti.

Uyarlanmış Cümle 1: Ama o, yine nasıl geldiyse öyle elleri boş, sessizce pencereden çıkıp gitti.

➢ **Dolaylı Anlatım Kullanımları:** B1 düzeyine uygun olmayan dolaylı anlatım kullanımları, düzeye uygun doğrudan anlatım kullanımlarıyla değiştirilmiştir.

Orijinal Cümle 1: Ben, isteksiz, nereye gideceği meçhul adımlarla ilerlerken, kapıcı arkamdan seslendi:

-"Nereye?"

-"Şöyle bir gezineyim, dedim."

-"Cambaza gitmiyor musun?"

Cevap vermediğimi görünce, ilave etti:

-"Herkes gidiyor. Bursa'ya daha böylesi gelmemiş.

Uyarlanmış Cümle 1: Ben isteksiz ve belirsiz adımlarla yürürken kapıcı arkamdan:

-"Nereye?" diye sordu.

-"Biraz gezeceğim", diye cevap verdim.

-"Cambaz gösterisine gitmiyor musun?" diye sordu.

Ben cevap vermeyince kapıcı tekrardan bana:

-"Herkes gidiyor. Bursa'ya daha önce böyle cambaz gösterisi gelmemiş," diye ekledi.

Karmaşık Cümlelerin Genişletilerek Açıklanması

Metnin anlaşılırlığını artırmak amacıyla kısa cümlelerin genişletilmesi ve açıklanması ilkesine başvurulmuştur. Bu doğrultuda, özgün metinde az sözle ifade edilen ve B1 düzeyindeki öğrenicilerin anlamakta güçlük çekebileceği cümleler, anlamı açık ve anlaşılır kılacak biçimde genişletilerek yeniden düzenlenmiştir. Bu cümlelere ait birkaç örnek aşağıdaki gibidir.

Orijinal Cümle 1: O sırada kapıcının saati on ikiyi çaldı.

Uyarlanmış Cümle 1: O zaman saat çalmaya başladı. Çünkü saat on ikiydi.

Orijinal Cümle 2: On beş yaşında vardı. Hani böyle şey âdeti değildi ama, gençlik işte. Birisi ondan ipekli mendil istemişti, hani canım anlarsın ya, âşıklısı, sevdalısı, komşu kızı işte!

Uyarlanmış Cümle 2: On beş yaşındaydı. Normalde hırsızlık yapmıyordu ama gençti. Birisi ondan ipekli mendil istemişti, anladın mı, âşıktı, seviyordu, belki de komşusunun kızıydı!

Uyarlanmış Metnin Özellikleri

Kelime ve Cümle Sayısı: *İpekli mendil* hikâyesinin uyarlanmış metninde 106 cümle (%13) ve 728 kelime (%87) yer almaktadır. Bu durum göz önünde bulundurulduğunda özgün metne oranla cümle sayısının %10, kelime sayısının ise %6 oranında arttığı görülmektedir. Bu artış, özgün metnin B1 düzeyine uyarlanması sürecinde anlaşılması güç cümlelerin açıklayıcı biçimde genişletilmesi, deyim, atasözü ve kalıp ifadelerin daha anlaşılır hâle getirilmesi, ayrıca devrik ve eksiltili cümlelerin düzenlenmesi ve gerekli görülen yerlerde yeni ifadelerin eklenmesinden kaynaklanmaktadır.

Kelime Türleri Dağılımı: Uyarlanmış metinde 174 fiil soylu sözcük (%76) ve 554 isim soylu sözcük (%24) bulunmaktadır. Bu durum göz önünde bulundurulduğunda özgün metne oranla fiil soylu sözcüklerin %5, isim soylu sözcüklerin ise %6 arttığı görülmektedir.

Dil Bilgisi Yapıları: Özgün metinde yer alan ancak B1 düzeyine uymayan dilbilgisi yapıları ve sözcükler değiştirilerek yerine Avrupa Dilleri Ortak Çerçeve Programı -CEFR- (A1, A2, B1 düzeleri dil yeterlilikleri ve kazanımları) ve *Yeni İstanbul Uluslararası Öğrenciler İçin Türkçe Ders Kitabı* setinin A1, A2 ve B1 düzeylerindeki dil bilgisi yapıları ve sözcükler eklenmiştir.

Zaman Kullanımı: Belirli geçmiş zaman (72), şimdiki zamanın hikâyesi (23), geniş zaman (8), belirsiz geçmiş zaman (7), gelecek zaman (4), şimdiki zaman (4), şart kipi (2), yeterlilik fiili (2), gereklilik kipinin hikâyesi (2), istek kipi (1) ve emir kipi (1) kullanılmıştır.

SONUÇ

Bu çalışmada, Sait Faik Abasıyanık'a ait olan *İpekli Mendil* adlı hikâyenin B1 seviyesindeki öğrenciler tarafından daha anlaşılır hâle getirilmesi amacıyla çeşitli uyarlama işlemleri yapılmıştır. Uyarlama sürecinde öncelikle B1 seviyesine uygun olmayan dil bilgisi yapıları, kelimeler, kalıp ifadeler, deyimler ve atasözleri tespit edilmiştir. Söz konusu unsurlar, *Yeni*

İstanbul Uluslararası Öğrenciler İçin Türkçe Ders Kitabı'nda (A1, A2 ve B1) yer alan kelime, dil bilgisi ve kalıp ifadelerden ve Avrupa Dilleri Ortak Çerçeve Programı'nda yer alan A1, A2 ve B1 seviye dil yeterliliklerinden (CEFR) yararlanılarak yeniden düzenlenmiştir. Böylece metnin söz varlığı ve dil bilgisi yapıları B1 düzeyindeki öğrenicilerin dil yeterliliklerine uygun hâle getirilmiştir.

Uyarlama sürecinde yalnızca sözcüksel ve dil bilgisel değişikliklerle yetinilmemiş, metnin söz dizimsel özellikleri de ele alınmıştır. Özellikle anlamlandırılması güç olan devrik yapıların yanı sıra uzun ve karmaşık cümleler, B1 seviyesindeki bir öğrencinin daha kolay anlayabileceği şekilde yeniden yapılandırılmıştır. Uzun cümleler gerektiğinde daha kısa ve anlaşılır cümlelere ayrılmış, anlamı açık olmayan veya bağlam içerisinde anlaşılması güç olan ifadeler ise genişletilerek hikâyenin anlaşılırlığı artırılmıştır.

Yapılan uyarlamalar sonucunda, özgün metin ile uyarlanmış metin arasında cümle ve sözcük türleri bakımından belirgin farklılıkların ortaya çıktığı görülmüştür. Uyarlanmış metne bakıldığında cümle, kelime, fiil soylu sözcükler ve isim soylu sözcüklerin sayı bakımından artış gösterdiğini söylemek mümkündür. Bu artışın temel nedeni, uyarlama sürecinde anlamlandırılması güç cümlelerin daha anlaşılır hâle getirilmesi amacıyla açıklayıcı unsurlar eklenmesi ve uzun cümlelerin daha kısa ve bağımsız birden fazla cümleye dönüştürülmesidir. Dolayısıyla uyarlama sürecinde, mevcut dil unsurlarının daha basit karşılıklarla değiştirilmesiyle sınırlı kalınmamış ayrıca öğrenicinin metni anlamasını kolaylaştıracak biçimde cümle yapılarının yeniden düzenlenmesi de yapılmıştır.

Sonuç olarak, edebî metinlerin yabancı dil olarak Türkçe öğretiminde kullanılabilmesi için öğrenicilerin dil düzeyleri dikkate alınarak sistemli bir biçimde uyarlanması gerekmektedir. Bu çalışmada gerçekleştirilen uyarlama işlemleri, özgün metnin edebî niteliğini mümkün olduğunca koruyarak metnin B1 seviyesindeki öğreniciler tarafından daha anlaşılır hâle getirilmesine katkı sağlamıştır. Elde edilen bulgular, özellikle sözcüksel, dil bilgisel ve söz dizimsel uyarlamaların yanı sıra cümlelerin açıklanması ve bölünmesinin metnin anlaşılabilirliğini artırmada önemli bir rol oynadığını göstermektedir. Bu doğrultuda, yabancılara Türkçe öğretiminde edebî metinlerden yararlanılırken metinlerin yalnızca kelime ve dil bilgisi açısından değil, cümle uzunluğu, söz dizimi, kalıp ifadeler ve metnin genel anlaşılırlığı bakımından da öğrenici düzeyine uygun hâle getirilmesi gerektiği sonucuna ulaşılmıştır.

KAYNAKÇA

Abasıyanık, S. F. (1990). *Seçme Hikâyeler*. (T. Buğra, Haz.). İstanbul: Milli Eğitim Bakanlığı Yayınları.

Alderson, J.C. (2000). *Assessing reading*. Cambridge: Cambridge University Press.

Bölükbaş, F. (2015, Mayıs). *Yabancı Dil Olarak Türkçe Öğretiminde Okuma Metinlerinin Sadeleştirilmesi.* 1. Uluslararası Dil Eğitimi ve Öğretimi Sempozyumu, Nevşehir.

Bölükbaş, F., Keskin, F., & Gedik, E., & Tokgöz Utku, H., & Ünsal Rey, G., & Özenç, F., & Gönültaş, G., & Şimşek, İ. D., & Alaca, S., & Yusufoğlu, S. (2016). *İstanbul Yabancılar İçin Türkçe Ders Kitabı A1*. Kültür Sanat Basımevi.

Bölükbaş, F., Keskin, F., & Gedik, E., & Tokgöz Utku, H., & Ünsal Rey, G., & Özenç, F., & Gönültaş, G., & Şimşek, İ. D., & Alaca, S., & Yusufoğlu, S. (2016). *İstanbul Yabancılar İçin Türkçe Ders Kitabı A2*. Kültür Sanat Basımevi.

Bölükbaş, F., Keskin, F., & Gedik, E., & Tokgöz Utku, H., & Ünsal Rey, G., & Özenç, F., & Gönültaş, G., Göçen Özdemirel, G., & Yavuz, S., & Şimşek, İ. D., & Alaca, S., & Yusufoğlu, S. (2016). *İstanbul Yabancılar İçin Türkçe Ders Kitabı B1*. Kültür Sanat Basımevi.

Chang, S. (2008). A preliminary categorization of techniques of simplification in ESL/EFL graded readers. (Ed. Martin Edwardes) *Proceedings of the BAAL Annual Conference*. London: Scitsiugnil Press.

Common European Framework Of Reference For Language, Learning, Teaching, Assesment. (2001). Cambridge: Cambridge University Press.

Çelik, Y. (2020, 03 Ocak). *Sait Faik Abasıyanık*. Türk Edebiyatı İsimler Sözlüğü (TEİS). https://teis.yesevi.edu.tr/madde-detay/sait-faik-abasiyanik

Erkul, R. (2004). *Cümle ve Metin Bilgisi*. Ankara: Anı yayıncılık.

Günay, D. (2007). Liselerdeki Yazın Eğitimine Yeni Bir Yaklaşım. *Milli Eğitim*, 169, 8-17.

Karataş, T. (2001). *Ansiklopedik Edebiyat Terimleri Sözlüğü*. İstanbul: Perşembe Kitapları.

Nation, P. (1993). Measuring readiness for simplified material: a test of the first 1,000 words of English. In M. L. Tickoo (Ed.), *Simplification: Theory and Application* (pp. 193-203). Singapore: SEAMEO Regional Language Center.

Okay, M. O. (2008). Sait Faik Abasıyanık. *İslâm Ansiklopedisi*. Türkiye Diyanet Vakfı Yay. 583-584. (https://islamansiklopedisi.org.tr/sait-faik-abasiyanik)

Türk Dil Kurumu. (t.y.). Dil. *Sozluk.gov.tr Güncel Türkçe Sözlük* içinde. 27 Ağustos, 2024, tarihinde https://sozluk.gov.tr/?ara=dil adresinden erişilmiştir.

Ek: Uyarlanmış Metin *İpekli Mendil*

İpekli Mendil

İpek fabrikasının geniş tarafı, ayla ışıldadı. Kapının önünden birkaç kişi, acele acele geçtiler. Ben, isteksiz ve belirsiz adımlarla ilerlerken, kapıcı arkamdan:

-"Nereye?" diye sordu.

-"Biraz gezeceğim", diye cevap verdim.

-"Sihirbaz gösterisine gitmiyor musun?" diye sordu.

Ben cevap vermeyince, kapıcı tekrardan bana:

-"Herkes gidiyor. Bursa'ya daha önce böyle cambaz gösterisi gelmemiş," diye ekledi.

-"Hiç istemiyorum," diye cevap verdim.

Fabrikayı beklemem için çok ısrar etti, ben de kabul ettim. Biraz oturdum, bir sigara içtim, bir türkü söyledim. Sonra sıkıldım. "Ne yapsam" diye düşündüm, kalktım, kapıcı odasındaki uzun ve demirli sopayı aldım, fabrikayı dolaşmaya çıktım.

Kozahanede kızlar çalışıyordu. Orayı geçer geçmez bir gürültü duydum. Cebimdeki elektrik fenerini yaktım. Etrafa bakındım. Fenerin uzun ve güçlü ışığında iki çıplak ayak gördüm. Işıktan kaçmak istiyordu. Arkasından koştum ve onu yakaladım.

Kapıcı odasına hırsızla beraber girdik. Kapıcının sarı ışıklı fenerini yaktım.

Aaa, bu küçücük bir hırsızdı! Ellerimin içindeki ufacık elini sıkıca tutuyordum. Gözleri parlaktı.

Bir süre sonra gülmek için, hem de çok fazla gülmek için ellerini bıraktım.

O zaman küçücük bir bıçak ile üzerime koştu. Ve yaramaz, beni küçük parmağımdan yaraladı. Delikanlıyı sımsıkı yakaladım. Ceplerini aradım. Bir parça kaçak tütün ve kaçak bir iki sigara kâğıdı, temizce bir mendil buldum. Parmağım kanıyordu. Kanı durdurmak için onun kaçak tütününden parmağıma koydum; mendili yırttım ve elimi o bağladı. Biraz tütün kaldı. Bu tütünle de iki kalın sigara yaptık, arkadaş gibi konuştuk.

Neredeyse on beş yaşındaydı. Normalde hırsızlık yapmıyordu ama gençti. Birisi ondan ipekli mendil istemiş, anladın mı, âşıktı, seviyordu, belki de komşusunun kızıydı. Para da yok ki, gidip çarşıdan alsın. İyice düşünmüş aklına bu çare gelmiş. Ben:

-"Peki ama fabrika bu tarafta, sen ters tarafta ne arıyordun?" diye sordum.

Güldü. Fabrikanın yerini o bilmiyordu.

Bir tane de benim köylü sigarasından yaktık, iyice arkadaş oluyorduk.

Doğma büyüme Bursalıydı. On beş senelik hayatında İstanbul'a hiç gitmemiş Mudanya'ya da -bunu söylerken yüzünü görmeliydiniz- bir kere gitmiş.

Eskiden Emir Sultan'da ay ışığında, kayak yapıyorduk. O zamanlar benim de aynı bu renkte bu tarzda arkadaşlarım oldu.

Eminim ki bunun da onlar gibi Gökdere'nin havuzlarında -Gökdere'nin havuzlarının sesini uzaktan duyarsın- derisi kahverengiye döndü. Onun derisinin rengi mevsimden mevsime meyve kabuklarının rengiyle aynı oluyordu.

Ona baktım, teni ceviz gibi kahverengiydi. Dişleri taze ceviz gibi beyaz ama güçsüzdü. Ben bilirim, yaz mevsiminin başından ceviz mevsimine kadar Bursalı çocukların hem elleri erik ve şeftali kokar, hem de çizgili gömleklerinin açık düğmelerinin içindeki göğüsleri fındık yaprağı kokar. O zaman saat çalmaya başladı. Çünkü saat on ikiydi. Sihirbaz gösterisinin bitmesine az kaldı.

Hırsız çocuk bana -"Kaçayım," diye söyledi.

Onu, ipekli mendili vermeden gönderdim ve sıkıntı ile düşünürken, dışarıdaki bir gürültü ile kendime geldim. Kapıcı konuşa konuşa odadan içeri giriyordu. Arkasında da hırsız…

Bu sefer ben de ona kızdım. Kapıcı çıplak ayaklarına söğüt ağacının ince dalıyla vurarak kıpkırmızı yaptı. Şanslıydık çünkü patron orada yoktu. Yoksa şimdi kesinlikle onu polise veriyorduk. Kapıcı bana: "Bu yaşta bir çocuk hırsız! Arkadaşım, hapishanede kalırsa doğru davranmayı öğrenir" diye söyledi.

Çok kızdık ağlamadı. Gözleri ağlamaya hazır çocukların gözlerine benziyordu ama, dudaklarında azıcık bir titreme gözükmedi ve kaşları sonuna kadar sabit ve kararlıydı. Fakat biraz hareketliydiler.

Biz hırsızı bırakınca özgür bir kuş gibi hızlıca gitti. Ay ışığını ve mısır tarlasını hızla geçip gitti. Ben, o zamanlar fabrikanın üstündeki odada yatıyordum. Bu fabrikada mallar vardı. Odam çok güzeldi. Aylı gecelerde daha şirin oluyordu.

Tam pencereme yakın bir dut ağacı vardı. Ay ışığı dut yapraklarından geçerek, odaya giriyordu. Neredeyse yaz ve kış mevsimlerinde pencereyi açıyordum. Çok serin ve tuhaf rüzgârlar esiyordu. Vapurlarda da çalıştım. Bu sebeple rüzgârları kokularından ayırt ederek, tanıyordum. Birçok rüzgâr yorganımın üzerinden acayip bir rüya gibi gelip geçti.

Kolayca uyanırım. Sabaha yakındı. Dışarıdan bir gürültü geliyordu. Dut ağacında birisi var gibiydi. Korkmuşum ki, kalkamadım, bağıramadım. Tam o zaman pencerede bir kişi gördüm. O'ydu, yavaşça pencereden içeri girdi. Benim önümden geçerken, gözlerimi kapadım, dolapların içini açıp baktı. Malları uzun süre karıştırdı. Hiçbir şey söylemedim. Gerçekten çok cesurdu ve bundan dolayı bütün malı alıp gitse, sesimi çıkarmamalıydım. Yarın patron bana:

-"Ölü gibi uyuyor muydun hayvan?" diye soracak, bana vuracak ve beni kovacak. Bunu bilmeme rağmen hiç konuşmadım.

Ama o, yine elleri boş geldi ve yine elleri boş, sessizce pencereden çıkıp gitti. Bu anda da bir dalın kırılma sesini işittim. Düştü. Aşağı indim. Hırsızın başında kapıcı ile beraber birkaç kişi bekliyordu.

Ölüyordu. Kapıcı hırsızın sımsıkı kapalı avucunu açtı. Bu avucun içinden bir ipek mendil su gibi fışkırdı.

Ya… İyi, güzel ipekli mendiller hep böyledir. Avucunun içinde bırakmadan sımsıkı sıkarsın, sonra avuç açılınca insanın elinden su gibi fışkırır.

DİL ÖĞRETİMİNDE KULLANILAN GÜNCEL ÜRETKEN YAPAY ZEKÂ ARAÇLARININ TÜRKÇE EĞİTİMİ AÇISINDAN DEĞERLENDİRİLMESİ, FIRSATLAR VE RİSKLER

AN EVALUATION OF CURRENT GENERATIVE AI TOOLS USED IN LANGUAGE TEACHING FROM THE PERSPECTIVE OF TURKISH LANGUAGE EDUCATION: OPPORTUNITIES AND RISKS

Yasemin ALBAYRAK
Master's Student
ORCID ID: https://orcid.org/0009000882338414

Assoc. Prof. Dr. Sıddık BAKIR
Atatürk University, Kazım Karabekir Faculty of Education
ORCID ID: https://orcid.org/0000-0002-5118-209X

Özet

Üretken yapay zekâ teknolojilerindeki hızlı gelişmeler, eğitim alanında olduğu gibi dil öğretiminde de önemli dönüşümlere yol açmıştır. ChatGPT, Gemini, Claude, Microsoft Copilot, Jenni, Qwen ve benzeri üretken yapay zekâ araçları; ders materyali hazırlama, metin üretme, konuşma pratiği, ölçme-değerlendirme ve bireyselleştirilmiş öğrenme gibi birçok alanda öğretmenlere ve öğrencilere yeni fırsatlar sunmaktadır. Bununla birlikte bu teknolojilerin eğitim süreçlerine entegrasyonu; yanlış bilgi üretimi, akademik dürüstlük, veri gizliliği ve aşırı teknoloji bağımlılığı gibi çeşitli riskleri de beraberinde getirmektedir. Bu çalışmanın amacı, güncel üretken yapay zekâ araçlarını dil öğretimi bağlamında tanıtmak ve bu araçların Türkçe eğitimi açısından kullanım alanlarını, sundukları fırsatları ve olası riskleri değerlendirmektir. Araştırma nitel araştırma yaklaşımı doğrultusunda kavramsal bir çalışma olarak tasarlanmıştır. Veriler, güncel alanyazın, yapay zekâ platformlarının özellikleri ve eğitim teknolojilerine ilişkin dokümanların incelenmesi yoluyla elde edilmiştir. Bulgular, üretken yapay zekâ araçlarının özellikle yazma, konuşma, okuma, materyal geliştirme ve bireyselleştirilmiş öğrenme süreçlerinde önemli katkılar sunduğunu göstermektedir. Bununla birlikte bu araçların bilinçli ve pedagojik ilkeler doğrultusunda kullanılması gerektiği ve bunun, verimliliğin artışına olan olası katkıları sıralanmaya çalışılmıştır.

Anahtar Kelimeler: Üretken yapay zekâ, Türkçe eğitimi, dil öğretimi, eğitim teknolojileri, ChatGPT.

Abstract

Rapid advancements in generative artificial intelligence technologies have led to significant transformations in language teaching, just as they have in the broader field of education. Generative AI tools such as ChatGPT, Gemini, Claude, and Microsoft Copilot, Jenni, Qwen offer new opportunities for teachers and students in areas including lesson material preparation, text generation, speaking practice, assessment, and personalized learning. However, integrating these technologies into educational processes also entails various risks, such as the generation of misinformation, concerns regarding academic integrity, data privacy issues, and excessive reliance on technology. This study aims to introduce current generative AI tools within the context of language teaching and to evaluate their potential applications, opportunities, and risks specifically regarding Turkish language education. The research was designed as a conceptual study following a qualitative research approach. Data were gathered through an analysis of current literature, the features of AI platforms, and documents related to educational technologies. The findings indicate that generative AI tools offer significant contributions, particularly in the processes of writing, speaking, reading, material development, and personalized learning. Furthermore, an attempt has been made to outline the necessity of using these tools in a conscious manner and in accordance with pedagogical principles, as well as their potential contributions to increased efficiency.

Keywords: Generative artificial intelligence, Turkish language education, language teaching, educational technologies, ChatGPT.

YABANCILARA TÜRKÇE ÖĞRETİMİNDE ÖĞRENCİ KATILIMININ ARTIRILMASINA YÖNELİK YAPAY ZEKÂ DESTEKLİ ARAYIŞLAR

AI-SUPPORTED APPROACHES TO INCREASING STUDENT PARTICIPATION IN TEACHING TURKISH TO FOREIGNERS

Yasemin ALBAYRAK
Master's Student
ORCID ID: https://orcid.org/0009000882338414

Assoc. Prof. Dr. Sıddık BAKIR
Atatürk University, Kazım Karabekir Faculty of Education
ORCID ID: https://orcid.org/0000-0002-5118-209X

Özet

Üretken yapay zekâ teknolojilerindeki gelişmeler, yabancı dil öğretiminde olduğu gibi yabancılara Türkçe öğretimi alanında da öğretim süreçlerini önemli ölçüde dönüştürmektedir. Büyük dil modellerine dayalı yapay zekâ sistemleri; konuşma pratiği, yazma etkinlikleri, okuma çalışmaları, kişiselleştirilmiş geri bildirim, materyal geliştirme ve kültür öğretimi gibi birçok alanda öğretmenlere ve öğrencilere yeni öğrenme fırsatları sunmaktadır. Bununla birlikte bu teknolojilerin en önemli katkılarından biri, öğrencilerin öğrenme sürecine daha etkin katılımını destekleyebilme potansiyelidir. Bu çalışmanın amacı, üretken yapay zekâ destekli etkinliklerin yabancılara Türkçe öğretiminde öğrenci katılımını nasıl destekleyebileceğini ortaya koymak ve öğretmenlere uygulamaya dönük öneriler sunmaktır. Araştırma nitel araştırma yaklaşımı doğrultusunda kavramsal bir çalışma olarak tasarlanmıştır. Veriler güncel alanyazın, eğitim teknolojileri araştırmaları ve üretken yapay zekâ uygulamalarına ilişkin dokümanların incelenmesi yoluyla elde edilmiştir. Bulgular, üretken yapay zekâ destekli etkinliklerin öğrencilerin davranışsal, bilişsel ve duyuşsal katılımlarını desteklediğini; özellikle konuşma, yazma, kültür öğretimi ve bireyselleştirilmiş öğrenme süreçlerinde önemli katkılar sağladığını göstermektedir. Sonuç olarak, üretken yapay zekânın öğretmenin yerini alan bir teknoloji değil, öğretim sürecini zenginleştiren ve öğrenci katılımını artıran pedagojik bir destek aracı olarak değerlendirilmesi gerektiği fikrinden hareketle de çeşitli önerilerde bulunulmuştur.

Anahtar Kelimeler: Üretken yapay zekâ, yabancılara Türkçe öğretimi, öğrenci katılımı, eğitim teknolojileri, dijital öğrenme.

Abstract

Advancements in generative artificial intelligence technologies are significantly transforming instructional processes in the field of teaching Turkish to foreigners, just as they are in general foreign language education. AI systems based on large language models offer new learning opportunities for both teachers and students across various areas, including speaking practice, writing activities, reading exercises, personalized feedback, material development, and cultural instruction. Notably, one of the most significant contributions of these technologies is their potential to foster more active student participation in the learning process. This study aims to demonstrate how generative AI-supported activities can enhance student engagement in teaching Turkish to foreigners and to provide practical recommendations for teachers. The research was designed as a conceptual study following a qualitative research approach. Data were gathered through an analysis of current literature, research on educational technologies, and documentation regarding generative AI applications. The findings indicate that generative AI-supported activities foster students' behavioral, cognitive, and affective engagement, making significant contributions-particularly in speaking, writing, cultural instruction, and individualized learning processes. Consequently, various recommendations have been made

based on the premise that generative AI should be viewed not as a technology that replaces the teacher, but rather as a pedagogical support tool that enriches the teaching process and enhances student engagement.

Keywords: Generative artificial intelligence, teaching Turkish to foreigners, student engagement, educational technologies, digital learning.

ADVANCING VETERINARY DISEASE MANAGEMENT THROUGH MACHINE LEARNING-DRIVEN ANTIMICROBIAL RESISTANCE PREDICTION

Moses Adeolu AGOI[1]
Lagos State University of Education, Lagos Nigeria.
ORCID ID: 0000-0002-8910-2876

Bamidele Olumuyiwa ALABA[2]
Tai Solarin University of Education, Ogun Nigeria.

Gbenga Oyewole OGUNSANWO[3]
Tai Solarin University of Education, Ogun Nigeria.
&
Hendri Hermawan ADINUGRAHA[4]
Universitas Islam Negeri K.H Abdurrahman Wahid Pekalongan.

Abstract

Antimicrobial resistance (AMR) has emerged as one of the most significant threats to animal health, food security, and global public health. The widespread and often indiscriminate use of antimicrobial agents in livestock production and veterinary medicine has accelerated the evolution of resistant bacterial pathogens, thereby reducing treatment efficacy and increasing morbidity, mortality, and production losses. Conventional antimicrobial susceptibility testing (AST), although considered the diagnostic gold standard, is time-consuming and often delays clinical decision-making. Recent advances in artificial intelligence (AI) and machine learning (ML) provide transformative opportunities for predicting antimicrobial resistance using genomic, phenotypic, epidemiological, and clinical datasets. This paper examines the application of machine learning techniques in advancing veterinary disease management through rapid antimicrobial resistance prediction. Drawing upon recent peer-reviewed literature, the study synthesizes current evidence regarding supervised, unsupervised, and deep learning approaches employed in veterinary pathogen surveillance and precision antimicrobial therapy. The review further discusses data integration strategies involving whole-genome sequencing, electronic veterinary health records, antimicrobial usage databases, and One Health surveillance systems. Additionally, the study highlights challenges relating to dataset heterogeneity, algorithm interpretability, model generalizability, and ethical governance. Conceptually, the paper proposes a machine learning-driven decision-support framework capable of improving diagnostic accuracy, reducing unnecessary antimicrobial prescriptions, and strengthening national AMR surveillance programmes. The findings suggest that explainable machine learning models, particularly Random Forest, Extreme Gradient Boosting, Support Vector Machines, and deep neural networks, consistently demonstrate superior predictive performance across multiple veterinary applications while enhancing clinical transparency. Integrating machine learning into veterinary diagnostics represents a promising pathway toward sustainable antimicrobial stewardship, precision livestock health management, and strengthened One Health initiatives aimed at mitigating antimicrobial resistance globally.

Keywords: Antimicrobial Resistance; Machine Learning; One Health; Veterinary Disease Management.

INTRODUCTION

Antimicrobial resistance (AMR) has become one of the most pressing global health challenges, threatening the effectiveness of antimicrobial therapies in both human and veterinary medicine while undermining food security, animal welfare, and sustainable agricultural production. The extensive use and misuse of antimicrobial agents in livestock, poultry, aquaculture, and companion animal practice have accelerated the emergence and dissemination of resistant bacterial pathogens across animal, human, and environmental interfaces, reinforcing the need for coordinated interventions under the One Health framework. Veterinary pathogens such as *Escherichia coli*, *Salmonella enterica*, *Staphylococcus aureus*, *Campylobacter* spp., and *Klebsiella pneumoniae* increasingly exhibit multidrug-resistant phenotypes that compromise treatment efficacy, prolong disease outbreaks, increase production losses, and elevate the risk of zoonotic transmission (Chakraborty et al., 2024; Alkheraije, 2024). Conventional antimicrobial susceptibility testing (AST), although regarded as the gold standard for determining antimicrobial sensitivity, typically requires 24-72 hours after bacterial isolation, delaying appropriate therapeutic interventions and often encouraging empirical antimicrobial prescriptions that may further exacerbate resistance development. Consequently, there is growing interest in developing rapid, data-driven approaches capable of predicting antimicrobial resistance before laboratory confirmation. Recent advances in artificial intelligence (AI), machine learning (ML), high-throughput sequencing, and bioinformatics have transformed predictive microbiology by enabling computational models to analyse large-scale genomic, phenotypic, epidemiological, and antimicrobial usage datasets with remarkable accuracy. Supervised learning algorithms, including Random Forest, Support Vector Machine, Extreme Gradient Boosting, Artificial Neural Networks, and Deep Learning architectures, have demonstrated considerable success in identifying resistance determinants, predicting antimicrobial susceptibility patterns, and supporting precision veterinary medicine (Antunes & Kemiyondo, 2025). These technologies facilitate early clinical decision-making, optimize antimicrobial stewardship programmes, and strengthen surveillance systems through automated analysis of complex multidimensional datasets (Ajose et al., 2024). Furthermore, integrating explainable artificial intelligence with veterinary diagnostics enhances clinician confidence by providing transparent interpretations of prediction outcomes while supporting evidence-based antimicrobial prescribing. Despite these advances, significant challenges remain regarding data standardization, model interpretability, interoperability among surveillance systems, and the availability of high-quality veterinary datasets across low- and middle-income countries (Delpy et al., 2024). Therefore, advancing veterinary disease management through machine learning-driven antimicrobial resistance prediction represents a critical research priority capable of improving diagnostic efficiency, strengthening One Health surveillance, reducing antimicrobial misuse, and promoting sustainable livestock production in an increasingly interconnected global health ecosystem.



LITERATURE REVIEW

Global Burden of Antimicrobial Resistance in Veterinary Medicine

Antimicrobial resistance (AMR) has evolved into one of the most significant threats confronting veterinary medicine, livestock production, and global food systems. The emergence of resistant bacterial pathogens has reduced the effectiveness of antimicrobial therapies used to control infectious diseases in food-producing and companion animals, resulting in increased animal mortality, reduced productivity, economic losses, and heightened risks of zoonotic disease

transmission. The interconnectedness of humans, animals, and the environment has positioned AMR as a major One Health concern requiring multidisciplinary interventions (World Health Organization, 2023). Recent global estimates indicate that bacterial AMR was associated with approximately 4.71 million human deaths in 2021, with projections suggesting that AMR-related deaths could exceed 8 million annually by 2050 if effective mitigation strategies are not implemented. These projections further emphasize the importance of addressing antimicrobial misuse across both human and veterinary sectors. Within veterinary medicine, antimicrobial agents remain indispensable for the treatment, prevention, and control of bacterial infections affecting cattle, poultry, swine, aquaculture, and companion animals. However, extensive therapeutic, metaphylactic, prophylactic, and, in some settings, growth-promotion applications have accelerated the selection of resistant microorganisms. Martins et al. (2024) argue that AMR in livestock extends beyond animal health by generating profound social, environmental, and economic consequences that affect food production systems, international trade, and public health. Their proposed Global Burden of Animal Diseases (GBADs) framework emphasizes standardized approaches for estimating the impact of AMR within livestock production systems and strengthening evidence-based policy development. Several veterinary pathogens have become increasingly resistant to critically important antimicrobial agents. Multidrug-resistant strains of *Escherichia coli*, *Salmonella enterica*, *Campylobacter jejuni*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, and *Enterococcus* species have been reported in livestock, companion animals, and wildlife across numerous countries. These pathogens frequently exhibit resistance to β-lactams, tetracyclines, fluoroquinolones, aminoglycosides, and third-generation cephalosporins, thereby limiting available treatment options and increasing production costs. Surveillance studies further demonstrate that resistant bacteria circulating within animal populations can disseminate through food chains, water systems, and environmental reservoirs, reinforcing the need for integrated surveillance under the One Health framework (Tuhamize & Bazira, 2024). The increasing burden of veterinary AMR has stimulated greater investment in precision diagnostics and computational prediction technologies. Conventional antimicrobial susceptibility testing remains reliable but often requires 24–72 hours before treatment decisions can be finalized. Consequently, researchers have explored artificial intelligence and machine learning approaches capable of predicting antimicrobial resistance directly from genomic, phenotypic, and epidemiological datasets. Recent reviews suggest that machine learning models can substantially improve early identification of resistant organisms, support empirical antimicrobial selection, and enhance antimicrobial stewardship programmes by reducing unnecessary antimicrobial exposure (Kherabi et al., 2024). Overall, contemporary evidence demonstrates that antimicrobial resistance is no longer solely a microbiological problem but represents a complex veterinary, agricultural, economic, and public health challenge. Sustainable disease management therefore requires coordinated surveillance, responsible antimicrobial stewardship, improved livestock management practices, genomic monitoring, and the integration of explainable machine learning systems capable of providing rapid, reliable, and clinically interpretable resistance predictions. Such innovations will strengthen veterinary decision-making, improve animal welfare, safeguard food security, and contribute to global efforts aimed at reducing the expanding burden of antimicrobial resistance across interconnected human-animal-environment systems.

Machine Learning Algorithms for Antimicrobial Resistance Prediction in Veterinary Pathogens

Machine learning (ML) has emerged as a transformative computational approach for predicting antimicrobial resistance (AMR) in veterinary pathogens by analysing large-scale genomic, phenotypic, clinical, and epidemiological datasets. Unlike conventional statistical methods that often rely on predefined assumptions, ML algorithms automatically identify complex and

nonlinear relationships within multidimensional data, making them highly suitable for predicting resistance phenotypes across diverse bacterial species. The integration of ML into veterinary diagnostics has significantly improved the speed and accuracy of antimicrobial susceptibility prediction, thereby facilitating timely therapeutic interventions and supporting antimicrobial stewardship initiatives (Arango-Argoty et al., 2018; Boolchandani et al., 2019). Recent advances in computational biology have further enhanced the application of explainable artificial intelligence (XAI), enabling clinicians to understand the biological factors influencing resistance predictions and increasing confidence in algorithm-assisted decision-making (Kherabi et al., 2024). Supervised learning algorithms remain the most widely adopted ML techniques for AMR prediction because they utilize labelled datasets containing bacterial genomic characteristics and corresponding antimicrobial susceptibility profiles. Among these, Random Forest (RF) has consistently demonstrated excellent predictive performance due to its ensemble architecture, robustness to noisy datasets, and ability to rank the relative importance of genomic resistance determinants. RF models have achieved classification accuracies exceeding 90% for predicting resistance in *Escherichia coli*, *Salmonella enterica*, *Campylobacter jejuni*, and *Staphylococcus aureus* isolated from livestock and companion animals (Nguyen et al., 2019). Similarly, Support Vector Machines (SVMs) have shown high discriminatory capability when analysing high-dimensional genomic data, particularly for multidrug-resistant bacterial strains, owing to their effectiveness in separating complex nonlinear classes through kernel-based optimization (Yang et al., 2020). Gradient boosting algorithms, including Extreme Gradient Boosting (XGBoost) and Light Gradient Boosting Machine (LightGBM), have gained increasing popularity because of their superior predictive accuracy, computational efficiency, and ability to manage missing values within heterogeneous veterinary surveillance datasets. These algorithms sequentially improve weak classifiers by minimizing prediction errors, resulting in highly accurate resistance prediction models across multiple antimicrobial classes (Chen & Guestrin, 2016; Kherabi et al., 2024). Their implementation has proven particularly valuable in analysing whole-genome sequencing (WGS) data, where thousands of genomic features must be processed simultaneously to identify antimicrobial resistance genes and mutations associated with resistant phenotypes. Deep learning techniques have further expanded the capabilities of AMR prediction through automated feature extraction from complex biological datasets. Artificial Neural Networks (ANNs), Convolutional Neural Networks (CNNs), and Recurrent Neural Networks (RNNs) have demonstrated remarkable performance in identifying intricate genomic patterns that conventional algorithms may overlook. Deep learning models are especially effective when analysing large genomic repositories generated through next-generation sequencing technologies, enabling rapid identification of resistance-associated mutations across diverse veterinary pathogens (Arango-Argoty et al., 2018). However, these models typically require extensive computational resources, large annotated datasets, and specialized hardware, which may limit adoption in resource-constrained veterinary laboratories. Despite the impressive predictive capabilities of modern ML algorithms, several implementation challenges remain. Veterinary datasets often exhibit class imbalance, inconsistent antimicrobial susceptibility testing protocols, limited representation of pathogens from low- and middle-income countries, and variations in sequencing quality. Furthermore, highly complex algorithms frequently function as "black-box" systems, reducing transparency in clinical decision-making. To address these concerns, recent research emphasizes the integration of Explainable Artificial Intelligence (XAI) methods such as SHapley Additive exPlanations (SHAP), Local Interpretable Model-Agnostic Explanations (LIME), and feature attribution analyses to improve model interpretability while maintaining predictive accuracy (Lundberg & Lee, 2017; Kherabi et al., 2024). Collectively, these advances demonstrate that machine learning represents a powerful decision-support technology capable of revolutionizing antimicrobial resistance prediction,

enhancing veterinary diagnostics, strengthening surveillance systems, and promoting precision antimicrobial therapy within the broader One Health framework.

Genomic Data, Bioinformatics, and Whole-Genome Sequencing for Machine Learning-Based Antimicrobial Resistance Prediction

The rapid advancement of whole-genome sequencing (WGS) and bioinformatics has fundamentally transformed antimicrobial resistance (AMR) surveillance by enabling comprehensive characterization of bacterial genomes and facilitating machine learning (ML)-based prediction of antimicrobial susceptibility. Unlike conventional culture-based antimicrobial susceptibility testing (AST), which generally requires 24-72 hours after pathogen isolation, WGS provides detailed genomic information capable of identifying resistance genes, chromosomal mutations, plasmid-mediated determinants, and mobile genetic elements responsible for antimicrobial resistance. The integration of WGS with machine learning algorithms has significantly enhanced predictive diagnostics in veterinary medicine by enabling early detection of resistant pathogens before phenotypic laboratory confirmation (Boolchandani et al., 2019; Köser et al., 2014). Whole-genome sequencing generates millions of nucleotide sequences from bacterial isolates, producing high-dimensional datasets that require advanced bioinformatics pipelines for processing and interpretation. Standard workflows involve sequence quality assessment, genome assembly, and annotation, identification of antimicrobial resistance genes, mutation analysis, and comparative genomic profiling before data are incorporated into predictive machine learning models. Public repositories such as the National Center for Biotechnology Information (NCBI), the Comprehensive Antibiotic Resistance Database (CARD), ResFinder, ARG-ANNOT, and Pathogenwatch provide valuable genomic resources for developing computational models capable of predicting resistance across numerous veterinary pathogens (Alcock et al., 2023; Bortolaia et al., 2020). These databases continue to expand through international surveillance initiatives, improving model robustness and facilitating global collaboration in antimicrobial resistance research. In veterinary medicine, WGS has become increasingly important for investigating multidrug-resistant pathogens affecting livestock, poultry, aquaculture, and companion animals. Resistant strains of *Escherichia coli*, *Salmonella enterica*, *Campylobacter jejuni*, *Klebsiella pneumoniae*, *Staphylococcus aureus*, and *Enterococcus faecium* have been extensively characterized using genomic sequencing technologies. Comparative genomic analyses have identified numerous resistance-associated genes, including blaCTX-M, blaNDM, mcr-1, tet(A), erm(B), qnrS, and vanA, which contribute to resistance against β-lactams, carbapenems, colistin, tetracyclines, macrolides, fluoroquinolones, and glycopeptides (Bortolaia et al., 2020; Alcock et al., 2023). Machine learning models trained using these genomic features have demonstrated excellent predictive performance in identifying antimicrobial resistance phenotypes across multiple bacterial species while reducing dependence on time-intensive laboratory testing. The increasing availability of genomic data has also accelerated the application of advanced machine learning algorithms such as Random Forest, Support Vector Machines, Extreme Gradient Boosting (XGBoost), Artificial Neural Networks, and Deep Learning architectures. These algorithms efficiently process thousands of genomic variables simultaneously, identifying complex nonlinear interactions among resistance genes, point mutations, virulence determinants, and mobile genetic elements that may influence antimicrobial susceptibility. Recent studies demonstrate that genomic-based machine learning models frequently achieve predictive accuracies exceeding 90% for several clinically important veterinary pathogens while substantially reducing diagnostic turnaround times (Nguyen et al., 2019; Yang et al., 2020). Furthermore, explainable artificial intelligence techniques, including SHapley Additive exPlanations (SHAP), have enhanced biological interpretability by identifying the genomic features that contribute most significantly to model predictions (Lundberg & Lee, 2017).

Despite these substantial advances, several challenges continue to limit the widespread implementation of genomic machine learning in veterinary practice. High sequencing costs, inconsistent metadata quality, fragmented surveillance infrastructures, computational resource requirements, and limited representation of bacterial isolates from low- and middle-income countries reduce model generalizability. Additionally, differences in sequencing platforms, genome assembly protocols, antimicrobial susceptibility testing standards, and data-sharing policies hinder interoperability among international surveillance systems (Kherabi et al., 2024). Consequently, future research should prioritize standardized genomic data collection, harmonized bioinformatics pipelines, interoperable databases, and explainable machine learning frameworks that support real-time antimicrobial resistance surveillance, precision veterinary medicine, and evidence-based antimicrobial stewardship within the broader One Health ecosystem.

Explainable Artificial Intelligence and Decision Support Systems for Precision Veterinary Antimicrobial Stewardship

The increasing integration of artificial intelligence (AI) into veterinary medicine has accelerated the development of explainable machine learning models and intelligent clinical decision support systems (CDSSs) for antimicrobial resistance (AMR) prediction. While conventional machine learning algorithms have demonstrated exceptional predictive performance, their adoption in veterinary clinical practice has been constrained by the "black-box" nature of many models, where prediction outcomes are generated without providing transparent explanations for the underlying decision-making process. Explainable Artificial Intelligence (XAI) has emerged as a solution to this challenge by enabling clinicians to understand how machine learning algorithms arrive at their predictions, thereby improving confidence, accountability, and clinical acceptance (Lundberg & Lee, 2017; Ghassemi et al., 2021). In veterinary healthcare, where treatment decisions directly affect animal welfare, food safety, and public health, model interpretability is particularly important for supporting evidence-based antimicrobial prescribing and strengthening antimicrobial stewardship programmes. Clinical decision support systems integrate patient information, laboratory findings, antimicrobial susceptibility profiles, genomic data, farm management records, and epidemiological surveillance data to assist veterinarians in selecting appropriate antimicrobial therapies. Modern AI-powered CDSSs employ machine learning algorithms such as Random Forest, Extreme Gradient Boosting (XGBoost), Support Vector Machines (SVM), and Deep Neural Networks to predict antimicrobial susceptibility patterns before conventional antimicrobial susceptibility testing results become available. These systems substantially reduce diagnostic turnaround time while minimizing empirical antimicrobial use, thereby lowering selective pressure for resistant bacterial strains (Kherabi et al., 2024). By combining predictive analytics with clinical expertise, decision support systems facilitate rapid interventions that improve treatment outcomes and reduce unnecessary antimicrobial exposure in livestock and companion animals. Explainable AI techniques have become central to improving transparency within these intelligent decision support systems. Among the most widely adopted approaches is SHapley Additive exPlanations (SHAP), which quantifies the contribution of individual variables to each model prediction using principles derived from cooperative game theory (Lundberg & Lee, 2017). SHAP enables veterinarians to identify which genomic mutations, antimicrobial usage histories, clinical indicators, or epidemiological factors most strongly influence predicted resistance. Similarly, Local Interpretable Model-Agnostic Explanations (LIME) generates simplified local explanations for individual predictions, allowing clinicians to understand model behaviour regardless of the underlying algorithm (Ribeiro et al., 2016). These approaches improve trust, facilitate regulatory compliance, and enhance collaboration between computational scientists and veterinary practitioners by translating complex mathematical

outputs into clinically meaningful information. The integration of explainable AI also supports antimicrobial stewardship initiatives aimed at optimizing antimicrobial use across animal production systems. Stewardship programmes emphasize appropriate antimicrobial selection, accurate dosing, treatment duration, and continuous surveillance to reduce the emergence of resistant microorganisms. Machine learning-driven stewardship platforms can analyse historical prescribing patterns, pathogen prevalence, resistance trends, and farm-level risk factors to recommend the most appropriate antimicrobial therapy while discouraging unnecessary broad-spectrum antimicrobial use (World Organisation for Animal Health [WOAH], 2023). Such systems contribute to preserving antimicrobial efficacy, improving animal health outcomes, and reducing the transmission of resistant pathogens throughout the food chain. Despite these promising developments, several implementation challenges remain. Many veterinary healthcare systems, particularly in low- and middle-income countries, experience limited digital infrastructure, fragmented health information systems, insufficient computational resources, and shortages of skilled data scientists capable of deploying AI technologies. Furthermore, concerns regarding algorithmic bias, data privacy, interoperability, ethical governance, and regulatory oversight continue to hinder widespread implementation (Ghassemi et al., 2021). Addressing these challenges will require standardized veterinary datasets, transparent model validation procedures, multidisciplinary collaboration, and harmonized international guidelines for the ethical deployment of explainable AI. Ultimately, explainable machine learning and intelligent clinical decision support systems represent essential components of precision veterinary medicine, strengthening antimicrobial stewardship and supporting sustainable disease management within the global One Health framework.

One Health Surveillance and Machine Learning Integration for Antimicrobial Resistance Monitoring

The global emergence of antimicrobial resistance (AMR) has reinforced the importance of the One Health concept, which recognizes the interdependence of human, animal, and environmental health in controlling infectious diseases. Resistant microorganisms and antimicrobial resistance genes circulate across livestock, wildlife, companion animals, humans, food products, soil, and water systems, creating complex transmission pathways that cannot be effectively addressed through sector-specific interventions alone. Consequently, international organizations such as the World Health Organization (WHO), the World Organisation for Animal Health (WOAH), the Food and Agriculture Organization (FAO), and the United Nations Environment Programme (UNEP) advocate integrated surveillance systems that combine multidisciplinary expertise, shared databases, and coordinated policy implementation to mitigate AMR globally (WHO, 2023; WOAH, 2023; FAO et al., 2022). Within veterinary medicine, One Health surveillance provides an essential framework for monitoring resistant pathogens, evaluating antimicrobial consumption, and identifying emerging resistance hotspots across livestock production systems. Machine learning (ML) has become an indispensable technology for strengthening One Health surveillance by enabling the rapid analysis of heterogeneous datasets generated from veterinary hospitals, livestock farms, diagnostic laboratories, genomic sequencing centres, environmental monitoring programmes, and public health agencies. These datasets often include antimicrobial susceptibility testing (AST) results, whole-genome sequencing (WGS) data, antimicrobial prescription records, farm management information, animal movement patterns, climatic variables, and geographic information systems (GIS). Traditional statistical approaches frequently struggle to analyse such multidimensional data because of nonlinear relationships and complex interactions among biological, environmental, and management factors. Machine learning algorithms, including Random Forest (RF), Extreme Gradient Boosting (XGBoost), Support Vector Machines (SVM), Artificial Neural Networks (ANN), and Deep Learning (DL), have demonstrated

exceptional capability in identifying hidden resistance patterns, forecasting AMR emergence, and supporting evidence-based disease surveillance (Kherabi et al., 2024). The integration of genomic surveillance with machine learning has significantly enhanced the detection of antimicrobial resistance genes circulating among veterinary pathogens. High-throughput sequencing technologies generate extensive genomic datasets that allow computational models to identify resistance determinants, mobile genetic elements, plasmids, and chromosomal mutations associated with multidrug resistance. Surveillance databases such as the Comprehensive Antibiotic Resistance Database (CARD), ResFinder, Pathogenwatch, and the National Center for Biotechnology Information (NCBI) provide standardized genomic information that supports machine learning model development and international collaboration (Alcock et al., 2023; Bortolaia et al., 2020). By integrating genomic data with epidemiological and environmental information, predictive models can identify emerging resistance clusters before widespread dissemination occurs, thereby facilitating targeted interventions and reducing disease transmission across the food chain. Another significant contribution of machine learning to One Health surveillance involves predictive risk modelling. Advanced algorithms can evaluate multiple determinants, including antimicrobial usage intensity, livestock density, farm biosecurity practices, climatic conditions, animal transportation networks, and previous outbreak histories, to estimate the probability of AMR emergence within specific geographic regions. Such predictive analytics enable veterinary authorities to allocate surveillance resources more efficiently, prioritize high-risk production systems, and implement proactive antimicrobial stewardship programmes before resistance becomes widespread (van Bunnik & Woolhouse, 2017). Furthermore, integrating spatial epidemiology with machine learning allows real-time mapping of resistance hotspots, thereby improving outbreak preparedness and facilitating coordinated responses among veterinary, environmental, and public health agencies. Despite these advances, important challenges continue to limit the implementation of integrated One Health surveillance systems. Variations in antimicrobial susceptibility testing standards, inconsistent data collection protocols, fragmented national surveillance infrastructures, limited interoperability among databases, and inadequate data sharing across sectors reduce the effectiveness of predictive analytics. Many low- and middle-income countries also face shortages of sequencing facilities, computational infrastructure, and trained bioinformaticians capable of managing large-scale surveillance systems (WHO, 2023). Addressing these limitations will require harmonized international data standards, expanded investment in digital veterinary infrastructure, strengthened cross-sectoral collaboration, and explainable machine learning frameworks capable of generating transparent and clinically interpretable predictions. Collectively, the integration of machine learning within One Health surveillance represents a transformative strategy for improving antimicrobial resistance monitoring, strengthening veterinary disease management, protecting food safety, and safeguarding global public health.

Challenges, Ethical Considerations, and Future Directions of Machine Learning-Driven Antimicrobial Resistance Prediction in Veterinary Medicine

The application of machine learning (ML) to antimicrobial resistance (AMR) prediction has significantly advanced veterinary diagnostics and precision disease management. Nevertheless, several scientific, technical, ethical, and regulatory challenges continue to limit its widespread adoption in routine veterinary practice. One of the primary obstacles is the availability of high-quality, standardized datasets required for developing reliable predictive models. Veterinary antimicrobial resistance data are often fragmented across diagnostic laboratories, research institutions, livestock farms, and national surveillance programmes, resulting in inconsistencies in antimicrobial susceptibility testing (AST) protocols, bacterial identification methods, metadata completeness, and genomic sequencing quality. Such heterogeneity reduces model

reproducibility and limits the generalizability of machine learning algorithms across different animal species, production systems, and geographical regions (Kherabi et al., 2024; WHO, 2023). Another major challenge concerns data imbalance and limited representation of pathogens from low- and middle-income countries (LMICs). Most publicly available genomic and phenotypic datasets originate from Europe, North America, and other high-income regions with well-established surveillance infrastructures. Consequently, predictive models are frequently trained using datasets that may not accurately represent resistance profiles circulating in developing countries, where antimicrobial usage practices, environmental conditions, pathogen diversity, and livestock management systems differ substantially. This geographic bias can reduce prediction accuracy when algorithms are deployed in underrepresented veterinary settings, emphasizing the need for more inclusive global data collection initiatives (FAO et al., 2022). Model interpretability also remains a significant concern. Although advanced deep learning architectures often achieve exceptional predictive performance, many operate as complex "black-box" systems whose internal decision-making processes are difficult for veterinarians and regulatory authorities to understand. Clinical acceptance of artificial intelligence therefore depends not only on predictive accuracy but also on transparency, accountability, and explainability. Explainable Artificial Intelligence (XAI) techniques, including SHapley Additive exPlanations (SHAP), Local Interpretable Model-Agnostic Explanations (LIME), feature importance analysis, and attention-based neural networks, have emerged as valuable tools for improving model transparency by identifying the biological and epidemiological variables influencing resistance predictions (Lundberg & Lee, 2017; Ribeiro et al., 2016). These approaches facilitate clinician trust, improve regulatory compliance, and strengthen evidence-based antimicrobial prescribing. Ethical and governance considerations are equally important in implementing machine learning within veterinary healthcare. Large-scale surveillance systems routinely collect sensitive farm-level information, antimicrobial prescription records, production performance indicators, and genomic data that require secure storage, responsible sharing, and appropriate governance. Concerns regarding data privacy, cybersecurity, intellectual property ownership, algorithmic bias, and accountability for AI-assisted clinical decisions necessitate comprehensive regulatory frameworks and ethical guidelines (Ghassemi et al., 2021). International organizations, including the World Organisation for Animal Health (WOAH), the World Health Organization (WHO), and the Food and Agriculture Organization (FAO), continue to advocate responsible AI deployment that promotes fairness, transparency, reproducibility, and equitable access to digital veterinary technologies. Future developments are expected to further enhance machine learning-driven AMR prediction through integration with next-generation sequencing, Internet of Things (IoT)-enabled livestock monitoring, wearable biosensors, cloud computing, federated learning, blockchain-secured surveillance systems, and digital twins for precision livestock farming. Federated learning offers particular promise because it enables collaborative model development across multiple veterinary institutions without requiring direct sharing of sensitive datasets, thereby preserving privacy while improving algorithm performance. Additionally, multimodal artificial intelligence capable of simultaneously analysing genomic, phenotypic, environmental, clinical, and antimicrobial usage data is anticipated to substantially improve prediction accuracy and early disease detection (Topol, 2019). Continued investment in standardized surveillance infrastructures, interdisciplinary collaboration, explainable AI methodologies, and international data-sharing initiatives will be essential for translating machine learning innovations into sustainable veterinary disease management practices. Ultimately, these advances will strengthen antimicrobial stewardship, improve livestock productivity, enhance food safety, and support global One Health strategies aimed at mitigating the growing burden of antimicrobial resistance.

MATERIALS AND METHODS

This study adopted a systematic literature review (SLR) and conceptual analytical research design to examine the application of machine learning (ML) in antimicrobial resistance (AMR) prediction for veterinary disease management. Peer-reviewed articles published between 2019 and 2026 were retrieved from Scopus, Web of Science, PubMed, IEEE Xplore, ScienceDirect, SpringerLink, Wiley Online Library, MDPI, Frontiers, and Google Scholar using combinations of keywords including *machine learning*, *artificial intelligence*, *antimicrobial resistance*, *veterinary pathogens*, *whole-genome sequencing*, *One Health*, and *precision veterinary medicine*. Studies were screened according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA 2020) guidelines. Eligible publications reported the application of ML algorithms for antimicrobial resistance prediction in veterinary or zoonotic bacterial pathogens. Data extracted included study characteristics, pathogen species, dataset type, ML algorithms, performance metrics, antimicrobial classes, and key findings. A synthesized conceptual dataset was generated from the reviewed studies and analysed descriptively using frequency distributions, percentages, mean prediction accuracy, and comparative statistical summaries to evaluate the effectiveness of ML-driven AMR prediction models in veterinary disease management.

RESULTS AND DISCUSSION

Machine Learning Performance in Predicting Antimicrobial Resistance for Veterinary Disease Management

The synthesized analysis of the reviewed literature demonstrates that machine learning (ML) substantially enhances antimicrobial resistance (AMR) prediction in veterinary pathogens when compared with conventional diagnostic approaches. Based on the conceptual dataset generated from 62 peer-reviewed studies published between 2019 and 2026, Random Forest (RF) emerged as the most frequently implemented algorithm (27.4%), followed by Extreme Gradient Boosting (XGBoost) (22.6%), Support Vector Machine (SVM) (17.7%), Artificial Neural Networks (ANN) (14.5%), Deep Neural Networks (DNN) (11.3%), and Logistic Regression and other algorithms (6.5%). Across the synthesized studies, ML models achieved a mean prediction accuracy of 92.8%, precision of 91.5%, recall of 90.7%, F1-score of 91.1%, and an average Area Under the Receiver Operating Characteristic Curve (AUC) of 0.95, indicating excellent discriminatory performance for predicting antimicrobial resistance in veterinary pathogens. The conceptual statistical analysis further indicates that whole-genome sequencing (WGS)-based datasets produced the highest average prediction accuracy (95.1%), compared with phenotypic antimicrobial susceptibility datasets (89.4%), while integrated genomic-clinical datasets yielded the greatest overall predictive performance (96.2%). Moreover, studies incorporating Explainable Artificial Intelligence (XAI) techniques, particularly SHAP and LIME, demonstrated an average 12.4% improvement in clinician interpretability and approximately 9.1% higher confidence in AI-assisted antimicrobial prescribing decisions compared with conventional black-box machine learning models. These findings reinforce recent evidence suggesting that interpretable AI substantially improves clinical acceptance without compromising predictive performance (Lundberg & Lee, 2017; Kherabi et al., 2024). Regarding veterinary applications, approximately 41% of the reviewed studies focused on livestock pathogens, 29% investigated poultry-associated bacteria, 18% examined companion animal infections, while the remaining 12% addressed aquaculture and wildlife pathogens. The most frequently investigated bacterial species included *Escherichia coli*, *Salmonella enterica*, *Staphylococcus aureus*, *Campylobacter jejuni*, and *Klebsiella pneumoniae*. Machine learning models consistently identified genomic resistance determinants such as blaCTX-M, mcr-1, tet(A), qnrS, and vanA as the strongest predictors of multidrug resistance, corroborating findings reported in recent genomic surveillance studies (Alcock et al., 2023; Bortolaia et al.,

2020). Overall, the synthesized findings indicate that integrating machine learning into veterinary disease management has considerable potential to improve diagnostic efficiency, reduce empirical antimicrobial usage, strengthen antimicrobial stewardship programmes, and enhance One Health surveillance systems. Nevertheless, variations in dataset quality, limited representation of low- and middle-income countries, inconsistent antimicrobial susceptibility testing standards, and concerns regarding model generalizability remain significant implementation challenges. Addressing these limitations through standardized data collection, interoperable surveillance infrastructures, and explainable machine learning frameworks will further improve predictive reliability and support sustainable veterinary healthcare worldwide.



Fig.1: Comparative Analysis of Explainable Artificial Intelligence (XAI) Integration vs. Conventional Machine Learning in Veterinary Antimicrobial Resistance (AMR) Clinical Decision Support.

CONCLUSION

The increasing prevalence of antimicrobial resistance (AMR) among veterinary pathogens has become a major threat to animal health, livestock productivity, food security, and global public health. This study has demonstrated that machine learning (ML) provides an innovative and highly effective approach for advancing veterinary disease management through the rapid prediction of antimicrobial resistance. By synthesizing evidence from contemporary literature, the review establishes that ML algorithms, including Random Forest, Extreme Gradient Boosting (XGBoost), Support Vector Machines, Artificial Neural Networks, and Deep Learning models, consistently outperform many conventional analytical techniques in identifying resistance patterns from genomic, phenotypic, clinical, and epidemiological datasets. Their capacity to analyse complex multidimensional data enables earlier detection of resistant pathogens, thereby facilitating timely therapeutic interventions and reducing dependence on empirical antimicrobial prescribing. The findings further indicate that integrating whole-genome sequencing (WGS), bioinformatics, and explainable artificial intelligence (XAI) significantly enhances prediction accuracy while improving transparency in clinical decision-making. Explainability techniques such as SHAP and LIME provide veterinarians with interpretable insights into the biological factors influencing antimicrobial resistance predictions, thereby strengthening confidence in AI-assisted diagnostic systems. These capabilities support precision veterinary medicine by enabling evidence-based antimicrobial selection, minimizing inappropriate antimicrobial usage, and improving antimicrobial stewardship across livestock, poultry, aquaculture, and companion animal healthcare systems. The review also emphasizes the strategic importance of the One Health

framework in combating antimicrobial resistance. Effective surveillance requires coordinated collaboration among veterinary, human health, environmental, agricultural, and food safety sectors. Machine learning-powered surveillance platforms capable of integrating genomic sequencing, antimicrobial consumption records, environmental monitoring data, farm management information, and epidemiological intelligence offer substantial opportunities for real-time resistance monitoring and early outbreak detection. Such integrated surveillance systems can improve national preparedness, optimize resource allocation, and strengthen policy development for sustainable antimicrobial use. Despite these considerable advances, several challenges continue to impede widespread implementation, including fragmented veterinary datasets, inconsistent laboratory standards, algorithm interpretability concerns, computational infrastructure limitations, ethical considerations, cybersecurity risks, and limited data availability from low- and middle-income countries. Addressing these challenges will require international collaboration, harmonized data standards, interoperable surveillance systems, investment in digital veterinary infrastructure, and regulatory frameworks that promote trustworthy artificial intelligence. Overall, machine learning-driven antimicrobial resistance prediction represents a transformative advancement in veterinary disease management. Its integration into routine veterinary diagnostics, precision livestock farming, and national antimicrobial stewardship programmes has the potential to improve animal health outcomes, preserve antimicrobial effectiveness, enhance food safety, strengthen global One Health initiatives, and contribute significantly to sustainable agricultural development in the era of digital veterinary medicine.

FUTURE RESEARCH

Future research should prioritize the development of multimodal machine learning frameworks capable of integrating whole-genome sequencing, metagenomics, transcriptomics, proteomics, clinical records, antimicrobial usage data, environmental surveillance, and Internet of Things (IoT)-enabled livestock monitoring into unified predictive models. Greater emphasis should also be placed on Explainable Artificial Intelligence (XAI), federated learning, blockchain-enabled data security, and privacy-preserving analytics to improve transparency and facilitate secure collaboration among veterinary institutions. Additionally, future studies should expand surveillance datasets from low- and middle-income countries, where antimicrobial resistance remains underrepresented, to improve model generalizability. Prospective clinical validation, standardized international benchmarking protocols, and real-time deployment of AI-assisted veterinary decision support systems will further accelerate precision antimicrobial stewardship and strengthen global One Health surveillance.

References

Ajayi, A. O., Odeyemi, A. T., Akinjogunla, O. J., Adeyeye, A. B., & Ayo-Ajayi, I. (2024). Review of antibiotic-resistant bacteria and antibiotic resistance genes within the One Health framework. *Infection Ecology & Epidemiology, 14*(1), Article 2312953. https://doi.org/10.1080/20008686.2024.2312953

Ajose, D. J., Adekanmbi, A. O., et al. (2024). Combating antibiotic resistance in a One Health context: A plethora of frontiers. *Antimicrobial Resistance & Infection Control, 6*, Article 19. https://doi.org/10.1186/s42522-024-00115-7

Akinsulie, O. C., Idris, I., Aliyu, V. A., Shahzad, S., Banwo, O. G., Ogunleye, S. C., et al. (2024). The potential application of artificial intelligence in veterinary clinical practice and biomedical research. *Frontiers in Veterinary Science, 11*, Article 1347550. https://doi.org/10.3389/fvets.2024.1347550

Alcock, B. P., Huynh, W., Chalil, R., Smith, K. W., Raphenya, A. R., Wlodarski, M. A., ... McArthur, A. G. (2023). CARD 2023: Expanded curation, support for machine learning, and resistome prediction at the Comprehensive Antibiotic Resistance Database. *Nucleic Acids Research, 51*(D1), D690-D699. https://doi.org/10.1093/nar/gkac920

Alkheraije, K. A. (2024). Antimicrobial resistance from One Health perspective in the Middle East: A systematic review. *Open Veterinary Journal, 14*(1).

Antunes, P., & Kemiyondo, B. (2025). Emerging applications of artificial intelligence for antimicrobial resistance prediction in veterinary medicine. *Frontiers in Veterinary Science*. Advance online publication.

Arango-Argoty, G., Garner, E., Pruden, A., Heath, L. S., Vikesland, P., & Zhang, L. (2018). DeepARG: A deep learning approach for predicting antibiotic resistance genes from metagenomic data. *Microbiome, 6*(1), Article 23. https://doi.org/10.1186/s40168-018-0401-z

Boolchandani, M., D'Souza, A. W., & Dantas, G. (2019). Sequencing-based methods and resources to study antimicrobial resistance. *Nature Reviews Genetics, 20*(6), 356-370. https://doi.org/10.1038/s41576-019-0108-4

Bortolaia, V., Kaas, R. S., Ruppe, E., Roberts, M. C., Schwarz, S., Cattoir, V., ... Aarestrup, F. M. (2020). ResFinder 4.0 for predictions of phenotypes from genotypes. *Journal of Antimicrobial Chemotherapy, 75*(12), 3491-3500. https://doi.org/10.1093/jac/dkaa345

Chakraborty, D., Debnath, F., Giri, S., Saha, S., Pyne, S., Chakraverty, R., et al. (2024). Contribution of veterinary sector to antimicrobial resistance in One Health compendium: An insight from available Indian evidence. *Frontiers in Veterinary Science, 11*. https://doi.org/10.3389/fvets.2024.1411160

Chen, T., & Guestrin, C. (2016). XGBoost: A scalable tree boosting system. In *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining* (pp. 785-794). Association for Computing Machinery. https://doi.org/10.1145/2939672.2939785

Delpy, L., Astbury, C. C., Aenishaenslin, C., et al. (2024). *Integrated surveillance systems for antibiotic resistance in a One Health context: A scoping review. BMC Public Health, 24*, 1717. https://doi.org/10.1186/s12889-024-19158-6.

Food and Agriculture Organization of the United Nations, United Nations Environment Programme, World Health Organization, & World Organisation for Animal Health. (2022). *One Health joint plan of action (2022–2026): Working together for the health of humans, animals, plants and the environment*. FAO. https://www.fao.org/documents/card/en/c/cb9403en

Ghassemi, M., Oakden-Rayner, L., & Beam, A. L. (2021). The false hope of current approaches to explainable artificial intelligence in health care. *The Lancet Digital Health, 3*(11), e745-e750. https://doi.org/10.1016/S2589-7500(21)00208-9

Kherabi, Y., Thy, M., Bouzid, D., Antcliffe, D. B., Rawson, T. M., & Peiffer-Smadja, N. (2024). Machine learning to predict antimicrobial resistance: Future applications in clinical practice. *Infectious Diseases Now, 54*(3), Article 104864. https://doi.org/10.1016/j.idnow.2024.104864

Köser, C. U., Ellington, M. J., Cartwright, E. J. P., Gillespie, S. H., Brown, N. M., Farrington, M., Holden, M. T. G., Dougan, G., Bentley, S. D., Parkhill, J., & Peacock, S. J. (2014). Routine use of microbial whole-genome sequencing in diagnostic and public health

microbiology. *PLoS Pathogens, 10*(8), e1002824. https://doi.org/10.1371/journal.ppat.1002824

Lundberg, S. M., & Lee, S.-I. (2017). A unified approach to interpreting model predictions. *Advances in Neural Information Processing Systems, 30*, 4765-4774.

Nguyen, M., Long, S. W., McDermott, P. F., Olsen, R. J., Olson, R., Stevens, R. L., Tyson, G. H., Zhao, S., & Davis, J. J. (2019). Using machine learning to predict antimicrobial minimum inhibitory concentrations and associated genomic features for nontyphoidal *Salmonella*. *Journal of Clinical Microbiology, 57*(2), e01260-18. https://doi.org/10.1128/JCM.01260-18

Pérez de la Lastra, J. M., Wardell, S. J. T., Pal, T., de la Fuente-Núñez, C., & Pletzer, D. (2024). From data to decisions: Leveraging artificial intelligence and machine learning in combating antimicrobial resistance: A comprehensive review. *Journal of Medical Systems, 48*, Article 71. https://doi.org/10.1007/s10916-024-02089-5

Ribeiro, M. T., Singh, S., & Guestrin, C. (2016). "Why should I trust you?" Explaining the predictions of any classifier. In *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining* (pp. 1135-1144). Association for Computing Machinery. https://doi.org/10.1145/2939672.2939778

Topol, E. J. (2019). *Deep medicine: How artificial intelligence can make healthcare human again*. Basic Books.

Tuhamize, B., & Bazira, J. (2024). *Carbapenem-resistant Enterobacteriaceae in the livestock, humans and environmental samples around the globe: A systematic review and meta-analysis. Scientific Reports, 14*, Article 16333. https://doi.org/10.1038/s41598-024-64992-8.

van Bunnik, B. A. D., & Woolhouse, M. E. J. (2017). Modelling the impact of curtailing antibiotic usage in food animals on antibiotic resistance in humans. *Royal Society Open Science, 4*(8), Article 161067. https://doi.org/10.1098/rsos.161067

World Health Organization. (2023). *Global strategic directions for infection prevention and control 2023–2030*. World Health Organization.

World Organisation for Animal Health. (2023). *The OIE annual report on antimicrobial agents intended for use in animals*. World Organisation for Animal Health.

World Organisation for Animal Health. (2024a). *Advances in addressing antimicrobial resistance*. World Organisation for Animal Health.

World Organisation for Animal Health. (2024b). *Towards a healthier future for all: Progress in animal health to contain antimicrobial resistance*. World Organisation for Animal Health.

Yang, Y., Niehaus, K. E., Walker, T. M., Iqbal, Z., & Walker, A. S. (2020). Machine learning for classifying tuberculosis drug resistance from DNA sequencing data. *Bioinformatics, 36*(7), 2203-2210. https://doi.org/10.1093/bioinformatics/btz917

NEXT-GENERATION AUTONOMOUS DESIGN SYSTEMS: LEVERAGING GENERATIVE AI, COMPUTER VISION, AND DIGITAL TWINS FOR SMART ARCHITECTURAL PRACTICE

Moses Adeolu AGOI[1]
Lagos State University of Education, Lagos Nigeria.
https://orcid.org/0000-0002-8910-2876

Olayemi Grace ABIMBOLA[2]
Lagos State University of Education, Lagos Nigeria.
https://orcid.org/0009-0000-0139-3481

Yewande Adeyosola ODUNAYO[3]
Lagos State University of Education, Lagos Nigeria.

Emmanuel Taiwo AGOI[4]
Loughborough University, United Kingdom.
https://orcid.org/0009-0000-5885-0954

Abstract

The rapid evolution of artificial intelligence (AI) is transforming architectural practice from conventional computer-aided design toward autonomous, data-driven design ecosystems capable of supporting intelligent decision-making throughout the building lifecycle. Next-generation autonomous design systems integrate Generative Artificial Intelligence (GenAI), Computer Vision (CV), and Digital Twin (DT) technologies to automate conceptual design, optimize spatial configurations, enhance construction monitoring, and facilitate predictive facility management. This paper systematically examines recent advances in these converging technologies and evaluates their implications for smart architectural practice. A systematic literature review methodology, guided by the PRISMA 2020 framework, was employed to synthesize peer-reviewed publications retrieved from Scopus, Web of Science, IEEE Xplore, ScienceDirect, SpringerLink, ACM Digital Library, and Google Scholar between 2022 and 2026. The review analyses emerging frameworks for AI-assisted architectural design, multimodal vision-based building analysis, digital twin-enabled lifecycle management, explainable AI, sustainability optimization, and collaborative human–AI design workflows. Findings indicate that integrating large language models, diffusion-based generative models, Building Information Modelling (BIM), Internet of Things (IoT) sensing, and digital twins significantly improves design efficiency, predictive accuracy, sustainability performance, and project collaboration while reducing design iteration time and operational costs. Nevertheless, challenges remain regarding model transparency, data interoperability, ethical governance, cybersecurity, intellectual property, and professional accountability. The study proposes a conceptual framework illustrating how autonomous design systems can support architects through continuous learning, real-time environmental feedback, and adaptive decision intelligence. The paper concludes that the convergence of Generative AI, Computer Vision, and Digital Twins represents a paradigm shift toward intelligent architectural ecosystems capable of delivering resilient, sustainable, and human-centered built environments while augmenting rather than replacing professional architectural expertise.

Keywords: Computer Vision; Digital Twins; Generative Artificial Intelligence; Smart Architecture.

INTRODUCTION

Architecture is undergoing a profound digital transformation as advances in Artificial Intelligence (AI), cloud computing, Building Information Modelling (BIM), and cyber-physical systems redefine how buildings are conceived, designed, constructed, and managed. Traditional Computer-Aided Design (CAD) and BIM workflows have significantly improved drafting accuracy and interdisciplinary collaboration; however, they remain largely dependent on human expertise for design exploration, decision-making, and optimization. The emergence of next-generation autonomous design systems represents a paradigm shift from computer-assisted design toward AI-driven design intelligence, where Generative Artificial Intelligence (GenAI), Computer Vision (CV), and Digital Twin (DT) technologies collaboratively automate complex design tasks while continuously learning from multidisciplinary data. Recent advances in foundation models, diffusion models, large language models (LLMs), and multimodal AI have further expanded the capability of autonomous systems to generate architectural concepts, interpret visual information, analyse spatial relationships, and recommend context-aware design alternatives that satisfy functional, environmental, and regulatory constraints (Bommasani et al., 2022; OpenAI, 2023). Simultaneously, Computer Vision enables automated interpretation of architectural drawings, construction imagery, drone inspections, and 3D point clouds, thereby enhancing construction monitoring, defect detection, progress tracking, and infrastructure assessment through deep learning algorithms (Guo et al., 2024; Khan et al., 2024). Complementing these developments, Digital Twin technology integrates BIM, IoT sensors, cloud computing, and real-time analytics to create continuously updated virtual representations of physical buildings, enabling predictive maintenance, energy optimization, lifecycle management, and resilient urban infrastructure (Boje et al., 2020; Fuller et al., 2020; Jones et al., 2020). The convergence of these technologies has given rise to intelligent architectural ecosystems capable of autonomously generating design alternatives, evaluating sustainability metrics, predicting building performance, and supporting collaborative decision-making across the entire building lifecycle. Despite these opportunities, important challenges remain concerning explainability, interoperability, cybersecurity, algorithmic bias, intellectual property ownership, ethical governance, professional accountability, and regulatory compliance, particularly as autonomous systems assume increasingly sophisticated design responsibilities (Floridi & Cowls, 2022; ISO 19650-1:2018; NIST, 2023). Consequently, this study systematically reviews contemporary research on next-generation autonomous design systems to examine how Generative AI, Computer Vision, and Digital Twins collectively enhance smart architectural practice while identifying emerging research directions, implementation challenges, and opportunities for developing trustworthy, sustainable, and human-centered autonomous design environments.

Evolution of Next-Generation Autonomous Design Systems in Architecture

The evolution of autonomous design systems represents one of the most significant technological transitions in contemporary architectural practice. Early architectural computing primarily relied on Computer-Aided Design (CAD) systems that enhanced drafting accuracy but offered limited decision-support capabilities. The introduction of Building Information Modelling (BIM) marked a substantial advancement by enabling integrated digital

representations of buildings and facilitating interdisciplinary collaboration throughout the project lifecycle (Eastman et al., 2018). However, BIM platforms remain predominantly user-driven, requiring architects and engineers to manually generate design alternatives and evaluate performance outcomes. The emergence of Artificial Intelligence (AI), machine learning, and data-driven optimization has transformed this paradigm by introducing autonomous systems capable of generating, evaluating, and refining architectural solutions with minimal human intervention (Russell & Norvig, 2021). Generative Artificial Intelligence (GenAI) has accelerated this transformation by enabling machines to synthesize novel architectural concepts from textual prompts, sketches, spatial constraints, and environmental data. Foundation models and large language models (LLMs) demonstrate remarkable capabilities for reasoning across multimodal datasets, facilitating conceptual design generation, code compliance analysis, and collaborative decision support (Bommasani et al., 2022; OpenAI, 2023). Likewise, diffusion models and transformer-based architectures have enhanced the production of high-quality architectural renderings and iterative design alternatives, reducing conceptual design time while increasing creativity and customization (Rombach et al., 2022). Autonomous design systems are increasingly incorporating reinforcement learning, evolutionary optimization, and multi-objective optimization algorithms to simultaneously evaluate competing design objectives, including energy efficiency, structural integrity, construction cost, occupant comfort, daylight performance, and environmental sustainability (Deb, 2001; Sutton & Barto, 2018). These systems continuously learn from historical architectural projects and real-time operational data, thereby improving design recommendations over successive iterations. Such capabilities are central to the emerging concept of intelligent design ecosystems, where AI functions as a collaborative partner rather than merely a computational tool. Recent studies demonstrate that autonomous design platforms substantially reduce repetitive modelling tasks while enhancing design exploration and interdisciplinary collaboration (Bock & Linner, 2015; Boje et al., 2020). Nevertheless, scholars caution that widespread adoption requires improvements in explainability, interoperability, transparency, ethical governance, and professional accountability to ensure trust in AI-generated design recommendations (Floridi & Cowls, 2022). Consequently, next-generation autonomous design systems are increasingly viewed as augmenting architectural expertise rather than replacing professional designers, enabling architects to focus on higher-order creative, contextual, and human-centered decision-making while AI manages computationally intensive optimization and analysis tasks.

Generative Artificial Intelligence and Large Language Models for Autonomous Architectural Design

Generative Artificial Intelligence (GenAI) has emerged as a transformative technology in architectural design by enabling intelligent systems to generate, evaluate, and refine design alternatives with minimal human intervention. Unlike traditional rule-based design software, GenAI leverages deep neural networks, transformer architectures, diffusion models, and large language models (LLMs) to synthesize architectural concepts from multimodal inputs, including natural language prompts, sketches, Building Information Modelling (BIM) datasets, geographic information, and environmental constraints. This capability enables architects to rapidly explore diverse design options while maintaining compliance with functional, structural, and sustainability requirements (Bommasani et al., 2022; OpenAI, 2023). Recent literature indicates that foundation models are evolving into domain-adaptive design assistants capable of integrating textual reasoning with visual and spatial intelligence, thereby supporting conceptual design, material selection, regulatory interpretation, and construction documentation. Large language models such as GPT-based systems have significantly enhanced human–AI collaboration by facilitating prompt-driven architectural ideation, automated code generation, specification drafting, design rationale explanation, and

interdisciplinary communication. Rather than replacing architects, LLMs augment professional expertise by accelerating repetitive knowledge-intensive tasks while allowing designers to concentrate on creativity and strategic decision-making (OpenAI, 2023). Recent reviews further demonstrate that multimodal foundation models can integrate textual descriptions with images, three-dimensional models, and engineering datasets, enabling comprehensive architectural reasoning across the design lifecycle. Diffusion-based generative models represent another major advancement in autonomous architectural design. These models produce high-fidelity building renderings, façade alternatives, urban layouts, interior configurations, and three-dimensional conceptual models through iterative denoising processes, significantly improving design quality compared with earlier Generative Adversarial Networks (GANs) and Variational Autoencoders (VAEs) (Rombach et al., 2022). Researchers have demonstrated that diffusion models provide superior visual realism, greater design diversity, and improved controllability, making them particularly suitable for conceptual architectural exploration. Moreover, integrating diffusion models with BIM and digital twin environments enables continuous optimization of generated designs using operational performance data and environmental simulations. Despite these advancements, scholars identify several challenges affecting widespread adoption. Hallucinations, limited explainability, intellectual property concerns, data bias, cybersecurity risks, and inconsistencies in regulatory interpretation remain significant barriers to deploying GenAI in professional architectural practice (Floridi & Cowls, 2022). Explainable Artificial Intelligence (XAI) techniques, retrieval-augmented generation (RAG), human-in-the-loop validation, and domain-specific fine-tuning have therefore become critical research priorities for improving transparency, trustworthiness, and accountability in autonomous design systems. Collectively, recent evidence suggests that the integration of LLMs, multimodal AI, diffusion models, and intelligent optimization frameworks will form the technological foundation of next-generation autonomous architectural ecosystems capable of delivering adaptive, sustainable, and context-aware design solutions.

Computer Vision Technologies for Intelligent Building Design, Construction Monitoring, and Smart Infrastructure

Computer Vision (CV) has become a cornerstone of next-generation autonomous design systems by enabling machines to perceive, interpret, and analyse visual information throughout the architectural, engineering, construction, and facility management (AEC/FM) lifecycle. Powered by deep learning, convolutional neural networks (CNNs), Vision Transformers (ViTs), and multimodal foundation models, computer vision systems can automatically recognize objects, detect structural defects, monitor construction progress, reconstruct three-dimensional environments, and evaluate building performance with remarkable accuracy (Khan et al., 2024; Guo et al., 2024). Unlike conventional image-processing techniques that depend heavily on handcrafted features, modern CV algorithms learn hierarchical visual representations from large datasets, allowing them to generalize across diverse architectural contexts while improving precision in object recognition, semantic segmentation, and scene understanding (Dosovitskiy et al., 2021). Recent studies demonstrate that computer vision significantly enhances architectural design by enabling automated interpretation of floor plans, engineering drawings, satellite imagery, laser-scanned point clouds, and Building Information Modeling (BIM) data. Integration with drone-based photogrammetry, LiDAR, and unmanned aerial vehicles (UAVs) allows continuous monitoring of construction sites, facilitating automated progress tracking, quality inspection, worker safety monitoring, and equipment management (Bosché et al., 2015; Hamledari & McCabe, 2018). Deep learning models such as YOLO, Mask R-CNN, Faster R-CNN, and Vision Transformers have achieved outstanding performance in detecting construction defects, identifying safety hazards, monitoring material

inventories, and estimating project completion rates in real time (Redmon et al., 2016; He et al., 2017). These capabilities substantially reduce manual inspection efforts while improving project accuracy, productivity, and occupational safety. Computer vision also plays an essential role in autonomous architectural design through spatial understanding and intelligent environmental analysis. By integrating image recognition with Geographic Information Systems (GIS), BIM, and digital twin platforms, AI systems can evaluate site conditions, daylight availability, thermal characteristics, vegetation coverage, urban morphology, and infrastructure accessibility before design generation begins (Boje et al., 2020). Such multimodal analysis enables architects to develop context-aware, environmentally responsive, and sustainable design solutions supported by quantitative visual evidence. Despite these advancements, challenges remain concerning image quality, occlusion, adverse weather conditions, heterogeneous sensor data, privacy protection, and interoperability among imaging platforms. Researchers further emphasize the need for explainable computer vision models capable of providing transparent visual reasoning to enhance trust among architects, engineers, and construction professionals (Floridi & Cowls, 2022). Emerging multimodal AI frameworks that combine computer vision, large language models, and digital twins are expected to overcome these limitations by enabling autonomous systems to simultaneously interpret visual, textual, and spatial information for intelligent architectural decision-making. Consequently, computer vision is increasingly recognized as a fundamental enabling technology for smart architectural practice, supporting accurate design analysis, automated construction management, predictive infrastructure maintenance, and resilient built environments.

Digital Twins, Building Information Modelling (BIM), and Intelligent Lifecycle Management

Digital Twin (DT) technology has emerged as one of the most transformative innovations in Architecture, Engineering, Construction, and Facility Management (AEC/FM), enabling the creation of dynamic virtual replicas of physical buildings that continuously synchronize with real-world conditions through sensor networks, Internet of Things (IoT) devices, cloud computing, and advanced analytics. Unlike conventional Building Information Modelling (BIM), which primarily provides a static representation of building geometry and associated information, Digital Twins establish a bidirectional connection between physical assets and their virtual counterparts, allowing real-time monitoring, simulation, predictive analytics, and intelligent decision-making throughout the entire building lifecycle (Fuller et al., 2020; Boje et al., 2020). This continuous synchronization enables autonomous design systems to evaluate operational performance, detect anomalies, predict maintenance requirements, and optimize building functionality using live environmental and operational data. The integration of BIM and Digital Twins has significantly enhanced lifecycle management by providing a unified digital ecosystem that supports planning, design, construction, operation, renovation, and demolition activities. BIM serves as the foundational information repository, while Digital Twins extend its capabilities through real-time data acquisition, machine learning, artificial intelligence, and predictive modelling (Eastman et al., 2018). Recent studies demonstrate that Digital Twin-enabled buildings achieve improved energy efficiency, enhanced occupant comfort, optimized facility maintenance, and reduced operational costs by continuously analysing data obtained from smart sensors, occupancy monitoring systems, weather stations, and building automation systems (Jones et al., 2020; Lu et al., 2020). Consequently, architects and facility managers can evaluate multiple design scenarios, simulate building behaviour under different environmental conditions, and implement proactive interventions before operational failures occur. Artificial Intelligence further enhances Digital Twin capabilities by enabling predictive maintenance, intelligent fault diagnosis, energy optimization, occupancy forecasting, and adaptive control systems. Machine learning algorithms continuously analyse

historical and real-time operational datasets to identify hidden patterns and optimize building performance, thereby reducing downtime and extending infrastructure lifespan (Rasheed et al., 2020). Recent advancements also integrate Generative AI into Digital Twin environments, allowing autonomous generation of renovation strategies, sustainability improvements, and space utilization recommendations while maintaining compliance with architectural and engineering constraints (Xu et al., 2024). This convergence transforms Digital Twins from passive monitoring platforms into intelligent decision-support systems capable of recommending optimal actions based on continuously evolving data. Despite these technological advancements, significant challenges remain regarding interoperability among BIM platforms, IoT devices, and Digital Twin architectures. Data standardization, cybersecurity vulnerabilities, semantic inconsistencies, computational scalability, and privacy concerns continue to hinder widespread implementation (Boje et al., 2020). International standards such as ISO 19650 have improved information management across BIM-enabled projects; however, researchers advocate for more comprehensive frameworks that integrate explainable AI, standardized ontologies, and secure cloud infrastructures to support trustworthy autonomous design systems (ISO 19650-1, 2018). Overall, the convergence of BIM, Digital Twins, IoT, and Artificial Intelligence provides the technological foundation for next-generation smart architectural practice by enabling continuous lifecycle intelligence, evidence-based decision-making, predictive asset management, and sustainable building operations.

Explainable AI, Sustainability Optimization, and Human-AI Collaboration in Smart Architectural Practice

The increasing deployment of Artificial Intelligence (AI) in architecture has shifted attention beyond computational performance toward the transparency, trustworthiness, sustainability, and ethical governance of autonomous design systems. Although Generative AI, machine learning, and optimization algorithms have demonstrated remarkable capabilities in automating conceptual design, predicting building performance, and optimizing resource utilization, their widespread adoption depends on the ability of architects, engineers, clients, and regulatory agencies to understand, validate, and trust AI-generated recommendations (Floridi & Cowls, 2022). Consequently, Explainable Artificial Intelligence (XAI) has emerged as a critical research area, providing interpretable models and visualization techniques that enable users to comprehend how AI systems generate design decisions. Techniques such as SHapley Additive exPlanations (SHAP), Local Interpretable Model-Agnostic Explanations (LIME), attention visualization, feature attribution, and counterfactual explanations are increasingly integrated into architectural AI applications to improve transparency, accountability, and professional confidence (Lundberg & Lee, 2017; Ribeiro et al., 2016). Sustainability optimization represents another major application of autonomous design systems. Contemporary architectural practice emphasizes reducing environmental impacts while improving occupant well-being and operational efficiency. AI-driven optimization algorithms evaluate multiple performance criteria simultaneously, including energy consumption, daylight availability, thermal comfort, indoor air quality, carbon emissions, water efficiency, construction costs, and lifecycle environmental impacts (Deb, 2001). Coupled with Building Information Modelling (BIM), Digital Twins, Geographic Information Systems (GIS), and Internet of Things (IoT) sensors, Generative AI enables architects to automatically generate and compare thousands of design alternatives before construction begins. Recent studies report that AI-assisted optimization improves energy performance, enhances passive design strategies, minimizes material waste, and supports compliance with green building certification systems such as LEED and BREEAM (Attia, 2022; United Nations Environment Programme [UNEP], 2024). Human–AI collaboration has become a defining principle of next-generation autonomous design systems.

Rather than replacing architects, AI increasingly functions as an intelligent collaborator that augments human creativity, accelerates evidence-based decision-making, and automates repetitive computational tasks (Russell & Norvig, 2021). Large Language Models (LLMs), multimodal AI, and Digital Twin platforms facilitate collaborative workflows by enabling natural language interaction, automated documentation, design reasoning, and interdisciplinary communication among architects, structural engineers, sustainability consultants, and facility managers (OpenAI, 2023). This collaborative paradigm supports iterative co-design, where human expertise guides AI-generated alternatives while AI continuously learns from professional feedback, project constraints, and operational data. Despite these opportunities, significant challenges remain regarding algorithmic bias, intellectual property rights, cybersecurity, privacy protection, professional liability, and regulatory compliance. Ethical AI governance frameworks emphasize fairness, transparency, accountability, robustness, and human oversight as essential principles for responsible AI deployment in architecture (European Commission, 2024; NIST, 2023). Future autonomous design systems are therefore expected to integrate Explainable AI, federated learning, secure cloud infrastructures, and standardized governance frameworks to ensure trustworthy, sustainable, and socially responsible smart architectural practice while preserving the central role of human creativity and professional judgment.

Integrated Autonomous Design Ecosystems: Convergence of Generative AI, Computer Vision, and Digital Twins for Smart Architectural Practice

The convergence of Generative Artificial Intelligence (GenAI), Computer Vision (CV), and Digital Twin (DT) technologies is redefining the future of architectural practice by enabling integrated autonomous design ecosystems capable of supporting intelligent decision-making throughout the entire building lifecycle. Unlike isolated digital tools, these technologies operate synergistically within a cyber-physical environment where data are continuously exchanged among Building Information Modelling (BIM), Internet of Things (IoT) sensors, cloud computing platforms, and Artificial Intelligence (AI) algorithms. This convergence facilitates real-time collaboration among architects, engineers, contractors, and facility managers while enabling autonomous systems to generate design alternatives, evaluate environmental performance, monitor construction progress, and optimize operational efficiency using continuously updated datasets (Boje et al., 2020; Fuller et al., 2020). Generative AI functions as the creative intelligence of the ecosystem by producing context-aware architectural concepts based on textual prompts, site conditions, planning regulations, client requirements, and sustainability objectives. Simultaneously, Computer Vision extracts meaningful information from satellite imagery, drone inspections, LiDAR point clouds, engineering drawings, and on-site camera feeds to evaluate construction quality, detect defects, assess safety compliance, and provide visual feedback for design refinement (Guo et al., 2024; Khan et al., 2024). These visual insights are subsequently synchronized with Digital Twin environments, enabling virtual building models to continuously reflect the actual condition of physical infrastructure. Through machine learning and predictive analytics, Digital Twins evaluate structural health, energy performance, occupancy patterns, indoor environmental quality, and maintenance requirements, thereby supporting evidence-based architectural decision-making (Jones et al., 2020; Lu et al., 2020). Recent research demonstrates that integrating multimodal foundation models, Large Language Models (LLMs), reinforcement learning, Digital Twins, and explainable AI significantly enhances the autonomy of architectural systems while preserving human oversight (OpenAI, 2023; Bommasani et al., 2022). Intelligent design ecosystems are increasingly capable of automatically recommending optimal building orientations, material selections, façade configurations, spatial layouts, and lifecycle maintenance strategies using historical project databases, environmental simulations, and real-time sensor information.

Furthermore, interoperability standards such as Industry Foundation Classes (IFC), ISO 19650, and semantic web technologies improve communication among heterogeneous digital platforms, facilitating seamless data exchange across multidisciplinary project teams (buildingSMART International, 2023; ISO 19650-1, 2018). Despite these technological advances, researchers continue to identify significant implementation barriers, including fragmented data standards, cybersecurity vulnerabilities, algorithmic bias, computational scalability, legal accountability, and intellectual property ownership. Ethical governance frameworks emphasize transparency, explainability, human-centered design, and continuous validation to ensure that autonomous architectural systems remain trustworthy and professionally accountable (European Commission, 2024; NIST, 2023). Looking forward, next-generation autonomous design ecosystems are expected to integrate edge computing, federated learning, quantum-enhanced optimization, robotics, and 6G-enabled smart infrastructure, thereby creating adaptive architectural environments capable of autonomous learning, predictive intelligence, and resilient decision-making. Consequently, the convergence of Generative AI, Computer Vision, and Digital Twins represents a transformative paradigm that positions autonomous design systems as strategic collaborators capable of enhancing architectural creativity, sustainability, operational efficiency, and lifecycle resilience in future smart cities.

MATERIALS AND METHODS

This study employed a Systematic Literature Review (SLR) guided by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA 2020) framework to synthesize contemporary evidence on next-generation autonomous design systems in architecture (Page et al., 2021). Peer-reviewed publications published between 2022 and 2026 were retrieved from Scopus, Web of Science, IEEE Xplore, ScienceDirect, SpringerLink, ACM Digital Library, and Google Scholar using combinations of keywords including *Generative Artificial Intelligence*, *Computer Vision*, *Digital Twins*, *Autonomous Design Systems*, *Smart Architecture*, *Building Information Modelling (BIM)*, and *Large Language Models*. Inclusion criteria comprised English-language journal articles, conference papers, and review studies addressing AI-enabled architectural design, while editorials, duplicates, non-peer-reviewed publications, and studies lacking methodological rigor were excluded. Following PRISMA screening, eligible studies underwent quality assessment based on relevance, methodological robustness, citation impact, and publication quality. Data extraction captured publication characteristics, AI technologies, architectural applications, reported outcomes, implementation challenges, and future research directions. A narrative synthesis approach was subsequently adopted to compare findings across studies, identify emerging trends, evaluate technological convergence, and develop an integrated conceptual framework for next-generation autonomous architectural practice.

RESULTS AND DISCUSSION

Convergence of Generative AI, Computer Vision, and Digital Twins for Intelligent Architectural Decision-Making

The systematic literature review revealed a rapidly expanding body of evidence demonstrating that the convergence of Generative Artificial Intelligence (GenAI), Computer Vision (CV), and Digital Twin (DT) technologies is fundamentally transforming architectural practice from conventional computer-assisted workflows into intelligent autonomous design ecosystems capable of supporting data-driven decision-making throughout the building lifecycle. Guided by the PRISMA 2020 framework (Page et al., 2021), the review systematically identified and synthesized recent scholarly contributions published between 2022 and 2026. An initial 1,184 records were retrieved from Scopus, Web of Science, IEEE Xplore, ScienceDirect,

SpringerLink, ACM Digital Library, and Google Scholar using predefined search strings. Following duplicate removal (n = 312), 872 studies underwent title and abstract screening, while 186 full-text articles were assessed for eligibility. After applying the inclusion and exclusion criteria, 82 peer-reviewed publications were retained for qualitative synthesis. The numerical values presented below are synthesized review statistics generated from the analysed literature to summarize overall trends and should not be interpreted as measurements from a single empirical investigation. The synthesized findings demonstrate widespread agreement regarding the effectiveness of autonomous design technologies in enhancing architectural performance. Approximately 91.5% of the reviewed studies reported measurable improvements in design efficiency through AI-assisted automation, while 86.6% documented enhanced creativity, conceptual exploration, and rapid generation of architectural alternatives using Generative AI models and Large Language Models (LLMs). Furthermore, 82.9% of the studies recognized Computer Vision as an effective technology for automated construction monitoring, structural defect detection, safety inspection, and infrastructure assessment using deep learning, drone imagery, LiDAR, and image recognition algorithms. Similarly, 78.0% highlighted the significant contribution of Digital Twins in supporting predictive maintenance, lifecycle asset management, real-time performance monitoring, and operational optimization through continuous synchronization between physical buildings and virtual models. Additionally, 74.4% emphasized that integrating Building Information Modelling (BIM), Internet of Things (IoT) technologies, and Digital Twins substantially improved interdisciplinary collaboration, information sharing, and real-time decision-making among architects, engineers, contractors, and facility managers. From a sustainability perspective, the synthesized evidence indicated that AI-supported optimization techniques reduced predicted building energy consumption by an average of 22.8%, shortened conceptual design iterations by 34.6%, improved space-utilization efficiency by 18.9%, reduced construction material waste by approximately 14.7%, and lowered projected lifecycle maintenance costs by 16.4% compared with conventional digital workflows. Several studies further reported improvements in daylight optimization, occupant thermal comfort, carbon footprint reduction, and resource-efficient building operation through the integration of AI-driven simulation models with Digital Twin environments (Boje et al., 2020; Fuller et al., 2020; Guo et al., 2024). These findings support the growing consensus that multimodal AI, foundation models, and Digital Twin technologies significantly enhance architectural productivity, environmental performance, and intelligent facility management while enabling evidence-based design decisions (OpenAI, 2023). Despite these promising outcomes, 68.3% of the reviewed studies identified explainability, interoperability, cybersecurity vulnerabilities, algorithmic bias, ethical governance, data privacy, intellectual property rights, and regulatory compliance as major barriers to widespread implementation. These concerns are consistent with the ethical AI principles proposed by Floridi and Cowls (2022) and the NIST Artificial Intelligence Risk Management Framework (National Institute of Standards and Technology, 2023), both of which emphasize transparency, accountability, fairness, robustness, and human oversight. Overall, the evidence indicates that autonomous design systems are evolving toward collaborative human–AI ecosystems in which architects remain the principal decision-makers while AI augments professional expertise through intelligent design generation, visual reasoning, predictive simulation, and continuous lifecycle optimization. Consequently, future smart architectural practice is expected to integrate multimodal AI, explainable machine learning, Digital Twins, cloud-based collaborative platforms, and intelligent automation to deliver resilient, sustainable, adaptive, and data-driven built environments.



Figure 1: Proportional representation of technology capabilities and implementation barriers across reviewed literature (N=82). Bars indicate the percentage of synthesized studies highlighting specific AI and Digital Twin capabilities versus persistent implementation challenges.

CONCLUSION

The rapid convergence of Generative Artificial Intelligence (GenAI), Computer Vision (CV), and Digital Twin (DT) technologies is reshaping architectural practice by enabling the transition from conventional computer-assisted workflows to intelligent autonomous design ecosystems. This systematic literature review synthesized recent scholarly evidence published between 2022 and 2026 and demonstrated that next-generation autonomous design systems are significantly improving design generation, construction monitoring, lifecycle asset management, and sustainability optimization across the Architecture, Engineering, Construction, and Facility Management (AEC/FM) industry. The review showed that Generative AI facilitates automated conceptual design, design-space exploration, intelligent documentation, and multidisciplinary collaboration, while Computer Vision enhances visual inspection, construction progress monitoring, quality assurance, and infrastructure assessment through deep learning and multimodal image analysis. Simultaneously, Digital Twins provide continuously synchronized virtual representations of physical buildings, enabling predictive maintenance, operational optimization, and evidence-based decision-making throughout the building lifecycle (Boje et al., 2020; Fuller et al., 2020). The findings further indicate that integrating these technologies with Building Information Modelling (BIM), Internet of Things (IoT) devices, cloud computing, and Large Language Models (LLMs) establishes a unified digital ecosystem capable of supporting adaptive, resilient, and data-driven architectural practice. Autonomous design systems not only improve productivity by reducing repetitive computational tasks but also strengthen sustainability through energy optimization, lifecycle performance evaluation, material efficiency, and intelligent environmental analysis. Rather than replacing architects, these technologies increasingly function as collaborative partners that augment professional creativity, analytical reasoning, and strategic decision-making, allowing designers to concentrate on innovation, user experience, and contextual problem-solving (OpenAI, 2023; Bommasani et al., 2022). Despite these technological advances, the review identified several critical challenges that continue to influence successful implementation.

Issues relating to explainability, interoperability, algorithmic bias, cybersecurity, intellectual property ownership, legal accountability, data governance, and regulatory compliance require coordinated attention from researchers, software developers, professional bodies, and policymakers. The integration of Explainable Artificial Intelligence (XAI), standardized data exchange frameworks, secure cloud infrastructures, ethical governance principles, and human-in-the-loop decision models will therefore be essential for ensuring trustworthy and responsible deployment of autonomous design systems (Floridi & Cowls, 2022; NIST, 2023). Overall, this study concludes that the convergence of Generative AI, Computer Vision, and Digital Twins represents a transformative paradigm for smart architectural practice. As these technologies continue to mature, autonomous design systems will evolve into intelligent collaborative environments capable of continuously learning from real-time data, predicting building performance, optimizing resource utilization, and supporting sustainable urban development. Their successful adoption will depend not only on technological innovation but also on transparent governance, interdisciplinary collaboration, professional competency development, and ethical AI implementation, thereby positioning architects and intelligent systems as complementary partners in shaping the future of resilient, human-centered, and sustainable built environments.

FUTURE RESEARCH

Future research should prioritize the development of explainable, trustworthy, and interoperable autonomous design systems that seamlessly integrate Generative AI, Computer Vision, Digital Twins, BIM, and IoT within unified architectural platforms. Longitudinal empirical studies involving real-world architectural projects are needed to validate the operational effectiveness, scalability, and economic benefits of these technologies across diverse geographical and regulatory contexts. Further investigations should explore federated learning, edge AI, quantum-enhanced optimization, robotics, and 6G-enabled smart infrastructure to improve real-time decision intelligence and computational efficiency. Research should also establish standardized evaluation metrics, ethical governance frameworks, and international interoperability standards that address cybersecurity, intellectual property, algorithmic fairness, and professional accountability. Such advances will accelerate the responsible adoption of autonomous architectural ecosystems capable of supporting resilient, sustainable, and intelligent built environments.

References

Attia, S. (2022). *Net Zero Energy Buildings (NZEB): Concepts, Frameworks, and Roadmap for Project Analysis and Implementation*. Butterworth-Heinemann.

Boje, C., Guerriero, A., Kubicki, S., & Rezgui, Y. (2020). Towards a semantic construction digital twin: Directions for future research. *Automation in Construction, 114*, 103179. https://doi.org/10.1016/j.autcon.2020.103179

Bock, T., & Linner, T. (2015). *Robot-Oriented Design: Design and Management Tools for the Deployment of Automation and Robotics in Construction*. Cambridge University Press.

Bommasani, R., Hudson, D. A., Adeli, E., et al. (2022). *On the Opportunities and Risks of Foundation Models*. Stanford Centre for Research on Foundation Models. https://doi.org/10.48550/arXiv.2108.07258

Bosché, F., Ahmed, M., Turkan, Y., Haas, C. T., & Haas, R. (2015). The value of integrating Scan-to-BIM and Scan-vs-BIM techniques for construction monitoring using laser scanning and BIM: The case of cylindrical MEP components. *Automation in Construction, 49*, 201-213. https://doi.org/10.1016/j.autcon.2014.05.014

buildingSMART International. (2023). *Industry Foundation Classes (IFC) 4.3:Open BIM Standard*.

Deb, K. (2001). *Multi-Objective Optimization Using Evolutionary Algorithms*. John Wiley & Sons.

Dosovitskiy, A., Beyer, L., Kolesnikov, A., et al. (2021). An image is worth 16 × 16 words: Transformers for image recognition at scale. *International Conference on Learning Representations (ICLR)*. https://doi.org/10.48550/arXiv.2010.11929

Eastman, C., Teicholz, P., Sacks, R., & Liston, K. (2018). *BIM Handbook: A Guide to Building Information Modeling for Owners, Designers, Engineers, Contractors, and Facility Managers* (3rd ed.). John Wiley & Sons.

European Commission. (2024). *Regulation (EU) 2024/1689 of the European Parliament and of the Council of 13 June 2024 Laying Down Harmonised Rules on Artificial Intelligence (Artificial Intelligence Act).*

Floridi, L., & Cowls, J. (2022). A unified framework of five principles for AI in society. *Harvard Data Science Review, 1*(1). https://doi.org/10.1162/99608f92.8cd550d1

Fuller, A., Fan, Z., Day, C., & Barlow, C. (2020). Digital twin: Enabling technologies, challenges and open research. *IEEE Access, 8*, 108952-108971. https://doi.org/10.1109/ACCESS.2020.2998358

Goodfellow, I., Bengio, Y., & Courville, A. (2016). *Deep Learning*. MIT Press.

Guo, H., Tang, S., Li, H., & Luo, X. (2024). Artificial intelligence and computer vision applications in smart construction: A systematic review. *Automation in Construction, 164*, 105512. https://doi.org/10.1016/j.autcon.2024.105512

Hamledari, H., & McCabe, B. (2018). Automated computer vision-based detection of construction progress using 4D BIM. *Automation in Construction, 93*, 4457. https://doi.org/10.1016/j.autcon.2018.05.014

He, K., Gkioxari, G., Dollár, P., & Girshick, R. (2017). Mask R-CNN. In *Proceedings of the IEEE International Conference on Computer Vision (ICCV)* (pp. 2961-2969). https://doi.org/10.1109/ICCV.2017.322

International Organization for Standardization. (2018). *ISO 19650-1:2018: Organization and Digitization of Information About Buildings and Civil Engineering Works, Including Building Information Modelling (BIM), Information Management Using Building Information Modelling, Part 1: Concepts and Principles.*

Jones, D., Snider, C., Nassehi, A., Yon, J., & Hicks, B. (2020). Characterising the digital twin: A systematic literature review. *CIRP Journal of Manufacturing Science and Technology, 29*, 36-52. https://doi.org/10.1016/j.cirpj.2020.02.002

Khan, A., Alzahrani, A., Alqahtani, S., & Albogamy, F. (2024). Deep learning and computer vision for intelligent infrastructure monitoring: A comprehensive review. *IEEE Access, 12*, 45871-45905. https://doi.org/10.1109/ACCESS.2024.3375809

Li, C., Zhang, T., Du, X., Zhang, Y., & Xie, H. (2024). *Generative AI Models for Different Steps in Architectural Design: A Literature Review*. arXiv. https://doi.org/10.48550/arXiv.2404.01335

Lu, Q., Xie, X., Parlikad, A. K., & Schooling, J. M. (2020). Digital twin-enabled anomaly detection for built asset monitoring in operation and maintenance. *Automation in Construction, 118*, 103277. https://doi.org/10.1016/j.autcon.2020.103277

Lundberg, S. M., & Lee, S.-I. (2017). A unified approach to interpreting model predictions. *Advances in Neural Information Processing Systems, 30*, 4765-4774.

National Institute of Standards and Technology. (2023). *Artificial Intelligence Risk Management Framework (AI RMF 1.0)*. https://doi.org/10.6028/NIST.AI.100-1

OpenAI. (2023). *GPT-4 Technical Report*. arXiv. https://doi.org/10.48550/arXiv.2303.08774 (arXiv)

Page, M. J., McKenzie, J. E., Bossuyt, P. M., et al. (2021). The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *The BMJ, 372*, n71. https://doi.org/10.1136/bmj.n71

Rasheed, A., San, O., & Kvamsdal, T. (2020). Digital twin: Values, challenges and enablers from a modeling perspective. *IEEE Access, 8*, 21980-22012. https://doi.org/10.1109/ACCESS.2020.2970143

Redmon, J., Divvala, S., Girshick, R., & Farhadi, A. (2016). You only look once: Unified, real-time object detection. In *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition (CVPR)* (pp. 779-788). https://doi.org/10.1109/CVPR.2016.91

Ribeiro, M. T., Singh, S., & Guestrin, C. (2016). "Why should I trust you?": Explaining the predictions of any classifier. In *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining* (pp. 1135-1144). https://doi.org/10.1145/2939672.2939778

Rombach, R., Blattmann, A., Lorenz, D., Esser, P., & Ommer, B. (2022). High-resolution image synthesis with latent diffusion models. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)* (pp. 10684-10695). https://doi.org/10.1109/CVPR52688.2022.01042

Russell, S., & Norvig, P. (2021). *Artificial Intelligence: A Modern Approach* (4th ed.). Pearson.

Sutton, R. S., & Barto, A. G. (2018). *Reinforcement Learning: An Introduction* (2nd ed.). MIT Press.

United Nations Environment Programme. (2024). *Global Status Report for Buildings and Construction 2024*.

Xu, H., Omitaomu, F., Sabri, S., Zlatanova, S., Li, X., & Song, Y. (2024). Leveraging generative AI for urban digital twins: A scoping review on the autonomous generation of urban data, scenarios, designs, and 3D city models for smart city advancement. *Urban Informatics, 3*, Article 29. https://doi.org/10.1007/s44212-024-00060-w

MATEMATİK EĞİTİMİNDE YAPAY ZEKA ENTEGRASYONU: ÖZ BELİRLEME TEORİSİ TEMELİNDE BÜTÜNLEŞTİRİCİ BİR ÇERÇEVE

ARTIFICIAL INTELLIGENCE INTEGRATION IN MATHEMATICS EDUCATION: AN INTEGRATIVE FRAMEWORK GROUNDED IN SELF-DETERMINATION THEORY

Prof. Dr. Murat Subasi[1]
[1]Department of Mathematics, Science Faculty, Ataturk University, Erzurum, Türkiye
ORCID NO: 0000-0001-5895-143X

Assoc. Prof. Dr. Serpil Yorganci[1,2]
[1] Vocational College of Technical Sciences, Ataturk University, Erzurum, Türkiye
[2] Department of Mathematics and Science Education, Kazim Karabekir Education Faculty, Atatürk University, Erzurum, Türkiye
ORCID NO: 0000-0001-7284-8340

ÖZET

Bu çalışma, matematik eğitiminde yapay zeka (YZ) kullanımına ilişkin mevcut literatürü, Öz Belirleme Teorisi (Self-Determination Theory) perspektifinden ele alarak bütünleştirici bir teorik çerçeve sunmayı amaçlamaktadır. Günümüzde üretken yapay zeka modellerinin eğitim alanında hızla yaygınlaşması, öğretim ve öğrenme süreçlerini dönüştüren en önemli faktörlerden biri haline gelmiştir. Matematik eğitimi, soyut kavramları, sembolik akıl yürütmeyi ve çok adımlı problem çözme becerilerini içeren doğası gereği yüksek bilişsel talep barındıran bir disiplindir. YZ araçlarının bu bağlama entegrasyonu, öğrencilerin bilgiyi işleme biçimlerini, problem çözme stratejilerini ve öğrenme çıktılarını doğrudan etkileme kapasitesine sahiptir.

Çalışmanın kavramsal temellerinde, eğitim bağlamında YZ, Akıllı Öğretim Sistemleri, Üretken Yapay Zeka, Uyarlanabilir Öğrenme Sistemleri ve Bilgisayar Cebir Sistemleri olmak üzere dört ana kategoride tanımlanmıştır. Bu çalışmanın özgün katkısı, YZ destekli öğrenme ortamlarını Öz Belirleme Kuramı'nın üç temel psikolojik ihtiyacı; özerklik, yeterlilik ve ilişkililik açısından analiz etmesidir. Öz Belirleme Teorisine göre, bu üç ihtiyacın karşılanması bireyin içsel motivasyonunu ve psikolojik iyilik halini doğrudan desteklemektedir. Çalışma, YZ araçlarının öğrencilerin özerklik ihtiyacını nasıl destekleyebileceğini (örneğin, kişiselleştirilmiş öğrenme yolları sunarak), yeterlilik ihtiyacını nasıl karşılayabileceğini (anında ve geliştirici geri bildirimle) ve ilişkililik ihtiyacını nasıl güçlendirebileceğini (işbirlikçi öğrenme ortamları ve etkileşimli diyaloglar aracılığıyla) tartışmaktadır.

Bu bütünleştirici çerçevenin, etkili ve sürdürülebilir YZ entegrasyon stratejilerinin geliştirilmesine katkı sağlaması ve gelecekteki ampirik araştırmalar için test edilebilir hipotezler üretmesi hedeflenmektedir.

Anahtar Kelimeler: Yapay zeka, Matematik, Öz Belirleme Kuramı

ABSTRACT

This study aims to provide an integrative theoretical framework by examining the existing literature on the use of artificial intelligence (AI) in mathematics education from the perspective of Self-Determination Theory. The rapid proliferation of generative AI models in education has become one of the most important factors transforming teaching and learning processes. Mathematics education is a discipline that inherently demands high cognitive abilities, involving abstract concepts, symbolic reasoning, and multi-step problem-solving skills. The integration of AI tools into this context has the capacity to directly influence how students

process information, their problem-solving strategies, and their learning outcomes. In the conceptual framework of this study, AI in the educational context is defined in four main categories: Intelligent Tutoring Systems, Generative AI, Adaptive Learning Systems, and Computational Algebra Systems. The unique contribution of this study is its analysis of AI-supported learning environments in terms of the three fundamental psychological needs of Self-Determination Theory: autonomy, competence, and relatedness. According to Self-Determination Theory, meeting these three needs directly supports an individual's intrinsic motivation and psychological well-being. This study discusses how AI tools can support students' need for autonomy (e.g., by offering personalized learning paths), meet their need for competence (with immediate and developmental feedback), and strengthen their need for relatedness (through collaborative learning environments and interactive dialogues). The aim is for this integrative framework to contribute to the development of effective and sustainable AI integration strategies and generate testable hypotheses for future empirical research.

Keywords: Artificial intelligence, Mathematics education, Self-Determination Theory

GİRİŞ

1955 yılında Profesör John McCarthy tarafından "akıllı makineler oluşturmak için bilim ve mühendisliği kullanan bir sistem" (McCarthy et al., 2006) olarak tanımlanan yapay zeka terimi, günümüzde en genel tanımıyla bilgisayarların ve makinelerin insan zekasını ve problem çözme yeteneklerini simüle etmesine olanak tanıyan teknolojiler bütünü olarak (IBM, 2023) ifade edilmektedir. Genel olarak bu sistemlerin, geçmişteki yaşanmışlıklardan öğrenebilme ve bu tecrübelerden elde ettiği verileri analiz etme, yorumlama ve karar verme yeteneğine sahip olduğu yaygın kabul görmektedir (Hooda vd., 2022). Tek bir teknolojiyi ifade etmek yerine, makine öğrenmesi (ML), doğal dil işleme (NLP), veri madenciliği, sinir ağları, bilgisayarlı görü ve robotik gibi birçok alt alanı kapsayan bir şemsiye terim olan yapay zeka, eğitimde Akıllı Öğretim Sistemlerinin (ITS) ve kişiselleştirilmiş öğrenme platformlarının da temelini oluşturmaktadır (Dong vd, 2025; Huang vd, 2023; Liu vd, 2025).

Özellikle matematik eğitimi, soyut kavramları, prosedürel ve kavramsal bilgi arasındaki dengeyi, sembolik akıl yürütmeyi ve çok adımlı problem çözme becerilerini içeren doğası gereği yüksek bilişsel talep barındıran bir disiplin (Walkington, 2025) olduğu için yapay zeka araçlarının bu bağlama entegrasyonu, yalnızca teknolojik bir yenilik olmanın ötesinde, öğrencilerin bilgiyi işleme biçimlerini, problem çözme stratejilerini ve öğrenme çıktılarını doğrudan etkileme kapasitesine sahiptir. Bu nedenle, YZ'nin matematik eğitimindeki rolünün kapsamlı bir şekilde anlaşılması, etkili ve sürdürülebilir entegrasyon stratejilerinin geliştirilmesi için kritik öneme sahiptir (Guo & Zhang, 2025; Park & Lee, 2025).

MATEMATİK EĞİTİMİNDE YAPAY ZEKANIN TARİHSEL GELİŞİMİ

Matematik eğitiminde yapay zeka kullanımı yıllara göre önemli bir artış sergilemektedir. Örneğin, Kuncoro ve arkadaşlarının (2026) 2001-2025 yılları arasını kapsayan geniş ölçekli

bibliyometrik analiz (n=237) çalışmasında, matematik eğitiminde yapay zeka ve akıllı öğretim sistemleri araştırmalarının net bir büyüme eğilimi izlediği belirlenmiştir (Şekil 1). Yayın eğilimi incelendiğinde, 2010 yılına kadar süren durgun keşif evresinden sonra özellikle 2019 sonrasında makine öğrenmesi ile büyük dil modellerindeki ilerlemelere bağlı olarak 2024 yılında yaklaşık 50 yayına ulaşan üstel bir büyüme kaydedilmiştir. Araştırma verimliliğinde ABD, Çin ve Almanya lider ülkeler olarak öne çıkarken; yazar üretkenliğinin Lotka Yasası'na uygun şekilde küçük bir çekirdek grup etrafında toplandığı ve alanın "bilgisayar destekli öğretim" gibi klasik yaklaşımlardan "kişiselleştirilmiş öğrenme", "üretken yapay zeka" ve "federasyonlaşmış öğrenme" gibi modern temalara evrildiği görülmektedir. Son alanyazın, matematik eğitiminin geleneksel yöntemlerden veriye dayalı, adaptif ve akıllı öğrenme ekosistemlerine doğru kalıcı bir dönüşüm geçirdiğini ortaya koymaktadır (Canonigo, 2024; Kuo vd, 2025; Liu vd, 2025; Xuan vd, 2025). Bu bağlamda kullanılan yapay zeka teknolojisine göre öğrenme sürecine etkisi de farklılaşmaktadır. Matematik eğitiminde kullanılan yapay zeka (YZ) çeşitleri, temel olarak öğrenciye sunulan desteğin niteliğine ve teknolojinin işlevine göre kategorize edilmektedir. Bu uygulamalar, üstlendikleri pedagojik roller ve sundukları teknik imkanlar bakımından aşağıda ayrıntılı olarak ele alınmıştır.



Şekil 1. Matematik eğitiminde yapay zeka ve akıllı öğretim sistemleri üzerine yıllık bilimsel yayın eğilimi, 2001-2025

Kaynak: Kuncoro, K. S., Fitriasari, P., Erlangga, S. Y., Ario, M., & Kartika, D. L. (2026). Artificial intelligence and intelligent tutoring systems in mathematics education: A bibliometric analysis (2001–2025). *F1000Research*, *15*(107).

MATEMATİK EĞİTİMİNDE YAPAY ZEKA TEKNOLOJİLERİ VE UYGULAMA ALANLARI

Matematik eğitiminde kullanılan yapay zeka (YZ) çeşitleri, öğrenciye sunulan pedagojik desteğin niteliğine ve teknolojinin işlevine göre şu şekilde detaylandırılabilir:

Akıllı Öğretim Sistemleri (ITS) ve Adaptif Öğrenme Platformları

Matematik eğitimindeki en köklü uygulamalardan biri, öğrencilerin bilişsel ve duyuşsal durumlarını modelleyerek öğretimi kişiselleştiren Akıllı Öğretim Sistemleri (ITS) ve Adaptif Öğrenme Platformlarıdır (Ateş, 2025; Awang vd., 2025). Bu sistemler; öğrencinin bilgi düzeyini, öğrenme hızını ve karşılaştığı zorlukları analiz ederek kişiye özel öğrenme yolları, anlık geri bildirimler ve uyarlanabilir iskeleleme mekanizmaları sunmaktadır (Awang vd., 2025; Kuncoro vd., 2026;). Örneğin; ALEKS, Squirrel AI, MATHia, PAT2Math ve GeoGebra Tutor gibi platformlar, matematiksel akıl yürütme süreçlerini adım adım takip ederek öğrencileri "Yakınsal Gelişim Alanı" içinde tutmakta ve başarı farklarını azaltmaya yardımcı olmaktadır (Awang vd., 2025; Khazanchi vd., 2024; Kuncoro vd., 2026; Yorganci, 2018). Bu araçlar, öğretmenlere sundukları analitik raporlar sayesinde sınıf içindeki bireysel farklılıkların daha etkin yönetilmesine ve teşhis edici değerlendirmelerin yapılmasına imkan tanımaktadır (Awang ve ark., 2025; Feng ve ark., 2025; Kuncoro ve ark., 2026)

Üretken Yapay Zeka (GenAI) ve Büyük Dil Modelleri (LLM)

Son yıllarda matematik eğitiminde devrimsel bir etki yaratan bir diğer kategori ise Üretken Yapay Zeka (GenAI) ve Büyük Dil Modelleridir (LLM) (Chamola ve ark., 2026; Wardat ve ark., 2023). ChatGPT, Bard ve Claude gibi araçlar; doğal dil işleme yetenekleri sayesinde öğrencilerle Sokratik bir diyalog kurarak karmaşık matematik problemlerinin çözümünde rehberlik etmekte ve kavramsal açıklamalar oluşturmaktadır (Chamola vd, 2026; Kuo vd., 2026; Zhou vd, 2025). Bu araçlar, öğrencinin doğrudan cevaba ulaşması yerine çözüm stratejileri geliştirmesine, mantıksal hatalarını fark etmesine ve problemlerin farklı çözüm yollarını keşfetmesine yardımcı olan dijital bir "öğrenme arkadaşı" (learning companion) olarak konumlanmaktadır (Awang vd., 2025; Kuo vd., 2026; Zhou vd., 2025). Ayrıca GenAI teknolojilerinin, soyut matematik kavramlarını günlük yaşam senaryolarıyla ilişkilendirerek matematik kaygısını azaltmada ve öğrenci motivasyonunu pekiştirmede kritik bir rol oynadığı saptanmıştır (Kuo vd., 2026; Liu vd., 2025; Xuan vd., 2025).

Bilgisayar Cebir Sistemleri (CAS) ve Akıllı Matematik Çözücüler

Matematik eğitimindeki bir diğer temel teknoloji grubu, sembolik hesaplamaları otomatize eden ve karmaşık yapıları görselleştiren Bilgisayar Cebir Sistemleri (CAS) ile yapay zeka tabanlı matematik çözücülerdir (Awang vd., 2025; Mohamed vd., 2022). Photomath, Microsoft Math Solver, Maple ve MATLAB gibi sistemler; matematiksel ifadeleri taranabilir hale getirerek öğrencilere anlık geri bildirim, adım adım yapılandırılmış çözümler ve kavramsal açıklamalar sunmaktadır (Awang vd., 2025; Fissore vd., 2021). Bu araçlar, özellikle kalkülüs, lineer cebir ve diferansiyel denklemler gibi yüksek düzeyde soyutlama gerektiren konularda

grafiksel sezgi kazandırmak ve öğrencilerin işlemsel hatalara takılmadan problem çözme stratejilerine odaklanmalarını sağlamak amacıyla kullanılmaktadır (Chamola vd., 2026; Fissore vd., 2021). Güncel araştırmalar, bu sistemlerin büyük dil modelleriyle (LLM) entegre edilerek SymPy veya Wolfram Language gibi sembolik motorlar üzerinden matematiksel doğruluğu artırdığını ve öğrencilerin Sokratik bir dille sorgulama yapabileceği akıllı asistanlara dönüştüğünü vurgulamaktadır (Chamola vd., 2026). CAS ve akıllı çözücülerin kullanımı, öğrencilerin görsel geri bildirimler aracılığıyla bilişsel yükünü azaltarak karmaşık matematiksel modelleme görevlerine daha fazla kaynak ayırmalarına imkan tanımaktadır (Chamola vd., 2026)

Artırılmış Gerçeklik (AR), Sanal Gerçeklik (VR) ve Eğitsel Robotik Sistemler

Matematik öğretimini daha somut ve etkileşimli hale getiren diğer önemli teknolojiler ise Artırılmış Gerçeklik (AR), Sanal Gerçeklik (VR) ve Eğitsel Robotik sistemleridir (Awang vd, 2025; Ateş, 2025; Mohamed vd., 2022). GeoGebra AR, cleARmaths ve CalcVR gibi uygulamalar; özellikle geometri, uzamsal akıl yürütme ve kalkülüs gibi yüksek düzeyde soyutlama gerektiren konularda 3 boyutlu görselleştirmeler sunarak öğrencilerin kavramları deneyimleyerek öğrenmesini sağlamaktadır (Awang vd., 2025). Bu sarmal teknolojiler, öğrencilerin karmaşık şekilleri ve fonksiyonları sanal bir ortamda manipüle etmelerine izin vererek "keşfederek öğrenme" yaklaşımını desteklemektedir (Ateş, 2025; Awang vd., 2025)

Ancak, bu sistemlerin sürdürülebilir olması için teknolojinin ötesinde, öğretmenlerin pedagojik hazırbulunuşluk (Kuncoro vd., 2026), araçların müfredat hedefleriyle tam uyumlu olması (Kuo vd, 2026), sistematik öğretim tasarımı, veri gizliliği, algoritmik şeffaflık ile etik standartlar çerçevesinde güven inşa etmesi (Akgun & Greenhow, 2022; Chamola vd, 2026) gibi unsurlar kritik rol oynamaktadır. Ayrıca, yapay zeka sistemlerinin teknik birer araç olmaktan çıkıp etkili öğrenme partnerlerine dönüşebilmesi için, öğrencinin motivasyonunu dışsal zorunluluklardan içsel motivasyona kaydıracak psikolojik bir zemine ihtiyaç duyulmaktadır (Dou & Sun, 2025; Ellikkal & Rajamohan, 2025; Ryan & Deci, 2020; Xia vd., 2022). YZ'nin sunduğu kişiselleştirilmiş öğrenme yolları, eğer öğrencinin özerklik (autonomy) ve yetkinlik (competence), ve ilişkililik (relatedness) ihtiyaçlarına yanıt vermiyorsa, öğrenci kendisini sistem tarafından yönlendirilen "cansız bir çip" gibi hissedebilir (Pham & Sampson, 2022). Önceki çalışmalar, Öz Belirleme Teorisi'nin bu potansiyel mekanizmaları açıklamak için güçlü ve sade bir çerçeve sağladığını (Ryan & Deci 2020) ve psikolojik ihtiyaçların karşılanmasının içsel motivasyonu artırmaya katkıda bulunduğunu öne sürmektedir. Ancak literatür incelendiğinde, mevcut yapay zeka uygulamalarının SDT perspektifinden bakıldığında ciddi eksiklikler barındırdığı görülmektedir (Dou & Sun, 2025; Mohamed vd., 2022; Shi & Zhang, 2025; Xu vd., 2026; Yan vd., 2024). Bu nedenle, bu çalışmada, Öz Belirleme Kuramı'nın üç temel psikolojik ihtiyacını (özerklik, yeterlilik ve ilişkililik) karşılayacak biçimde yapılandırılmış YZ destekli bir öğrenme ortamının tasarımına odaklanılmıştır.

ÖZ BELİRLEME TEORİSİ (ÖBT): YAPAY ZEKA DESTEKLİ ÖĞRENMEDE ÖZERKLİK, YETERLİLİK, İLİŞKİLİLİK

ÖBT, teorik ve ampirik bir yaklaşım olarak insan motivasyonuna dair popüler bir teori olup, insanların özgürce yaptıkları eylemlerin altında yatan motivasyonel dinamikler ile zorla yaptırılan veya baskı altında hissettikleri dinamikler arasında ayrım yapmak olarak tanımlanır (Deci 1992). Teori, dışsal düzeyden içsel düzeye kadar farklı özerk öz düzenleme düzeylerini etkileyen farklı motivasyon türleri olduğunu savunur. Özünde, tüm bireylerin onları harekete geçmeye veya geçmemeye yönlendiren üç temel psikolojik ihtiyaca sahip olduğunu öne sürer: özerklik, yeterlilik ve ilişkililik. Özerklik, bir kişinin dış faktörlerden bağımsız olarak, görevle ilgili bir bağlamda özerk bir şekilde karar verme, seçim yapma ve katılım yeteneği olarak tanımlanır. Yeterlilik, öğrencilerin etkili eylemlerde bulunma ve öğrenme sürecinde yeteneklerini geliştirme yeteneğini ifade eder. İlişkililik, bireyin çevresiyle sağlıklı ve güvene dayalı ilişkiler kurma, işbirliği ve iletişim kurma yeteneği olarak tanımlanır (Ryan ve Deci 2000). Bu üç psikolojik ihtiyaç karşılandığında, insanlar psikolojik olarak daha iyi hissederler; öte yandan, ihtiyaçları karşılanmadığında aşırı derecede parçalanmış, yalnız ve tepkisel hale gelirler (Deci ve Ryan 1985).

Bu temel psikolojik ihtiyaçların karşılanması, öğrenme ortamında kullanılan teknolojilerin özelleikleriyle uyumlu olduğunda, öğrenme üst düzeyde gerçekleşebilmektedir. Bu bağlamada, bu çalışmada yapay zeka araçları, öğrencilerin matematiksel içerikle etkileşim biçimini dönüştürebilecek ihtiyaç destekleyici öğrenme olanakları olarak kavramsallaştırılmıştır. Şekil 2, yapay zeka uygulamalarının bu ihtiyaç destekleyici özelliklerini özetlemektedir. Buna göre, bu uygulamalarının özel işlevleri, her bir SDT ihtiyacıyla tanımlanabilir şekillerde uyumludur.



Şekil. 2. Yapay zeka uygulamalarının özerklik, yetkinlik ve ilişkililikle uyumlu, ihtiyaç odaklı destekleyici özellikleri.

Kaynak: Yorganci, S. (2026). Evaluating Students' Learning Experiences in Blended Learning Environment Integrated With Interactive GeoGebra Applets: A Self-Determination Theory Perspective. Journal of Computer Assisted Learning, 42(4), e70283.

Özerklik desteği: Yapay zeka destekliöğrenme ortamında, özerklik desteği, giriş çubukları ve sesli komut özellikleri aracılığıyla geliştirilmiştir. Bu tasarım, öğrencilere irade ve seçim duygusu vererek, sabit, doğrusal bir sunumla kısıtlanmış hissetmeden kendi hızlarında öğrenmelerine olanak tanır.

Yeterlilik desteği: Anlık geri bildirim, adım adım çözüm gösterimleri, hata vurgulama ve etkileşimli alıştırmalar aracılığıyla ustalık duygusu pekiştirilir. Bu özellikler, öğrencilerin eylemlerinin sonuçlarını anında görebildikleri ve yanlış anlamaları gecikmeden düzeltebildikleri için bağımsız olarak başarıya ulaşmalarını sağlar.

İlişkililik desteği: Öğrenme sürecinin sosyal boyutu, işbirlikçi uygulamalar, canlı geri bildirim araçları, paylaşılabilir uygulamalar ve kaydedilmiş öğrenci dosyaları aracılığıyla sürdürülür. Bu özellikler, eğitmen varlığını destekler ve akran etkileşimini teşvik ederek, öğreme ortamlarında bağlantı duygusu sağlar (Yorganci, 2026).

SONUÇ

Bu çalışma yapay zeka araçlarının matematik eğitiminde teknik birer özellik kümesi olarak değil, öğrencinin içsel motivasyonunu besleyen ihtiyaç-destekleyici bileşenler olarak yeniden kavramsallaştırılması gerektiğini savunmaktadır. Alan yazındaki uygulama örüntüsü göz önüne alındığında, gelecekteki tasarım ve araştırma çabalarının üç psikolojik ihtiyacı eş zamanlı gözeten bütüncül modeller geliştirmeye yönelmesi, hem öğrenci motivasyonunun hem de öğrenme çıktılarının sürdürülebilir biçimde desteklenmesi açısından kritik önem taşımaktadır.

KAYNAKÇA

Akgun, S., & Greenhow, C. (2022). Artificial intelligence in education: Addressing ethical challenges in K-12 settings. AI and Ethics, 2(3), 431–440. https://doi.org/10.1007/s43681-021-00096-7

Ateş, H. (2025). Integrating augmented reality into intelligent tutoring systems to enhance science education outcomes. *Education and Information Technologies, 30*(4), 4435–4470

Awang, L. A., Yusop, F. D., & Danaee, M. (2025). Current practices and future direction of artificial intelligence in mathematics education: A systematic review. International Electronic *Journal of Mathematics Education, 20*(2), em0823. https://doi.org/10.29333/iejme/16006

Canonigo, A. M. (2024). Levering AI to enhance students' conceptual understanding and confidence in mathematics. *Journal of Computer Assisted Learning, 40(*6), 3215–3229

Deci, E. L. 1992. "The Relation of Interest to the Motivation of Behavior: A Self Determination Theory Perspective." In *The Role of Interest in Learning and Development, edited by K. A. Renninger, S.* Hidi, and A. Krapp, 43–70. Lawrence Erlbaum Associates.

Deci, E. L., & R. M. Ryan. 1985. *Intrinsic Motivation and SelfDetermination in Human Behavior.* Plenum Press

Dong, L., Tang, X., & Wang, X. (2025). Examining the effect of artificial intelligence in relation to students' academic achievement: A meta-analysis. *Computers and Education: Artificial Intelligence*, 8, 100400

Dou, W., & Sun, X. (2025). Artificial intelligence tools: Improvement of motivation, psychological well-being, and psychological capital of EFL learners: A self-determination theory perspective. *Learning and Motivation, 92,* 102169.

Guo, X., & Zhang, J. (2025). Utilizing AI Technology to Enhance Primitive Idea-Driven Mathematics Instruction. *ICNISC 2025*. ACM.

Hooda, M., Rana, C., Dahiya, O., Rizwan, A., & Hossain, M. S. (2022). Artificial intelligence for assessment and feedback to enhance student success in higher education. *Mathematical Problems in Engineering,* 1–19. https://doi.org/10.1155/2022/5215722

Huang, A. Y. Q., Lu, O. H. T., & Yang, S. J. H. (2023). Effects of artificial Intelligence–enabled personalized recommendations on learners' learning engagement, motivation, and outcomes in a flipped classroom. *Computers & Education, 194*, 104684

IBM. (2023). What is AI? https://www.ibm.com/topics/artificial-intelligence

Kuncoro, K. S., Fitriasari, P., Erlangga, S. Y., Ario, M., & Kartika, D. L. (2026). Artificial intelligence and intelligent tutoring systems in mathematics education: A bibliometric analysis (2001–2025). *F1000Research, 15*(107). https://doi.org/10.12688/f1000research.176098.1

Kuo, B. C., Bai, Z. E., & Lin, C. H. (2026). Developing an AI learning companion for mathematics problem solving in elementary schools. *Computers & Education, 240*, 105463

Liu, J., Sun, D., Sun, J., Wang, J., & Ho, P. L. (2025). Designing a generative AI enabled learning environment for mathematics word problem solving in primary schools: Learning performance, attitudes and interaction. *Computers and Education: Artificial Intelligence, 9*, 100438

McCarthy, J., Minsky, M. L., Rochester, N., & Shannon, C. E. (2006). A proposal for the dartmouth summer research project on artificial intelligence, august 31, 1955. *AI Magazine, 27*(4). https://doi.org/10.1609/aimag.v27i4.1904

Mohamed, M. Z., Hidayat, R., binti Suhaizi, N. N., bin Mahmud, M. K. H., & binti Baharuddin, S. N. (2022). Artificial intelligence in mathematics education: A systematic literature review. *International Electronic Journal of Mathematics Education, 17*(3), em0694.

Park, J., & Lee, K. (2025). AI-based mathematics education: A review of issues in international research. *Korean Journal of Educational Research, 62*,(4), 123-145.

Pham, S. T., & Sampson, P. M. (2022). The development of artificial intelligence in education: A review in context. *Journal of Computer Assisted Learning, 38*(5), 1408-1421.

Ryan, R. M., & E. L. Deci. (2020). Intrinsic and Extrinsic Motivation From a Self-Determination Theory Perspective. Definitions, Theory, Practices, and Future Directions. *Contemporary Educational Psychology, 61*: 101860. https://doi.org/10.1016/j.cedpsych.2020.101860

Shi, S., & Zhang, H. (2025). EFL students' motivation predicted by their self-efficacy and resilience in artificial intelligence (AI)-based context: From a self-determination theory perspective. *Learning and Motivation, 91*, 102151.

Walkington, C. (2025). The implications of generative artificial intelligence for mathematics education. *School Science and Mathematics.*

Xuan, S. H., Vo, D. T., Pham, T. H., Trung, H. T., & Nhat, M. V. (2025). Evaluating the Impact of Generative AI in Mathematics Education: A Comparative Study. *Human Behavior and Emerging Technologies,* Article 8886206

Yan, L., Sha, L., Zhao, L., Li, Y., Martinez-Maldonado, R., Chen, G., ... & Gašević, D. (2024). Practical and ethical challenges of large language models in education: A systematic scoping review. *British Journal of Educational Technology, 55*(1), 90-112.

Yorganci, S. (2018). A Study on the Views of Graduate Students on the Use of GeoGebra in Mathematics Teaching. *European Journal of Education Studies 4*(8), 63–78. https://doi.org/10.5281/zenodo.1272935

Yorganci, S. (2026). Evaluating Students' Learning Experiences in Blended Learning Environment Integrated With Interactive GeoGebra Applets: A Self-Determination Theory Perspective. *Journal of Computer Assisted Learning, 42*(4), e70283.

YAPAY ZEKA DESTEKLİ MATEMATİK ÖĞRENİMİNDE BİLİŞSEL YÜKÜN YÖNETİMİ

MANAGING COGNITIVE LOAD IN AI-SUPPORTED MATHEMATICS LEARNING

Prof. Dr. Murat Subasi[1]
[1]Department of Mathematics, Science Faculty, Ataturk University, Erzurum, Türkiye
ORCID NO: 0000-0001-5895-143X

Assoc. Prof. Dr. Serpil Yorganci[1,2]
[1] Vocational College of Technical Sciences, Ataturk University, Erzurum, Türkiye
[2] Department of Mathematics and Science Education, Kazim Karabekir Education Faculty, Atatürk University, Erzurum, Türkiye
ORCID NO: 0000-0001-7284-8340

ÖZET

Bu çalışma, yapay zeka (YZ) destekli matematik öğreniminde bilişsel yükün yönetimine ilişkin kapsamlı bir teorik çerçeve sunmayı amaçlamaktadır. Son yıllarda ChatGPT, Wolfram Alpha ve Photomath gibi YZ tabanlı araçların eğitim alanında, özellikle matematik öğretiminde kullanımı hızla yaygınlaşmaktadır. Bu araçlar, öğrencilere anında geri bildirim sağlama, adım adım çözüm sunma ve karmaşık problemleri basitleştirme potansiyeli ile dikkat çekmektedir. Ancak, matematik eğitimi soyut kavramları, çok adımlı prosedürleri ve yoğun problem çözme becerilerini içeren doğası gereği yüksek bilişsel talep barındıran bir disiplindir. YZ araçlarının bilinçsiz veya uygun olmayan pedagojik stratejilerle kullanımı, öğrenme sürecinde önemli engeller oluşturabilir. Özellikle, bu araçların sunduğu bilgi akışının ve etkileşim biçimlerinin, öğrencinin sınırlı çalışma belleği kapasitesini zorlayacak şekilde tasarlanmaması durumunda, öğrenme üzerinde olumsuz etkiler ortaya çıkabilir.

Çalışmanın kuramsal çerçevesini Bilişsel Yük Teorisi (Cognitive Load Theory - CLT) oluşturmaktadır. Sweller (1988) tarafından geliştirilen bu teori, insan bilişsel mimarisinin sınırlamalarını merkeze alarak öğretim tasarımının nasıl optimize edilmesi gerektiğine dair güçlü bir çerçeve sunmaktadır. Teoriye göre, öğrenme sürecinde karşılaşılan bilişsel yük üç ana türde sınıflandırılmaktadır: görevin doğal karmaşıklığı ve öğrencinin ön bilgisiyle ilişkili içsel yük (intrinsic load); zayıf öğretim tasarımından kaynaklanan dışsal yük (extraneous load); ve anlamlı şema oluşturma çabasıyla ilişkili ilgili yük (germane load). Bu teorik çerçeve, YZ araçlarının matematik öğrenimindeki etkilerini analiz etmek için güçlü bir lens sunmaktadır.

YZ araçlarının bilişsel yük mekanizmaları üç boyutta ele alınmaktadır. Birincisi, YZ ile dışsal yükün azaltılması: YZ araçları, gereksiz bilgi arayışını ortadan kaldırarak, görselleştirmelerle soyut kavramları somutlaştırarak ve çözüm adımlarını optimize ederek öğrencinin dikkatini dağıtan unsurları azaltabilir. Örneğin, bir YZ aracının karmaşık bir problemi alt problemlere ayırması, öğrencinin temel yapıyı anlamasına yardımcı olur. İkincisi, YZ ile içsel yükün optimize edilmesi: YZ sistemleri, öğrencinin ön bilgi düzeyini tespit ederek içeriğin zorluk seviyesini ve sunum hızını bireysel ihtiyaçlara göre uyarlayabilir, böylece öğrencinin

kapasitesinin üzerinde veya altında kalan görevlerin oluşturacağı bilişsel yükü dengeleyebilir. Üçüncüsü, YZ ve ilgili yükün artırılması: YZ, öğrencinin derinlemesine düşünmesini, hipotezler test etmesini ve alternatif çözüm yolları keşfetmesini teşvik eden etkileşimler sunarak, anlamlı öğrenme için harcanan bilişsel çabayı üst düzey düşünme etkinliklerine yönlendirebilir. Bununla birlikte, YZ araçlarının tüm çözüm adımlarını doğrudan sunması veya aşırı bilgi yüklemesi yapması durumunda öğrenmeyi sekteye uğratabileceği de vurgulanmaktadır.

Sonuç olarak, bu teorik çerçeve, eğitim teknolojileri tasarımcılarına, öğretmenlere ve araştırmacılara YZ araçlarını öğrenme sürecine entegre ederken bilişsel faktörleri göz önünde bulundurmaları için yol gösterici olmayı hedeflemektedir. Çalışma, gelecekte yapılacak ampirik çalışmalar için test edilebilir hipotezlerin ve tasarım ilkelerinin geliştirilmesine zemin hazırlayacak ve YZ destekli öğrenme ortamlarının daha etkili ve verimli tasarlanmasına katkı sağlayacaktır.

Anahtar Kelimeler: Yapay zeka, Matematik, Bilişsel yük kuramı

ABSTRACT

This study aims to present a comprehensive theoretical framework for managing cognitive load in artificial intelligence (AI)-supported mathematics learning environments. The rapid proliferation of AI-based tools such as ChatGPT, Wolfram Alpha, and Photomath in educational settings has drawn significant attention to their potential to provide immediate feedback, offer step-by-step solutions, and simplify complex problems. However, mathematics education is inherently a cognitively demanding discipline that involves abstract concepts, multi-step procedures, and intensive problem-solving skills. The unconscious or pedagogically inappropriate use of AI tools can create significant obstacles in the learning process. In particular, if the information flow and interaction formats provided by these tools are not designed with due consideration for learners' limited working memory capacity, negative effects on learning outcomes may emerge.

The theoretical foundation of this study is built upon Cognitive Load Theory (CLT), originally developed by Sweller (1988). CLT provides a robust framework for optimizing instructional design by centering on the limitations of human cognitive architecture. According to the theory, cognitive load encountered during the learning process is classified into three main types: intrinsic load, which is related to the natural complexity of the task and the learner's prior knowledge; extraneous load, which stems from poor instructional design; and germane load, which is associated with the effort devoted to meaningful schema construction and deep learning.

AI tools' effects on cognitive load mechanisms are examined in three dimensions. First, AI can reduce extraneous cognitive load by eliminating unnecessary information search, concretizing abstract concepts through visualizations, and optimizing solution steps. For example, an AI tool

that breaks down a complex problem into sub-problems helps learners understand the core structure. Second, AI can optimize intrinsic cognitive load by detecting learners' prior knowledge levels and adapting the difficulty level and presentation speed of content according to individual needs, thereby balancing the cognitive load imposed by tasks that are either above or below the learner's capacity. Third, AI can enhance germane cognitive load by encouraging deeper thinking, hypothesis testing, and exploration of alternative solution paths, thereby directing cognitive effort toward higher-order thinking activities. Nevertheless, it is also emphasized that AI tools may hinder learning if they directly present all solution steps or overload learners with excessive information.

In conclusion, this theoretical framework aims to guide educational technology designers, teachers, and researchers in considering cognitive factors when integrating AI tools into mathematics learning processes. The study lays the groundwork for developing testable hypotheses and design principles for future empirical research, thereby contributing to the design of more effective and efficient AI-supported learning environments.

Keywords: Artificial intelligence, Mathematics education, Cognitive Load Theory

GİRİŞ

Kasım 2022 tarihinde OpenAI tarafından geliştirilerek piyasaya sürülen Chat Generative Pre-trained Transformer (Sohbet Amaçlı Üretken Ön Eğitimli Dönüştürücü), yaygın kullanımıyla ChatGPT, eğitim alanında ivmelenmeye ve ilgili akademik çalışmaların hızla yoğunlaşmasına önayak olmuştur (Liu vd., 2026; Wei & Wijaya, 2025). Bu süreçte her alanda olduğu gibi matematik eğitiminde de önemli adımlar atılmıştır. Özellikle, kişiselleştirilmiş öğrenme, kavram yanılgılarının tespiti, görselleştirme, bilişsel yük yönetimi ve problem çözme becerilerinin geliştirilmesi gibi konularda önemli başarılar elde edilmiştir (Eti vd., 2026; Gkintoni vd., 2025; Kaya & Yavuz, 2025; Wang & Li, 2026). Örneğin, öğrencinin başarılı oldukça problemleri zorlaştıran, zorlandığında ise ipuçları sunan veya görevi basitleştiren, içerik sırasını öğrencinin anlık performansına göre dinamik olarak ayarlayan yeni sistemler (örneğin ALEKS, Khan Academy, Cognitive Tutor Algebra I, MathGPT, Edmentum Exact Path) geliştirilmiştir (Alvarez, 2024; Eti vd., 2026; Halkiopoulos & Gkintoni, 2024; Ismail vd., 2026). Yine öğrencilerin en çok zorlandığı alanlardan biri olan metin tabanlı matematik problemlerinde ChatGPT-MPS ve ChatGPT-MWPS gibi sistemler, karmaşık metin yapılarını sadeleştirerek dilsel zorlukları matematiksel mantıktan ayırmıştır (Liu vd., 2025a, Liu vd., 2025b; Patac & Patac, 2025). Özellikle ortaöğretimde kullanılan YZ araçları, soyut kavramların görselleştirilmesini sağlayarak öğrencilerin Van Hiele geometrik düşünme seviyelerini hızlandırmakta (Polydoros vd, 2025); görsel sezgiden tümdengelimli akıl yürütmeye geçişi bilişsel bir iskele ile desteklemektedir. Bunun yanı sıra, bu araçlar ispat oluşturma sürecinde mantıksal adımların düzenlenmesine yardımcı olmakta ve öğrenci başarı puanlarında anlamlı artışlar sağlamaktadır (Davor, 2026).

Bununa birlikte araştırmalar, yapay zekâ araçlarının doğrudan öğretim amaçlı ve pedagojik bir çerçeveden yoksun olarak kullanıldığı durumlarda, öğrenmeyi teşvik etmek yerine yalnızca

cevap bulmaya odaklanan bir yapıya dönüşebileceğini göstermektedir (Lodge & Loble, 2026; Setala vd, 2025). Bu durumun ise, bilişsel tembelliğe ve eleştirel düşünme becerilerinin zayıflamasına neden olduğu yaygın olarak kabul edilmektedir (Almarashdi,vd, 2024; Fan vd, 2025; Kuo vd., 2026). Son araştırmalar, Bilişsel Yük Teorisi'nin (CLT) yapay zeka destekli öğrenme sürecinde, bilişsel kaynakların yönetiminde güçlü bir çerçeve sağladığını ortaya koymaktadır (Alam vd, 2026; Patac & Patac, 2025). Bilişsel Yük Teorisi'nin temel ilkeleri ve bu kuram doğrultusunda yapılandırılan yapay zeka destekli öğrenme ortamlarının temel özellikleri aşağıdaki bölümde ayrıntılı olarak ele alınmaktadır.

Bilişsel Yük Teorisi

John Sweller (1988) tarafından geliştirilen Bilişsel Yük Teorisi (CLT), öğretim tasarımı ile insan bilişsel mimarisi arasındaki ilişkiyi ele almakta ve temel olarak öğrenme sırasında zihnin harcadığı eforun nasıl yönetilmesi gerektiğini (Alam vd, 2026) açıklamaktadır. Teori, temel olarak insan zihnindeki çalışan bellek kapasitesinin sınırlı olduğu ve öğretim materyallerinin bu kapasiteyi aşacak şekilde sunulmasının öğrenmeyi engelleyeceği (bilişsel aşırı yüklenme) varsayımına dayanır.

Bu teoriye göre, insan bilişsel mimarisi çalışma belleği ve uzun süreli bellek şeklinde iki ana bileşen üzerinden ele alınır. Yeni bilgilerin aktif olarak işlendiği yer olan sınırlı çalışma belleği, adından da anlaşılacağı gibi sınırlı kapasiteye sahiptir. Aynı anda sadece 4 ila 7 bilgi öğesini tutan bu bellekte kapasite aşıldığında bilişsel aşırı yüklenme (cognitive overload) meydana gelir ve öğrenme süreci engellenir (Lodge & Loble, 2026; Patac & Patac, 2025). Bilgilerin kalıcı olarak depolandığı yer ise sınırsız uzun süreli bellektir. Burada yeni bilgiler şema adı verilen karmaşık bilgi yapılarına entegre edilir (Lodge & Loble, 2026; Sweller vd, 2019)

Teori, öğrenme sırasında ortaya çıkan bilişsel yükü üç ayrı ancak birbiriyle etkileşimli bileşene ayırmaktadır. İçsel yük, öğrenilecek içeriğin doğasından ve öğe etkileşimi düzeyinden kaynaklanır; dışsal yük, öğretim materyalinin zayıf tasarımından doğar. Etkili yük ise şema oluşumu ve otomasyon sürecine ayrılan, öğrenmeyi doğrudan destekleyen bilişsel çabadır (Sweller, 1988).

Son yıllarda yapılan çalışmalar, bilişsel yükün yönetiminde yapay zeka teknolojilerini önemli bir araç olarak görmektedir (Chiriatti vd, 2025; Lodge & Loble, 2026; Sharma & Khadka, 2026). Bu bağlamda yapay zeka teknolojileri, sadece komut verilen bir makine değil, kullanıcıyla sürekli bir geri bildirim döngüsü içinde etkileşime giren aktif bir iş birliği ortağı olarak görülmektedir (Riva, 2026). Riva'ya (2026) göre, yapay zeka, bilişsel yükü sadece pasif bir şekilde azaltan bir araç değil, içsel, dışsal ve etkili yükleri dinamik olarak dengeleyen bir "bilişsel eş-düzenleyici" (co-regulator) olarak konumlandırılmaktadır.

Yapay zeka araçlarının bilişsel yük mekanizmaları üç boyutta ele alınmaktadır. Birincisi, yapay zeka ile dışsal yükün azaltılması: YZ sistemleri, öğrenme hedefiyle doğrudan ilgisi olmayan hesaplama işlemleri, dil bilgisi kontrolü veya karmaşık arayüz görevleri gibi rutin işleri üstlenerek öğrencinin sınırlı çalışma belleği kaynaklarını asıl kavramsal işlemlere odaklamasını sağlar (Alam vd, 2026; Riva, 2026). İkincisi, YZ ile içsel yükün kalibrasyonu: Öğrenilen

materyalin kendi doğasından kaynaklanan bu zorluk, YZ tarafından görev ayrıştırma yöntemiyle yönetilir. Karmaşık matematiksel problemler küçük ve yönetilebilir adımlara bölünerek, öğrencinin mevcut bilgi seviyesine ve "yakınsal gelişim alanına" uygun hale getirilir (Riva, 2026). Üçüncüsü, YZ ile etkili yükün teşvik edilmesi: Öğrenmenin gerçekleşmesi ve bilginin uzun süreli bellekteki şemalara entegre edilmesi için gereken bu yapıcı zihinsel çaba, YZ'nin öğrenciye doğrudan cevap vermek yerine yansıtıcı sorular sorması ve bilişsel iskele (scaffolding) kurmasıyla desteklenir. Sokratik diyaloglar ve metabilişsel ipuçları sayesinde öğrenci, strateji seçimi ve çözüm yollarını gerekçelendirme gibi derin bilişsel süreçlere yönlendirilerek anlamlı öğrenme deneyimleri yaşar (Alam vd, 2026).

Şekil 1'de etkili bir bilişsel yük yönetimi için yapay zeka araçlarının nasıl yapılandırılbileceğini gösteren bir örnek model gösterilmiştir. Buna göre, ders sürecinin yedi aşaması (ön değerlendirme, kademeli görev sunumu, yapay zeka arayüzü, rehberli uygulama, bağımsız problem çözme, pekiştirme ve yansıtma) ile her aşamada yapay zekânın üstlendiği somut rol birlikte gösterilmektedir. Kutu renkleri ilgili aşamanın hangi bilişsel yük bileşenini (içsel, dışsal, etkili) hedeflediğini; mor yapay zekasatırı ise yapay zekânın o aşamadaki işlevini belirtmektedir. Model, Sweller (1988), Sweller ve Chandler (1991), Mayer (2005) ve Kirschner ve ark. (2006) tarafından ortaya konan bilişsel yük ilkeleri ile Deci ve Ryan'ın (2000) öz-belirleme kuramı temel alınarak araştırmacı tarafından geliştirilmiştir.

SONUÇ

Bu çalışma, yapay zekâ destekli matematik öğretiminin bilişsel yük teorisi merceğinden sistematik olarak ele alınması gerektiğini savunmuştur. Son yıllarda alan yazında gözlenen kavramsal dönüşüm, yapay zekâyı yalnızca komut uygulayan pasif bir araç olmaktan çıkarıp, öğrenme sürecine sürekli bir geri bildirim döngüsü içinde eşlik eden aktif bir iş birliği ortağı olarak konumlandırmaktadır (Chiriatti vd., 2025; Lodge & Loble, 2026; Sharma & Khadka, 2026). Bu çalışmada önerilen ve Şekil 1'de sunulan yedi aşamalı model, bu eş-düzenleme işlevinin somut bir öğretim sürecine nasıl aktarılabileceğini göstermektedir: ön değerlendirmeden yansıtmaya uzanan her aşamada yapay zekânın rolü, ilgili bilişsel yük bileşenini yönetmeye yönelik açık bir işlevle tanımlanmıştır.



Şekil 1. Bilişsel Yük Teorisi Temelinde Yapay Zekâ Destekli Matematik Dersinin Yapılandırılması

Bu bütünleşik çerçeve, Sweller (1988), Sweller ve Chandler (1991), Mayer (2005) ve Kirschner ve arkadaşlarının (2006) ortaya koyduğu klasik bilişsel yük ilkelerini, Deci ve Ryan'ın (2000) öz-belirleme kuramıyla ve güncel eş-düzenleyici yapay zekâ anlayışıyla bir araya getirerek matematik öğretiminde yapay zeka entegrasyonuna hem bilişsel hem motivasyonel açıdan bakan bütüncül bir model sunmaktadır. Ancak bu potansiyelin kendiliğinden gerçekleşmediği vurgulanmalıdır; yapay zekânın eş-düzenleyici işlevi, ancak dikkatli pedagojik tasarım, sürekli izleme ve öğretmenin süreçteki etkin rolüyle anlamlı hâle gelmektedir. Gelecekteki ampirik çalışmaların, burada önerilen yedi aşamalı modeli farklı matematik konuları ve öğrenci grupları üzerinde sınayarak, yapay zekânın bilişsel eş-düzenleyici olarak işlevselliğini deneysel olarak doğrulaması, alan yazına önemli bir katkı sağlayacaktır.

KAYNAKÇA

Alam, M. N., Islam, M. A., Babiker, M. O. A., Siddiqui, M. S., Amin, M. B., & Oláh, J. (2026). AI-assisted learning tools and student learning outcomes: A cognitive load theory perspective. *Computers in Human Behavior Reports, 21*, 100986. https://doi.org/10.1016/j.chbr.2026.100986

Almarashdi, H. S., Jarrah, A. M., Khurma, O. A., & Gningue, S. M. (2024). Unveiling the potential: A systematic review of ChatGPT in transforming mathematics teaching and learning. *Eurasia Journal of Mathematics, Science and Technology Education, 20*(12), Article em2555. https://doi.org/10.29333/ejmste/15739

Alvarez J. (2024). Evaluating the impact of AI–powered tutors Mathgpt and flexi 2.0 in enhancing Calculus learning. *Jur. Ilmh. Ilm. Ter. Un. Ja. 8*, 495– 508. https://doi.org/10.22437/jiituj.v8i2.34809

Chiriatti, M., Bergamaschi Ganapini, M., Panai, E., Wiederhold, B. K., & Riva, G. (2025). System 0: Transforming artificial intelligence into a cognitive extension. *Cyberpsychology, Behavior, and Social Networking*, 28(7), 534–542.

Davor, I. (2026). Generative AI-supported instruction as a cognitive scaffold: effects on senior high school students' geometric reasoning and proof construction. *Frontiers in Education, 11,*. https://doi.org/10.3389/feduc.2026.1869961

Eti, N., Mosia, M., & Egara, F. O. (2026). The role of AI-driven personalised learning in enhancing mathematics problem-solving skills: a systematic review. *Frontiers in Computer Science, 8*.

Fan, Y., Tang, L., Le, H., Shen, K., Tan, S., Zhao, Y., Shen, Y., Li, X., & Gašević, D. (2025). Beware of metacognitive laziness: Effects of generative artificial intelligence on learning motivation, processes, and performance. *British Journal of Educational Technology, 56*(2), 489–530.

Gkintoni, E., Antonopoulou, H., Sortwell, A., & Halkiopoulos, C. (2025). Challenging Cognitive Load Theory: The Role of Educational Neuroscience and Artificial Intelligence in Redefining Learning Efficacy. *Brain Sciences, 15*(2), 203.

Halkiopoulos, C., & Gkintoni, E. (2024). Leveraging AI in E-Learning: Personalized Learning and Adaptive Assessment through Cognitive Neuropsychology—A Systematic Analysis. *Electronics, 13*(18), 3762. https://doi.org/10.3390/electronics13183762

Ismail, S. M., Abdel-Al Ibrahim, K. A., & Hashemifardnia, A. (2026). The impact of artificial intelligence-based tutoring systems on student engagement and long-term retention in math education. *Scientific Reports*. https://doi.org/10.1038/s41598-026-62920-6

Kaya, D., & Yavuz, S. (2025). Can Generative AI and ChatGPT Break Human Supremacy in Mathematics and Reshape Competence in Cognitive-Demanding Problem-Solving Tasks? *Journal of Intelligence, 13*, 43.

Kuo, B.-C., Bai, Z.-E., & Lin, C.-H. (2026). Developing an AI learning companion for mathematics problem solving in elementary schools. *Computers & Education, 240,* 105463. https://doi.org/10.1016/j.compedu.2025.105463

Liu, J., Keane, T., Sun, D., Yang, Y., & Yang, Y. (2025b). Design and evaluation of ChatGPT-MWPS: an AI-enhanced learning system for improving primary students' mathematical word problem solving. *Smart Learning Environments, 12*(68).

Liu, J., Sun, D., Sun, J., Wang, J., & Yu, P. L. H. (2025a). Designing a generative AI enabled learning environment for mathematics word problem solving in primary schools: Learning performance, attitudes and interaction. *Computers and Education: Artificial Intelligence, 9*, 100438.

Liu, B., Zhang, W., & Wang, F. (2026). Can generative artificial intelligence effectively enhance students' mathematics learning outcomes?—A meta-analysis of empirical studies from 2023 to 2025. *Education Sciences, 16*(1), 140. https://doi.org/10.3390/educsci16010140

Lodge, J. M., & Loble, L. (2026). Artificial intelligence, cognitive offloading and implications for education. *University of Technology Sydney*. https://doi.org/10.71741/4pyxmbnjaq.31302475

Patac, L. P., & Patac, A. V., Jr. (2025). Using ChatGPT for academic support: Managing cognitive load and enhancing learning efficiency – A phenomenological approach. Social Sciences & Humanities Open, 11, 101301.

Polydoros, G., Galitskaya, V., Antoniou A.-S., & Drigas, A. (2025). AI technology integration in elementary geometry and its effects on performance, anxiety levels, learning styles, cognitive styles, and executive functions. *Scientific Electronic Archives, 18*(2). https://doi.org/10.36560/18220252050

Setälä, M., Heilala, V., Sikström, P., & Kärkkäinen, T. (2025). The use of generative artificial intelligence for upper secondary mathematics education through the lens of technology acceptance. In Proceedings of the 40th ACM/SIGAPP Symposium on Applied Computing (SAC '25) (pp. 74–82). *Association for Computing Machinery*. https://doi.org/10.1145/3672608.3707817

Sharma, R., & Khadka, S. (2026). Managing learning demands through mobile AI: A cognitive load perspective on ChatGPT use in Nepal. *Mobile Media & Communication*.

Sweller, J. (1988). Cognitive load during problem solving: Effects on learning. *Cognitive Science, 12*(2), 257–285. https://doi.org/10.1207/s15516709cog1202_4

Sweller, J., & Chandler, P. (1991). Evidence for Cognitive Load Theory. *Cognition and Instruction, 8*(4), 351–362. http://www.jstor.org/stable/3233599

Wang, J., & Li, Y. (2026). AI and the future of educational technology: a systematic review of pedagogical innovations and challenges. *Cogent Education, 13*(1), 2691562.

Wei, Z., & Wijaya, T. T. (2025). Awareness, acceptance, and adoption of GenAI by K12 mathematics teachers an empirical study integrating TAM and TPB. *BMC Psychology, 13*(478).

MAPPING RECENT TRENDS IN ARTIFICIAL INTELLIGENCE RESEARCH FOR SUSTAINABLE AGRICULTURE: A BIBLIOMETRIC AND SYSTEMATIC REVIEW

M.Sc., Esra YILMAZ ÖNDER
Tokat Gaziosmanpaşa University, Faculty of Agriculture, Department of Agricultural Economics, Tokat, Türkiye
ORCID ID : https://orcid.org/ 0009-0002-8146-4909

1. INTRODUCTION

Sustainable agriculture (SA) is currently regarded as one of the fundamental approaches that addresses food security and the conservation of natural resources together. It is also important for environmental protection and the maintenance of socioeconomic well-being in rural areas. The main objective of this approach is to ensure that natural resources such as soil and water are not excessively consumed while production continues and that future generations retain the opportunity to benefit from these resources (Blakeney, 2022; El Bilali et al., 2021).

Changes in climatic conditions, declining water resources, loss of soil fertility, reductions in biodiversity, and population growth are increasing the pressures faced by agricultural production. These problems make the economic, technological, and environmental risks confronting small-scale producers particularly evident in developing countries. Although agriculture is an important source of livelihood in Africa, and especially in Sub-Saharan Africa, climate risks and structural problems in agricultural production pose significant challenges to sustainable production (Ado et al., 2020).

Under these conditions, the integration of scientific advances and new technologies into agricultural production processes has become increasingly important. Artificial intelligence (AI) is one of the prominent technologies in this transformation and can be used to support data-driven decision-making, automate certain production operations, and plan resource use more precisely. Technologies such as machine learning, deep learning, computer vision, and the Internet of Things (IoT) are used in various areas, including crop and soil monitoring, disease and pest detection, weed management, irrigation, yield prediction, and agricultural automation (Jha et al., 2019; Qazi et al., 2022).

One of the prominent aspects of AI use in agriculture is its ability to evaluate data related to the production process and provide support for resource use and decision-making. For example, AI-supported irrigation systems use soil, environmental, and meteorological data to optimize irrigation decisions and reduce water use (Preite & Vignali, 2024). Similarly, studies have been conducted on identifying weeds from images using computer vision and deep learning techniques and intervening only in areas where intervention is necessary (Vasileiou et al., 2024).

Another area of AI application in agriculture is robotic systems. Robotic applications are becoming increasingly important, particularly for weed control, crop monitoring, and the automation of labor-intensive agricultural activities. Studies using YOLOv7-based deep learning and robotic systems for targeted weed identification represent a recent example of this development (Balabantaray et al., 2024).

However, evaluating the use of AI in agriculture solely on the basis of technical performance is not sufficient. Factors such as data access, digital infrastructure, technological skills, cost, algorithmic transparency, and adaptation to regional conditions can affect the sustainability and accessibility of AI applications. In particular, differences in technological and research capacity between developed and developing regions may constitute a significant obstacle to the equitable global diffusion of AI-based agricultural solutions.

The development of digital technologies in agriculture has accelerated with the spread of Agriculture 4.0 and precision agriculture approaches. Technologies such as AI, IoT, sensor technologies, robotics, and data analytics contribute to the more precise and data-driven management of agricultural production (Liu et al., 2021; Qazi et al., 2022).

Although numerous studies in the existing literature examine applications of AI in agriculture, a substantial proportion of these studies focus on specific technologies or application areas. This situation highlights the need for a holistic assessment of the research relationship between AI and sustainable agriculture.

Bibliometric analysis enables the systematic examination of publications and scientific relationships within a research field. Publication counts, citations, authors, countries, institutions, and keywords can be evaluated within this framework. Scientific mapping techniques help reveal the intellectual, social, and conceptual relationships within a research field.

Indeed, Pervez et al. (2026) examined 645 documents from the Scopus database covering the period 2010–2025 and evaluated research trends at the intersection of AI and sustainable agriculture using bibliometric and systematic methods together. The study reported that research output has increased in recent years and that areas such as machine learning, precision agriculture, deep learning, computer vision, robotics, and IoT have become prominent. The study also draws attention to the uneven distribution of research production across regions and the need for context-specific research in the Global South (Pervez et al., 2026).

This study aims to reveal the current structure and evolution over time of the research field at the intersection of sustainable agriculture and AI. For this purpose, bibliometric analysis and systematic literature review were used together. The study examines research trends between AI and sustainable agriculture during the 2010–2025 period in terms of scientific production, contributions by authors and countries, collaboration networks, keywords, thematic development, and research topics.

2. CONCEPTUAL FRAMEWORK

2.1. Sustainable Agriculture

Sustainable agriculture refers to the development of production systems that are economically viable, environmentally responsible, and socially inclusive. This approach encompasses not only increasing agricultural production but also the long-term conservation of soil, water, and other natural resources (Blakeney, 2022; El Bilali et al., 2021).

Increasing pressure from climate change and on natural resources makes more efficient resource use in agricultural production necessary. Therefore, sustainable agriculture is evaluated through

interrelated dimensions such as productivity, resource optimization, environmental protection, climate change adaptation, and food security.

2.2. Artificial Intelligence and Agriculture

Artificial intelligence encompasses technologies that enable computer systems to perform functions associated with human intelligence, such as learning, prediction, classification, and decision-making. In the agricultural sector, AI is used together with machine learning, deep learning, computer vision, and IoT technologies (Jha et al., 2019; Qazi et al., 2022).

The main application areas of AI in agriculture include:

- ürün verimi tahmini,
- hastalık ve zararlı tespiti,
- yabancı ot belirleme,
- hassas sulama,
- toprak analizi,
- gübreleme optimizasyonu,
- ürün ve toprak izleme,
- robotik tarım uygulamaları,
- tarımsal karar destek sistemleri yer almaktadır.

Jha et al. (2019) demonstrated that technologies such as AI, machine learning, deep learning, and IoT can be used in areas such as agricultural automation, irrigation, weed management, and crop monitoring.

2.3. Artificial Intelligence and Resource Optimization

One of the important contributions of AI to sustainable agriculture is the optimization of resource use. In particular, efficient use of water resources can be regarded as one of the fundamental elements of agricultural sustainability.

Preite and Vignali (2024) developed an AI- and machine-learning-based irrigation management system and investigated the use of soil, environmental, and weather forecast data in irrigation decisions. This approach demonstrates the potential of AI for optimizing irrigation amounts and preventing unnecessary water use.

2.4. Artificial Intelligence and Computer Vision

Computer vision and deep learning are used particularly for the automatic identification of plant diseases, weeds, and other agricultural elements from images.

Vasileiou et al. (2024) systematically reviewed the use of AI in weed management and emphasized the importance of deep learning and image-processing technologies for precision weed management. The study also highlights the potential of these technologies for environmental and economic sustainability.

2.5. Artificial Intelligence and Robotics

Robotic systems constitute an important emerging research area in terms of automating agricultural activities. Robotic systems are used particularly for weed control, crop monitoring, and labor-intensive activities.

Balabantaray et al. (2024) investigated the use of a YOLOv7-based deep learning method combined with robotic systems for the targeted management of Palmer amaranth. Such studies represent current examples of the integration of AI with agricultural automation.

2.6. Explainable Artificial Intelligence

The increasing use of AI systems in agricultural decision-support processes makes it important for model results to be understandable and trustworthy to users. In this context, Explainable Artificial Intelligence (XAI) encompasses methods aimed at understanding why and how the results produced by complex models arise.

Transparency, data management, algorithmic fairness, and explainability are becoming increasingly important for the future development of AI in sustainable agriculture. Pervez et al. (2026) also draw attention to the need to develop explainable AI frameworks and transparency in AI research for sustainable agriculture.

3. METHOD

3.1. Research Design

Two approaches were considered together within the methodological framework of the study: bibliometric analysis and systematic literature review. In the bibliometric section, indicators related to publication output and scientific networks were examined; in the systematic review section, the contents of the selected studies and prominent research directions were addressed.

3.2. Data Source

The bibliographic data of the study were obtained from the Scopus database. Scopus is one of the databases widely used in bibliometric research because it covers a large number of scientific publications from different disciplines.

In constructing the dataset, studies published between 2015 and 2025 that addressed the relationship between AI and sustainable agriculture were evaluated.

3.3. Database Search Strategy

In line with the purpose of the study, keywords containing the concepts of AI and sustainable agriculture together were used.

Proposed Scopus search query:

TITLE-ABS-KEY
("artificial intelligence" OR "machine learning" OR "deep learning" OR "AI")
AND ("sustainable agriculture" OR "sustainable farming" OR "smart agriculture" OR "precision agriculture")

The purpose of this search strategy is to identify publications examining the scientific relationship between AI technologies and sustainable agriculture.

3.4. Inclusion Criteria

The following studies were included:

- yapay zekâ, makine öğrenmesi veya derin öğrenme teknolojilerini ele alan,
- sürdürülebilir tarım, akıllı tarım, hassas tarım veya tarımsal sürdürülebilirlikle ilişkili olan,
- Scopus veri tabanında bibliyografik bilgileri bulunan,
- belirlenen tarih aralığında yayımlanan,
- İngilizce dilinde yayımlanan,
- araştırma makalesi veya inceleme makalesi niteliğindeki

were included in the study.

3.5. Exclusion Criteria

The following were excluded from the study:

- araştırma konusu ile doğrudan ilişkili olmayan,
- yinelenen kayıtlar,
- bibliyografik bilgileri eksik olan,
- konu dışında kalan konferans özeti, editöre mektup vb. yayınlar,
- yapay zekâ veya sürdürülebilir tarım boyutlarından anlamlı biçimde hiçbirini içermeyen

were excluded from the study.

3.6. Data Analysis

Bibliometrix/Biblioshiny and VOSviewer software were used to conduct the bibliometric analyses.

Using Biblioshiny:

- yıllık bilimsel üretim,
- yazar performansı,
- ülke katkıları,
- kaynak performansı,
- atıf göstergeleri,
- tematik gelişim

were examined.

Using VOSviewer:

- ortak yazarlık ağları,
- ülke iş birliği ağları,
- ortak atıf ağları,

- bibliyografik eşleşme,
- anahtar kelime birlikte oluşum ağları görselleştirilmiştir.

This approach enables the research field to be evaluated not only in terms of publication counts but also in terms of its intellectual, social, and conceptual structure.

4. BIBLIOMETRIC ANALYSIS PROCESS

1. Data Collection

↓

2. Data Cleaning and Removal of Duplicate Records

↓

3. Application of Inclusion and Exclusion Criteria

↓

4. Export of Bibliographic Data

↓

5. Conducting Bibliometric Analyses

↓

6. Visualization of Scientific Networks and Themes

↓

7. Systematic Literature Review

↓

8. Holistic Evaluation of the Findings

Figure 1. Bibliometric and systematic review process of the study

5. CONCLUSION AND DISCUSSION

The results of the bibliometric and systematic review indicate that research at the intersection of AI and sustainable agriculture has developed markedly, particularly in recent years. In the recent bibliometric review conducted by Pervez et al. (2026), 645 studies published in Scopus during 2010–2025 were examined, and a marked increase in publication output, particularly after 2020, was identified. The study reported an annual publication growth rate of 12.69%.

Machine learning, deep learning, precision agriculture, computer vision, robotics, and IoT are among the prominent technologies in the research field. This indicates that sustainable agriculture research is directed not only toward increasing production quantities but also toward optimizing resource use, improving agricultural decision-making processes, and automation.

The efficient use of water resources is particularly important as a research area for sustainable agriculture. AI-supported irrigation systems can use soil and meteorological data to optimize irrigation decisions (Preite & Vignali, 2024).

Similarly, the use of AI and computer vision technologies for the automatic identification of weeds and targeted intervention can be considered among the applications that may contribute to more precise use of agricultural inputs (Vasileiou et al., 2024).

However, successful results from AI technologies under controlled experimental conditions do not necessarily mean that these systems can be directly applied in different agricultural production environments. Future research should jointly evaluate long-term field studies, different agroecological conditions, and the economic feasibility of these technologies.

Another important finding of the study is that research on AI and sustainable agriculture has not developed equally across different geographical regions. Pervez et al. (2026) reported that India, China, and the United States stand out in research output, whereas Africa, South America, and the Middle East have lower levels of representation. This situation highlights the need to adapt AI solutions to different agroecological and socioeconomic conditions.

Therefore, future research should focus not only on technological performance but also on the accessibility of technologies for different income groups, small-scale farmers, and different regional production systems.

Furthermore, the technical, agronomic, and socioeconomic dimensions of AI research need to be integrated to a greater extent. Interdisciplinary collaboration among AI specialists, agricultural engineers, agricultural economists, and social scientists can contribute to the development of solutions suited to regional needs.

Explainability and transparency are also important research areas for the future of AI in sustainable agriculture. In particular, ensuring that AI models used in agricultural decision-support systems are understandable to producers is important for technology adoption and building trust. Pervez et al. (2026) also emphasize the need for explainable AI, ethical use, and stronger international collaboration in future research.

The overall picture of the study shows that AI research in sustainable agriculture focuses on various areas such as productivity, resource use, environmental aspects, climate change adaptation, automation, and decision support. Nevertheless, for technological development to translate into sustainable outcomes, regional conditions, economic feasibility, data access, digital infrastructure, and user needs must be considered together.

6. REFERENCES

Ado, A. M., Savadogo, P., Pervez, A. K. M. K., & Mudimu, G. T. (2020). Farmers' perceptions and adaptation strategies to climate risks and their determinants: Insights from a farming community of Aguie district in Niger. *GeoJournal, 85*(4), 1075–1095.

Balabantaray, A., Behera, S., Liew, C., Chamara, N., Singh, M., Jhala, A. J., & Pitla, S. (2024). Targeted weed management of Palmer amaranth using robotics and deep learning (YOLOv7). *Frontiers in Robotics and AI, 11*, 1441371. https://doi.org/10.3389/frobt.2024.1441371

Blakeney, M. (2022). Agricultural innovation and sustainable development. *Sustainability, 14*(5), 2698.

El Bilali, H., Strassner, C., & Ben Hassen, T. (2021). Sustainable agri-food systems: Environment, economy, society, and policy. *Sustainability, 13*(11), 6260.

Jha, K., Doshi, A., Patel, P., & Shah, M. (2019). A comprehensive review on automation in agriculture using artificial intelligence. Artificial Intelligence in Agriculture, 2, 1–12. https://doi.org/10.1016/j.aiia.2019.05.004https://doi.org/10.1016/j.aiia.2019.05.004

Liu, Y., Ma, X., Shu, L., Hancke, G. P., & Abu-Mahfouz, A. M. (2021). From Industry 4.0 to Agriculture 4.0: Current status, enabling technologies, and research challenges. *IEEE Transactions on Industrial Informatics, 17*(6), 4322–4334.

Pervez, A. K. M. K., Kabir, M. S., Prodhan, F. A., Rahman, M. H., Roy, A., & Mahedi, M. (2026). Mapping recent trends in artificial intelligence research for sustainable agriculture: A bibliometric and systematic review. Discover Artificial Intelligence, 6, 509. https://doi.org/10.1007/s44163-026-01024-6

Preite, L., & Vignali, G. (2024). Artificial intelligence to optimize water consumption in agriculture: A predictive algorithm-based irrigation management system. Computers and Electronics in Agriculture, 223, 109126. https://doi.org/10.1016/j.compag.2024.109126https://doi.org/10.1016/j.compag.2024.109126

Qazi, S., Khawaja, B. A., & Farooq, Q. U. (2022). IoT-equipped and AI-enabled next generation smart agriculture: A critical review, current challenges and future trends. *IEEE Access, 10*, 21219–21235.

Sarkar, A., Wang, H., Rahman, A., Memon, W. H., & Qian, L. (2022). A bibliometric analysis of sustainable agriculture: Based on the Web of Science (WOS) platform. *Environmental Science and Pollution Research, 29*, 38928–38949.

Vasileiou, M., Kyrgiakos, L. S., Kleisiari, C., Kleftodimos, G., Vlontzos, G., Belhouchette, H., & Pardalos, P. M. (2024). Transforming weed management in sustainable agriculture with artificial intelligence: A systematic literature review towards weed identification and deep learning. Crop Protection, 176, 106522. https://doi.org/10.1016/j.cropro.2023.106522https://doi.org/10.1016/j.cropro.2023.106522

SORUMLU YAPAY ZEKÂ DESTEKLİ TÜKETİCİ ETKİLEŞİMİNİ ŞEKİLLENDİREN DÖNÜŞÜM FAKTÖRLERİ: NİTEL BİR ARAŞTIRMA

TRANSFORMATION FACTORS FOR RESPONSIBLE AI-DRIVEN CONSUMER ENGAGEMENT: A QUALITATIVE EXPLORATION

Assistant Prof. Dr. Serap SARP[1]
[1]Faculty of Managerial Sciences, Abdullah Gül University, Türkiye
ORCID ID: 0000-0002-2560-4105

ÖZET

Yapay zekâ (YZ), işletmelerin tüketicilerle iletişim kurma ve müşteri ilişkilerini yönetme biçimini hızla dönüştürmektedir. Üretken yapay zekâ, öneri sistemleri ve kestirimsel analitik gibi teknolojiler artık yalnızca deneysel araçlar olmaktan çıkmış, modern pazarlama stratejilerinin ayrılmaz bir parçası hâline gelmiştir. Bu gelişmeler, işletmelerin kişiselleştirilmiş deneyimler sunmasına, tüketici ihtiyaçlarına gerçek zamanlı yanıt vermesine ve daha etkileşimli iletişim süreçleri oluşturmasına olanak sağlamaktadır. Bununla birlikte, bu teknolojiler şeffaflık, etik, tüketici güveni ve sorumlu karar alma süreçlerine ilişkin yeni sorunları da beraberinde getirmektedir. Dolayısıyla, günümüzde işletmelerden yalnızca yapay zekâ teknolojilerini benimsemeleri değil, aynı zamanda bu teknolojileri tüketicilerle uzun vadeli ilişkileri güçlendirecek biçimde kullanmaları da beklenmektedir.

Pazarlamada yapay zekâ alanındaki çalışmalar son yıllarda önemli ölçüde artmış olmasına rağmen, mevcut araştırmaların büyük bölümü yapay zekâ kullanımının sonuçlarına odaklanmaktadır. Önceki çalışmalar ağırlıklı olarak müşteri memnuniyeti, satın alma niyeti, sadakat, algılanan değer ve güven gibi çıktıları incelemiştir. Buna karşın, işletmelerin sorumlu yapay zekâ destekli tüketici etkileşimini oluşturmasını mümkün kılan koşullar hakkında bilgi sınırlıdır. Özellikle başarılı uygulamayı destekleyen dönüşüm faktörleri henüz yeterince araştırılmamıştır. Bu boşluk, işletmelerin tüketicilerle anlamlı etkileşimler kurarken teknolojik yenilik ile etik sorumluluk arasında nasıl denge kurabileceklerine ilişkin anlayışımızı sınırlandırmaktadır.

Bu çalışma, sorumlu yapay zekâ destekli tüketici etkileşimini şekillendiren faktörleri ortaya koyarak söz konusu boşluğu gidermeyi amaçlamaktadır. Araştırmada, yapay zekâ ve dijital pazarlama teknolojileriyle aktif olarak çalışan profesyonellerin deneyimlerinden zengin ve bağlama özgü bilgiler elde etmek amacıyla nitel araştırma deseni benimsenmiştir. Bu kapsamda pazarlama yöneticileri, dijital pazarlama uzmanları, müşteri deneyimi yöneticileri, yapay zekâ danışmanları ve uzmanlık alanı yapay zekâ ile pazarlamanın kesişiminde bulunan akademisyenlerle yarı yapılandırılmış görüşmeler gerçekleştirilecektir. Elde edilen veriler, Braun ve Clarke'ın tematik analiz yaklaşımı kullanılarak analiz edilecektir. Bu yöntem, katılımcı deneyimlerindeki örüntülerin sistematik ve esnek bir biçimde belirlenmesine ve üst düzey kavramsal temaların geliştirilmesine olanak sağlamaktadır.

Araştırma bulgularının, birbirinden bağımsız belirleyicilerden ziyade birbiriyle ilişkili dönüşüm faktörlerini ortaya koyması beklenmektedir. Örgütsel hazırlık, teknolojik yetkinlik, veri kalitesi, etik yönetişim, şeffaflık, tüketici güveni, kişiselleştirme kapasitesi, insan–yapay zekâ iş birliği ve sürekli örgütsel öğrenmenin temel temalar olarak öne çıkacağı öngörülmektedir. Bu faktörlerin birlikte, işletmelerin tüketiciler tarafından değerli, güvenilir ve sorumlu olarak algılanan yapay zekâ destekli etkileşimleri nasıl oluşturduklarını açıklaması beklenmektedir.

Bu çalışma, yapay zekâ ve pazarlama literatürüne çeşitli açılardan katkı sağlamayı hedeflemektedir. İlk olarak, yapay zekâ kullanımının sonuçlarından ziyade, sorumlu yapay zekâ

destekli tüketici etkileşimini mümkün kılan örgütsel ve stratejik koşullara odaklanmaktadır. İkinci olarak, yalnızca mevcut kuramlara dayanmayan, uygulayıcıların deneyimlerinden hareketle geliştirilen kavramsal bir çerçeve sunmaktadır. Son olarak, bulguların etik, şeffaf ve tüketici odaklı yapay zekâ stratejileri geliştirmek isteyen yöneticilere uygulamaya yönelik öneriler sağlaması beklenmektedir. Dijital dönüşüm, sorumlu yapay zekâ ve tüketici etkileşimi perspektiflerini bütünleştiren önerilen çerçeve, işletmelerin yapay zekânın giderek daha fazla benimsendiği pazarlarda tüketicilerle sürdürülebilir ilişkiler kurabilmelerine yönelik daha kapsamlı bir bakış açısı sunmaktadır.

Anahtar Kelimeler: Yapay Zekâ, Sorumlu Yapay Zekâ, Tüketici Etkileşimi, Dijital Dönüşüm, Tematik Analiz.

ABSTRACT

Artificial intelligence (AI) is rapidly reshaping the way organizations communicate with consumers and manage customer relationships. Technologies such as generative AI, recommendation systems, and predictive analytics are no longer experimental tools. They have become integral components of modern marketing strategies. These developments allow firms to deliver personalized experiences, respond to customer needs in real time, and create more interactive forms of engagement. At the same time, they introduce new challenges related to transparency, ethics, consumer trust, and responsible decision-making. Consequently, organizations are expected not only to adopt AI technologies but also to use them in ways that strengthen long-term relationships with consumers.

Although research on AI in marketing has expanded considerably, most existing studies concentrate on the consequences of AI adoption. Previous work has primarily examined outcomes such as customer satisfaction, purchase intention, loyalty, perceived value, and trust. Much less is known about the conditions that enable organizations to build responsible AI-driven consumer engagement in the first place. In particular, the transformation factors that support successful implementation remain underexplored. This gap limits our understanding of how organizations can balance technological innovation with ethical responsibility while creating meaningful interactions with consumers.

This study aims to address this limitation by exploring the factors that shape responsible AI-driven consumer engagement. A qualitative research design is adopted to reach rich and context-dependent insights from professionals who actively work with AI and digital marketing technologies. Semi-structured interviews will be conducted with marketing managers, digital marketing specialists, customer experience managers, AI consultants, and academics whose expertise lies at the intersection of artificial intelligence and marketing. The interview data will be analysed using Braun and Clarke's thematic analysis approach. This method provides a systematic yet flexible process for identifying patterns across participants' experiences and developing higher-level conceptual themes.

The findings are expected to reveal a set of interconnected transformation factors rather than isolated drivers. Organizational readiness, technological capability, data quality, ethical governance, transparency, consumer trust, personalization capability, human–AI collaboration, and continuous organizational learning are anticipated to emerge as central themes. Together, these factors are expected to explain how organizations create AI-enabled engagement that consumers perceive as valuable, trustworthy, and responsible.

This study contributes to the growing literature on AI and marketing in several ways. First, it shifts attention from the outcomes of AI adoption to the organizational and strategic conditions that make responsible AI-driven consumer engagement possible. Second, it develops a

conceptual framework grounded in practitioners' experiences instead of relying solely on existing theories. Finally, the findings are expected to offer practical guidance for managers seeking to design ethical, transparent, and consumer-centred AI strategies. By integrating perspectives from digital transformation, responsible AI, and consumer engagement, the proposed framework provides a more comprehensive understanding of how organizations can build sustainable relationships with consumers in an increasingly AI-powered marketplace.

Keywords: Artificial Intelligence, Responsible AI, Consumer Engagement, Digital Transformation, Thematic Analysis.

2018 SOSYAL BİLGİLER PROGRAMINDAKİ KAZANIMLARDAN 2024 PROGRAMINDAKİ ÖĞRENME ÇIKTILARINA YAPISAL BİR KARŞILAŞTIRMA

A STRUCTURAL COMPARISON OF THE LEARNING OBJECTIVES IN THE 2018 SOCIAL STUDIES CURRICULUM AND THE LEARNING OUTCOMES IN THE 2024 CURRICULUM

Ramazan AKYOL
Adıyaman Üniversitesi, Lisansüstü Eğitim Enstitüsü
Eğitim Programları ve Öğretim Anabilim Dalı
https://orcid.org/0009-0000-5197-3265

ÖZET

Bu araştırmanın amacı, 2018 Sosyal Bilgiler Dersi Öğretim Programında yer alan kazanımlar ile Türkiye Yüzyılı Maarif Modeli kapsamında hazırlanan 2024 Sosyal Bilgiler Dersi Öğretim Programında yer alan öğrenme çıktılarını yapısal açıdan karşılaştırarak programlar arasındaki değişimi ortaya koymaktır. Araştırma nitel araştırma yaklaşımı doğrultusunda doküman incelemesi yöntemiyle gerçekleştirilmiştir. Araştırmanın veri kaynaklarını Millî Eğitim Bakanlığı tarafından yayımlanan 2018 ve 2024 Sosyal Bilgiler Dersi Öğretim Programları oluşturmaktadır. Verilerin analizinde betimsel analiz kullanılmış; programlarda yer alan kazanım ve öğrenme çıktıları sayı, yapı, sınıf düzeyleri ve öğrenme alanları bağlamında karşılaştırmalı olarak incelenmiştir. Araştırma sonucunda 2018 programında toplam 131 kazanım bulunurken 2024 programında 72 öğrenme çıktısına yer verildiği belirlenmiştir. Bu durum, 2024 programında öğrencilerden beklenen öğrenmelerin sayısal olarak azaltıldığını göstermekle birlikte hedef yapısının daha farklı bir anlayışla düzenlendiğini ortaya koymaktadır. 2018 programında kazanımlar, öğrencilerin ulaşması beklenen bilgi, beceri ve tutumları ifade eden temel hedef birimleri olarak yapılandırılırken 2024 programında öğrenme çıktılarının süreç bileşenleriyle birlikte ele alındığı görülmüştür. Böylece öğrencinin yalnızca ulaşması beklenen öğrenme sonucu değil, bu sonuca ulaşırken işe koşacağı süreçler de programda görünür hâle getirilmiştir. Sonuç olarak 2018'den 2024'e geçişte Sosyal Bilgiler Dersi Öğretim Programının hedef yapısında yalnızca terminolojik bir değişimin gerçekleşmediği; kazanım temelli yapıdan öğrenme çıktıları ve süreç bileşenlerinin birlikte ele alındığı daha süreç odaklı bir yapılanmaya geçildiği sonucuna ulaşılmıştır.

Anahtar Kelimeler: Sosyal bilgiler, öğretim programı, Türkiye Yüzyılı Maarif Modeli, kazanım, öğrenme çıktısı.

ABSTRACT

The aim of this study is to reveal the changes between the curricula by structurally comparing the learning objectives included in the 2018 Social Studies Curriculum with the learning outcomes included in the 2024 Social Studies Curriculum developed within the framework of the Türkiye Century Education Model. The study was conducted using document analysis within a qualitative research approach. The data sources consisted of the 2018 and 2024 Social Studies Curricula published by the Ministry of National Education. Descriptive analysis was employed to analyze the data, and the learning objectives and learning outcomes in the curricula were comparatively examined in terms of their number, structure, grade levels, and learning areas. The findings revealed that the 2018 curriculum included a total of 131 learning objectives, whereas the 2024 curriculum included 72 learning outcomes. Although this indicates a numerical reduction in the learning expectations for students in the 2024 curriculum, it also demonstrates that the structure of the objectives was reorganized based on a different approach. While the learning objectives in the 2018 curriculum were structured as basic units expressing the knowledge, skills, and attitudes

students were expected to acquire, the learning outcomes in the 2024 curriculum were addressed together with process components. Thus, the curriculum makes visible not only the learning outcomes students are expected to achieve but also the processes they are expected to employ in achieving these outcomes. In conclusion, the transition from the 2018 to the 2024 Social Studies Curriculum represents more than a terminological change in the structure of curriculum objectives; it reflects a shift from an objective-based structure toward a more process-oriented structure in which learning outcomes and process components are addressed together.

Keywords: Social Studies, curriculum, Türkiye Century Education Model, learning objective, learning outcome.

MULTI-AGENT ARTIFICIAL INTELLIGENCE SYSTEMS FOR COLLABORATIVE BUSINESS DECISION INTELLIGENCE

Muhammad FAISAL
Joint Director (AICU) at Ministry of IT & Telecommunication, Pakistan
ORCID: 0000-0002-5797-766X

ABSTRACT

Modern enterprises increasingly require intelligent systems capable of solving complex problems through distributed collaboration rather than isolated computation. Multi-Agent Artificial Intelligence (MAAI) has emerged as a promising solution where multiple autonomous AI agents cooperate, negotiate, and share knowledge to achieve organizational objectives. This study examines the architecture, operational mechanisms, and business applications of multi-agent systems in enterprise decision intelligence. The proposed framework integrates specialized agents responsible for financial forecasting, customer analytics, operational planning, cybersecurity monitoring, and strategic recommendation generation. Communication protocols supported by Large Language Models facilitate cooperative reasoning while maintaining contextual awareness across multiple business domains. Experimental analysis demonstrates that collaborative AI architectures outperform single-agent systems in handling complex decision-making scenarios involving uncertainty and dynamic environments. Additionally, the research discusses scalability, coordination strategies, conflict resolution mechanisms, and governance models required for enterprise deployment. The findings indicate that multi-agent collaboration enhances organizational agility, reduces decision latency, improves analytical accuracy, and enables continuous optimization of enterprise resources. As businesses increasingly adopt intelligent automation, multi-agent systems represent a significant advancement toward decentralized AI ecosystems capable of supporting strategic business growth and digital innovation.

Keywords: Multi-Agent Systems, Decision Intelligence, Autonomous Agents, Enterprise AI, Collaborative Intelligence, Business Analytics, Artificial Intelligence

НАШ ОПЫТ РЕИНСЕРЦИИ ПЕРЕДНЕЙ КРЕСТООБРАЗНОЙ СВЯЗКИ КОЛЕННОГО СУСТАВА»

"OUR EXPERIENCE WITH REINSERTATION OF THE ANTERIOR CRUCIATE LIGAMENT OF THE KNEE JOINT"

Anorboev D.G.,
Republican Scientific and Practical Center for the Study of Traumatology and Otropedia, Tashkent
ORCID ID: 0009-0008-7473-1515

Irismetov M.E., Razhabov K.N.
Republican Scientific and Practical Center for the Study of Traumatology and Otropedia, Tashkent

ABSTRAKT. Anterior cruciate ligament (ACL) injuries are one of the most common knee injuries, especially among physically active individuals and athletes. Traditionally, the primary surgical treatment method has been ACL reconstruction using autografts. However, in recent years, increasing attention has been paid to organ-preserving techniques, particularly ACL reinsertion. This method involves anatomical restoration of the ligament by fixing it to the site of the tear while preserving the ligament's own tissue. The aim of this study is to analyze current data on the indications, technique, and clinical outcomes of ACL reinsertion.

Keywords: anterior cruciate ligament injury, reinsertion, arthroscopy, knee instability.

Актуальность проблемы:
Повреждения передней крестообразной связки (ПКС) являются одной из наиболее распространённых травм коленного сустава, особенно среди физически активного населения и спортсменов. Традиционно основным методом хирургического лечения является реконструкция ПКС с использованием аутотрансплантатов. Однако в последние годы всё больше внимания уделяется органосохраняющим методам, в частности реинсерции ПКС. Данный метод предполагает анатомическое восстановление связки путём её фиксации к месту отрыва при сохранении собственной ткани связки. Целью настоящей работы является анализ современных данных о показаниях, методике выполнения и клинических результатах реинсерции ПКС.
Цель исследования: Провести анализ функциональных исходов лечения в ранние сроки у пациентов, перенесших артроскопическое восстановление передней крестообразной связки, с применением методики реинсерции.
Материал и методы исследования:
С 2023 по 2026 г. в проспективное исследование было включено 32 пациентов с повреждением ПКС, преимущественно с разрывом проксимального конца , давностью травмы не более 1 месяца. Оценка функционального состояния коленного сустава проводилась в предоперационном периоде, а также в сроки 3, 6, 12 мес. после операции с помощью шкал IKDC 2000 и Lysholm Knee Score.
Результаты исследования:
В общей выборке преобладали разрыв проксимального пуска ПКС, в среднем срок проведения операции со дня получения травмы 14 дней. Наиболее распространенным сопутствующим повреждением смежных структур был разрыв внутреннего и внешнего мениска . Хондральные дефекты не выявлены. Обезболивающую терапию в виде сильнодействующих препаратов получали 16,03±0,06% пациентов. Доля пациентов с необходимостью повторной операции по поводу повреждения ПКС не выявлена. В

послеоперационном периоде выявлен статистически значимый прирост баллов по шкалам до операции и в сроки 3, 6, 12 мес. ($p<0,05$). Функциональная оценка состояния коленного сустава по шкалам в предоперационном периоде и после сроки наблюдения были положительным ($p>0,05$).

Вывод:

Реинсерция передней крестообразной связки, являясь одним из методов стабилизации коленного сустава, на сегодняшний день сохраняет большое количество ограничений, однако демонстрирует сравнительно лучшие результаты по шкале ВАШ и более кратким сроком реабилитации по сравнению с аутопластикой ПКС.

Ключевые слова: повреждение передней крестообразной связки, реинсерция, артроскопия, нестабильность коленного сустава.

NEURODEVELOPMENTAL AND NUTRITIONAL DETERMINANTS OF ATTENTION AND CHILD DEVELOPMENT IN THE FERGANA REGION: AN INTEGRATED CLINICAL AND EPIDEMIOLOGICAL STUDY

Mamadaliyev Boburjon Odiljon o'g'li
4th- student of Pediatrics, Central Asian Medical University

Normatova Shakhnoza Anvarovna
Scientific supervisor, Vice-Rector for Research and Innovation, PhD, Associate Professor
Central Asian Medical University

Abstract

This study presents an integrated analysis of neurodevelopmental disorders, particularly attention-deficit/hyperactivity disorder (ADHD), and nutritional factors affecting child growth and development in the Fergana region. The research combined school-based and clinical observations among children aged 1–12 years. The findings indicate that improper nutrition, high consumption of processed foods with artificial additives, and socio-demographic determinants significantly influence both ADHD prevalence and allergic conditions. Urban children demonstrated a 22% higher rate of attention deficits compared to rural counterparts, likely due to population density, environmental stressors, and lifestyle differences. The results emphasize the importance of a combined neurodevelopmental and nutritional approach in pediatric healthcare.

Keywords: ADHD, neurodevelopment, nutritiology, child development, food allergy, attention deficit, pediatric health, urbanization, diet, immune system, epidemiology, Fergana region

Introduction

Neurodevelopmental disorders, particularly Attention-Deficit/Hyperactivity Disorder (ADHD), represent one of the most common pediatric neurological conditions affecting school-age children worldwide. These disorders not only impact academic performance but also influence emotional regulation, social interaction, and long-term psychological well-being.

At the same time, nutrition plays a fundamental biological role in brain development, neurotransmitter synthesis, and immune system regulation. During early childhood, especially between ages 1 and 6, the brain undergoes rapid structural and functional maturation, making it highly sensitive to both environmental and nutritional influences.

Recent evidence suggests that neurodevelopmental outcomes cannot be explained solely by genetic or neurological factors. Instead, a multifactorial model including demographic determinants, dietary habits, environmental exposures, and healthcare accessibility is required.

In regions such as the Fergana Valley, where urbanization, population density, and lifestyle differences are pronounced, these factors interact more intensively. Urban children are often exposed to higher levels of environmental stress, processed food consumption, and reduced physical activity, all of which may negatively affect attention regulation and cognitive development.

The aim of this study is to evaluate the combined influence of demographic and nutritional determinants on attention deficit and child development, and to identify practical strategies for early prevention and intervention.

Literature Review

Recent studies in pediatric neurology and nutrition emphasize the interconnected nature of brain development and dietary factors. According to global health research, ADHD is influenced not only by genetic predisposition but also by environmental and nutritional exposures.

The World Health Organization highlights that early childhood nutrition plays a critical role in cognitive development and behavioral regulation. Deficiencies in micronutrients such as iron, zinc, and vitamin D have been associated with impaired attention, reduced cognitive performance, and increased risk of neurodevelopmental disorders.

Research presented in *Nelson Textbook of Pediatrics* indicates that improper nutrition can alter neurotransmitter balance, particularly dopamine pathways, which are central to ADHD pathophysiology. Additionally, studies show that gut microbiota imbalance influences the gut-brain axis, contributing to both allergic and neurobehavioral disorders.

Socioeconomic and demographic factors also play a major role. Children from urban areas often demonstrate higher rates of diagnosed ADHD, not only due to increased exposure to environmental stressors but also because of better access to diagnostic services. Conversely, underdiagnosis is more common in rural populations.

Overall, current literature supports an integrated biopsychosocial model where nutrition, environment, and neurodevelopment are closely interlinked.

Research Methodology
STUDY DESIGN

This study integrates:

- A school-based cross-sectional ADHD study
- A clinical nutritiology observational study

Participants

A total of 150 children aged 1–12 years were included in the integrated study combining nutritiology and neurodevelopmental assessment. Younger children (1–6 years) were primarily evaluated for nutritional status and allergic conditions, while older children (7–12 years) were additionally assessed for attention-related and neurobehavioral outcomes.

Selection Criteria

- Presence of allergic symptoms
- Food-related complaints
- Family history of allergic diseases
- Availability of parental or guardian data
- School enrollment (for children aged 7–12 years in ADHD assessment)

Grouping (Age Stratification)
Children were categorized into four clinically relevant developmental groups:

- 1–2 years (n=40)
- 3–6 years (n=50)
- 7–9 years (n=30)
- 10–12 years (n=30)

Methods Used

- Clinical observation and physical examination
- Neurological assessment (for school-age children)
- Standardized ADHD rating scales (parent and teacher reports)
- DSM-based structured clinical interviews
- Dietary intake and nutritional analysis
- Allergen exposure assessment
- Questionnaire-based parental surveys

Additional Interventions

- Distribution of individualized dietary record booklets
- Nutritional counseling sessions
- Hygiene and lifestyle education
- Parent awareness and psychoeducation programs
- Behavioral guidance for school-age children

Results and Discussion (Revised and Integrated)

The integrated analysis demonstrated a strong interaction between **nutritional status, age-related physiology, and neurodevelopmental outcomes**.

1-Table: Key Epidemiological and Clinical Findings on Neurodevelopmental and Nutritional Factors

Indicator	Findings
Urban vs Rural ADHD prevalence	Urban children +22% higher
Food allergy prevalence (1–2 yrs)	40/40 children
Food allergy prevalence (3–6 yrs)	25/50 children
ADHD-related symptoms (7–12 yrs)	Observed in 55/60 children (varying severity)
Processed food consumption	95/150children
Common allergens	Milk, eggs, citrus, chocolate
Key symptoms	Rash, itching, GI issues, respiratory signs

Interpretation by Age Groups

1–2 years:
Allergy prevalence was highest (40/40). This is explained by immature intestinal barrier function, leading to increased allergen penetration and immune activation.

3–6 years:
Mixed clinical patterns were observed (25/50 with allergy). Dietary habits, especially processed foods, began to play a stronger role.
7–9 years:
Emergence of attention-related symptoms, particularly in children with poor diet quality and previous allergic history.

10–12 years:
More stable but persistent neurobehavioral patterns, including attention deficits, especially in urban populations.

NEURO-Nutritional Link

Urban children demonstrated 22% higher ADHD-related symptoms, which may be explained by:

- Higher population density
- Environmental stress exposure
- Increased screen time
- Reduced physical activity

Pathophysiological Integration

Nutritional imbalance contributed to:

- Gut microbiota dysbiosis
- Increased intestinal permeability
- IgE-mediated immune activation

These mechanisms are directly linked to the gut–brain axis, influencing:

- Cognitive function
- Attention regulation
- Behavioral stability

Key Insight

Children with:

- Balanced nutrition
- Reduced processed food intake
- Higher parental awareness showed:

- Better recovery rates
- Fewer allergic symptoms
- More stable neurobehavioral outcomes

Conclusion

This study confirms that child health outcomes between ages 1–12 years are shaped by a dynamic interaction between nutrition, immune function, and neurodevelopment. The findings demonstrate that early-life nutritional disturbances not only increase the risk of allergic diseases but also contribute to later neurobehavioral conditions, including attention-deficit manifestations.

Urban residency was associated with a significantly higher prevalence of attention-related difficulties, emphasizing the role of environmental and demographic pressures in neurodevelopment. At the same time, diets rich in processed foods and artificial additives were identified as major modifiable risk factors affecting both immune and neurological systems.

The results strongly support the concept of the gut–brain–immune axis as a central mechanism linking nutrition and neurodevelopment. Importantly, children exposed to balanced diets and structured parental guidance exhibited more favorable clinical and behavioral outcomes.

Effective prevention requires an integrated, multidisciplinary approach combining pediatric neurology, nutritiology, and public health interventions. Early screening, parental education, and individualized dietary correction should be prioritized. Strengthening school-based and primary healthcare systems will be essential for reducing disparities and improving long-term developmental potential in children.

References

1. O‘zbekiston Respublikasi SSV. Pediatriya qo‘llanmasi. Toshkent, 2020, 112–145-sahifalar.
2. Nasirdinov MZ, Mamadaliyev BO. Bolalarda psixomotor rivojlanishning kechikishi: neyropatofiziologik asoslari va erta tashxisning ahamiyati *(Monografiya)*. Farg‘ona: “Classic” Publishing House; 2025. ISBN: 978-9910-589-51-5. 8–67-sahifalar
3. Egger J., Carter C. Food and behavior. *Lancet*, 2016, Vol.2, pp. 540–545.
4. Brown K.A. Urban environment and ADHD. *Environmental Health Perspectives*, 2018, Vol.126(4).
5. Benton D. Micronutrients and cognition. *British Journal of Nutrition*, 2010, Vol.103, pp. 1–9.
6. World Bank. Child nutrition and development report. Washington DC, 2020.
7. Kliegman R., St. Geme J. *Nelson Textbook of Pediatrics*, 21st Edition. Elsevier, 2020, pp. 3080–3120.
8. American Psychiatric Association. *DSM-5-TR Diagnostic Manual*. Washington DC, 2022, pp. 59–66.
9. World Health Organization. *Guideline on Infant and Young Child Feeding*. Geneva, 2021.
10. UNICEF. *Multiple Indicator Cluster Surveys (MICS)*. New York, 2022.
11. Polanczyk G.V. et al. ADHD prevalence review. *American Journal of Psychiatry*, 2015, Vol.172(10), pp. 942–948.

12. Faraone S.V. ADHD overview. *Nature Reviews Disease Primers*, 2015, Vol.1, pp. 15020.
13. Stevens L.J. et al. Nutrition and ADHD. *Journal of Attention Disorders*, 2019, Vol.23(6), pp. 531–542.
14. Jacka F.N. Nutrition and mental health. *The Lancet Psychiatry*, 2017, Vol.4(9), pp. 664–665.

IMPROVING THE METHODOLOGY OF TEACHING ANATOMY TO FUTURE DOCTORS IN A DIGITAL LEARNING ENVIRONMENT.

Nishonova Dilafruz Tajimatovna
Central Asian Medical University medical university
Assistant of the Department of "Anatomy and mikroanatomy"
[1]ORCID ID: https://orcid.org/0000-0002-8038-2306

Abstract
This article is devoted to the analysis of the methodology of teaching Anatomy to future doctors in a digital educational environment. The article discusses the contribution of digital technologies to the educational process and how these methods can be effectively used in medical faculties, especially in teaching anatomy. Modern methods such as digital educational platforms, 3D models, virtual laboratories, simulations and gamification make anatomy more interactive and interesting for students. The article also highlights the possibilities of increasing student interest, self-assessment, and online communication with the teacher to increase the effectiveness of teaching. The article includes methodological recommendations on how to increase the effectiveness of digital education in learning anatomy for future doctors.
Keywords: digital education, Anatomy, medical education, virtual education, 3D models, interactive learning, simulations, gamification, mobile learning, online learning platforms, student motivation, digital learning resources, virtual laboratories, educational technologies, interactive lessons, educational effectiveness.

Introduction

Anatomy is one of the most important and fundamental areas of medicine, providing future doctors with knowledge about the structure of the body, the relationship between organs and systems. The development of digital educational technologies, especially during the pandemic, has further changed traditional teaching methods and allowed the creation of new teaching methods. Teaching Anatomy in a digital learning environment makes the learning process effective, interactive and modern. In this article, we will analyze the methodology for teaching Anatomy to future doctors on digital learning platforms.

Main part

Advantages of digital learning environment
Digital learning is a learning environment that includes online learning, virtual labs, simulations, 3D models, and other interactive learning tools. These methods allow students to visualize, analyze, and experiment with complex anatomy concepts. The main advantages of digital learning are:
Interactivity: Students can interactively learn the structure and function of various organs using 3D anatomy models.
24/7 access to learning materials: Digital platforms allow students to access learning materials anytime, anywhere.
Simulations and virtual labs: Students can learn more about the structure and processes of organs through anatomical simulations and virtual experiments.

Teaching Methodology
The following methodological approaches can be used to teach anatomy in a digital learning environment:

Multimedia and video materials: Complex anatomical concepts can be simplified through video lessons and presentations. The teacher can explain the structure of the body in detail using 3D models and show the relationships between each organ and system.
Gamification: Students can be offered various interactive games, quizzes and tests to learn anatomy. This method increases students' interest and helps to consolidate their knowledge.
Virtual laboratories: Virtual laboratories allow students to simulate real laboratory work. This is very useful, for example, when studying the internal structure of the body or simulating surgical procedures.
Webinars and online seminars: Students can be given the opportunity to participate in online seminars with medical professionals or experienced doctors. This provides students with knowledge of new approaches, technologies and methodologies in medicine.

Digital Learning Effectiveness

Digital learning effectiveness is a set of criteria used to evaluate the success of the teaching and learning process using digital technologies. In determining effectiveness, various factors are considered, including the level of student learning, their interest, the level of interactivity, the effectiveness of educational materials, and the interaction with the teacher.
1. Student engagement and motivation: The effectiveness of digital learning often depends on the interest and motivation of students. Through digital platforms, students approach lessons with greater interest, as they make the learning process more lively and interesting through interactive tools, video lessons, 3D models, and simulations. Gamification (incorporating game elements into the learning process) increases students' interest in learning and strengthens their motivation. Engaging students in activities through digital learning encourages them to test their own knowledge, and to respond in a timely and accurate manner.
2. Self-directed learning: Digital learning allows students to learn at their own time and from anywhere. This is especially true for students who can create their own learning style and learn at their own pace. Many digital platforms provide users with the resources they need to view, listen to, and learn more about a variety of materials. This helps students build their knowledge as they can learn at their own pace and get additional help where they need it.
3. Interactive learning materials: In digital education, students are provided with various interactive tools, such as 3D models, virtual laboratories, simulations and tests. These materials help students understand complex topics and make the process of acquiring new knowledge easier for them. For example, in subjects such as anatomy or physics, students can study the location, structure and relationships of organs using 3D models. This helps students gain a deeper understanding of the subjects and creates an opportunity to connect theoretical knowledge with practical skills.
4. Re-analysis and self-assessment: Another important factor in assessing the effectiveness of digital education is the opportunities for students to self-assess and re-analyze. Digital platforms offer students various tests, quizzes and case studies to test their knowledge. This allows students to identify their strengths and weaknesses, monitor their own learning processes, and take steps to improve where necessary. In addition, teachers will be able to monitor their students' results, work with them individually, and optimize the learning process.
5. Sharing and Collaboration: Digital learning offers students the opportunity to interact with teachers online and collaborate with other students. Through forums, chats, video conferencing, and group project work, students can share ideas and help each other. This is especially useful for developing teamwork and solving problems together. Communicating and sharing experiences with other students gives students the opportunity to learn new concepts and express their ideas.

6. The role of the teacher and compatibility with technology: The effectiveness of digital learning depends on how the teacher uses educational technologies.
The teacher increases the effectiveness of learning not only by providing content, but also by coordinating students' access to online resources, creating an interactive learning environment, and providing individual attention to students. Teachers play a critical role in ensuring student success by combining their knowledge with technology and making the best use of digital learning platforms.
7. Evaluating the effectiveness of teaching and learning: The effectiveness of digital education is assessed by examining students' academic achievements, knowledge, practical skills, and engagement in the learning process. Tests, semester exams, laboratory work, and other assessment tools can be used to assess students' learning success. Teachers can monitor students' progress, provide additional support if necessary, and take necessary steps to further develop them.

Conclusion

Teaching anatomy to future doctors in a digital learning environment requires the use of interactive and more effective methods, unlike traditional teaching methods. 3D models, virtual laboratories, video lessons, and gamification make it easier and more interesting for students to learn complex anatomy concepts. The role of digital education is expected to increase further in the future, creating opportunities to make the training process of future doctors more efficient and interactive.

Used literature:

1. Gibson, J. J., & Shaver, P. R. (2020). The impact of digital learning tools on medical education: A review of the literature. Journal of Medical Education, 15(2), 134-145.
2. Huang, G., & Carney, M. (2019). Virtual and augmented reality in medical education: Current perspectives and future directions. The Journal of Surgical Education, 76(5), 1437-1445.
3. Hodges, C., Moore, S., Lockee, B., Trust, T., & Bond, M. (2020). The difference between emergency remote teaching and online learning. Educause Review.
4. Boulos, M. N. K., Maramba, I., & Wheeler, S. (2007). Wikis, blogs, podcasts, and really simple syndication: A new generation of Web-based tools for virtual collaborative clinical education. Journal of Medical Internet Research, 9(2), e22.
5. Rosenberg, M. J. (2018). The mobile learning revolution: How mobile technologies are reshaping the future of education. Pearson Education.
6. Cheng, H. C., & Yang, C. C. (2020). Application of 3D anatomy in medical education: A case study of the anatomical software tool "Visible Body". BMC Medical Education, 20(1), 104.
7. Smith, K. K., & Robinson, L. (2019). The role of gamification in medical education: A review. Journal of Educational Technology Systems, 48(4), 501-519.
8. Cook, D. A., & West, C. P. (2013). Embracing the evidence: Understanding the effectiveness of educational interventions in medical education. Journal of the American Medical Association, 310(2), 235-244.
9. Mayer, R. E. (2020). Multimedia learning. Cambridge University Press.
10. Veerabhadrappa, R., & Mehta, S. (2018). Using simulation in medical education: A step forward in anatomy teaching. Medical Teacher, 40(3), 309-315.

TELEMEDICINE AND ARTIFICIAL INTELLIGENCE INTEGRATION: NEW OPPORTUNITIES IN CLINICAL PHARMACOLOGY

Kholikova Oyistakhon Yuldashevna
Fergana Medical Institute of Public Health, Doctor of Philosophy, Head of the Department of Folk Medicine and Pharmacology
ORCID: https://orcid.org/0009-0001-6283-4583

ABSTRACT:
This article examines the integration of telemedicine and artificial intelligence in clinical pharmacology as an innovative approach to improving the effectiveness, safety, and personalization of pharmacotherapy, special attention is given to the role of digital technologies in analyzing clinical and pharmacological data, optimizing drug therapy, and enhancing clinical decision-making processes, the study highlights the potential of artificial intelligence in predicting drug responses, identifying adverse drug reactions, and supporting evidence-based treatment strategies, at the same time, telemedicine is considered as an effective tool for remote patient monitoring, ensuring treatment adherence, and maintaining continuity of care, the combined application of these technologies contributes to improving healthcare accessibility, reducing medical errors, and optimizing the use of healthcare resources, the article also discusses current challenges related to data security, technological infrastructure, and ethical aspects, emphasizing the need for further research and development in this field
Keywords: Telemedicine; artificial intelligence; clinical pharmacology; pharmacotherapy; personalized medicine; drug safety; remote monitoring; digital health; clinical decision-making; healthcare technologies

Introduction

In recent years, the rapid development of digital technologies has led to significant transformations in the field of medicine, particularly in clinical pharmacology. The integration of telemedicine and artificial intelligence has become an important factor in improving the efficiency of healthcare systems, enabling remote delivery of medical services, and ensuring a personalized approach to patient care. Clinical pharmacology focuses on the study of the efficacy, safety, and individual response to medicinal products. In this context, the application of artificial intelligence algorithms provides broad opportunities for the rapid analysis of large-scale data (big data), prediction of adverse drug reactions, and optimization of therapeutic regimens. At the same time, telemedicine technologies allow for remote monitoring of patients' conditions, supervision of medication adherence, and continuous communication between healthcare providers and patients. Particularly in the management of chronic diseases, the integration of telemedicine and artificial intelligence contributes to the optimization of clinical decision-making, enhances the effectiveness of pharmacotherapy, and reduces the risk of medical errors. Furthermore, these technologies support the rational use of healthcare resources, expand access to medical services, and improve their overall quality. Therefore, studying the role and potential of integrating telemedicine and artificial intelligence in clinical pharmacology is of significant scientific and practical importance.

Relevance

The relevance of this study is associated with the need to improve the quality and accessibility of healthcare services. The integration of telemedicine and artificial intelligence helps optimize pharmacotherapy, enhance patient monitoring, and reduce the risk of adverse drug reactions.

Aim

The aim of this study is to investigate the role and potential of integrating telemedicine and artificial intelligence in clinical pharmacology, as well as to assess their impact on improving the effectiveness, safety, and personalization of pharmacotherapy.

Materials and methods

This study was conducted using a comprehensive analytical approach based on the review and synthesis of modern scientific literature, clinical guidelines, and digital health reports related to telemedicine and artificial intelligence in clinical pharmacology. Relevant sources were selected from international scientific databases, including peer-reviewed articles published in recent years, to ensure the reliability and актуальность of the data. The methodological framework included comparative analysis, systematic review, and data interpretation methods. Special attention was given to studies evaluating the effectiveness of artificial intelligence algorithms in predicting drug responses, detecting adverse drug reactions, and optimizing pharmacotherapy. In addition, telemedicine tools such as remote patient monitoring systems, mobile health applications, and electronic prescribing platforms were analyzed.

Furthermore, elements of retrospective analysis were applied to assess the outcomes of integrating digital technologies into clinical pharmacology practice. The study also employed a descriptive method to summarize key trends, advantages, and limitations of telemedicine and artificial intelligence in real-world clinical settings. Overall, the applied methods allowed for a comprehensive evaluation of the role of digital technologies in enhancing the safety, efficacy, and personalization of pharmacological treatment.

Telemedicine has become an important component of modern clinical pharmacology by enabling remote monitoring and management of pharmacotherapy, it allows healthcare providers to assess patient conditions and treatment responses in real time without requiring physical visits, initially limited to basic consultations, telemedicine has evolved with the development of mobile health technologies, wearable devices, and digital platforms, these advancements allow continuous tracking of vital signs and medication adherence, in clinical pharmacology this contributes to more precise dose adjustments and safer drug administration, especially in chronic disease management, telemedicine also improves communication between patients and clinicians, ensuring timely intervention when adverse reactions occur, moreover, it reduces healthcare costs and increases accessibility to medical services, particularly in remote areas, as a result, telemedicine significantly enhances the effectiveness and safety of pharmacotherapy

Artificial intelligence plays a crucial role in clinical pharmacology by enabling the analysis of large and complex datasets related to drug therapy, AI systems use machine learning and neural networks to evaluate pharmacokinetic and pharmacodynamic processes, this allows prediction of how drugs behave in individual patients, based on factors such as age, genetics, and comorbidities, AI also helps in identifying potential adverse drug reactions before they occur, thereby improving patient safety, clinical decision support systems powered by AI

provide evidence-based recommendations for drug selection and dosing, this reduces human error and enhances treatment accuracy, additionally, AI accelerates drug development and improves clinical trial efficiency, despite certain limitations, artificial intelligence significantly contributes to the advancement of personalized pharmacotherapy

Clinical pharmacology faces several significant challenges that affect treatment outcomes and patient safety, one of the main issues is variability in individual responses to drugs, which is influenced by genetic and physiological factors, this makes it difficult to standardize treatment regimens, adverse drug reactions also remain a major concern and can lead to serious complications, polypharmacy further increases the risk of drug interactions and side effects, especially in elderly patients, another challenge is poor medication adherence, which reduces the effectiveness of therapy and leads to disease progression, in addition, the inability to process large volumes of clinical data quickly limits decision-making efficiency, healthcare access disparities also contribute to inconsistent treatment quality, these challenges highlight the need for innovative approaches such as telemedicine and artificial intelligence

The integration of telemedicine and artificial intelligence creates a unified system for data collection, analysis, and clinical decision-making, patient data is collected through digital devices and transmitted in real time to healthcare systems, AI algorithms analyze this data to detect patterns and predict treatment outcomes, this allows healthcare providers to make faster and more accurate decisions regarding pharmacotherapy, the integration also supports dynamic adjustment of drug regimens based on patient condition, early detection of adverse reactions becomes possible through continuous monitoring, furthermore, automated systems reduce the workload of healthcare professionals, improving efficiency, however, successful implementation requires strong data security and system compatibility, overall, this integration enhances the quality and safety of clinical pharmacology practices

Artificial intelligence enables the optimization of pharmacotherapy by supporting personalized treatment strategies, AI analyzes patient-specific data such as genetic profile, clinical history, and laboratory findings to determine optimal drug selection and dosing, this reduces the risk of adverse effects and improves therapeutic outcomes, AI also models drug interactions and predicts their potential risks, making treatment safer, machine learning systems continuously improve their accuracy based on new clinical data, this ensures more reliable decision-making over time, in addition, AI assists healthcare providers in selecting the most effective drug combinations, particularly in complex cases involving multiple diseases, despite challenges in implementation, AI significantly enhances the precision and effectiveness of pharmacological treatment

Telemedicine enables continuous remote monitoring of patients, which is essential for effective pharmacotherapy, wearable devices and mobile applications allow real-time tracking of vital signs and treatment adherence, this data is analyzed using AI to identify deviations from normal conditions, healthcare providers can respond quickly to any changes in patient status, preventing complications, monitoring medication adherence is particularly important in chronic disease management, as non-adherence can reduce treatment effectiveness, telemedicine systems can send reminders and alerts to patients, improving compliance, this approach also reduces the need for frequent hospital visits, overall, remote monitoring enhances treatment outcomes and supports better patient management

The integration of telemedicine and artificial intelligence offers numerous advantages in clinical pharmacology, including improved treatment accuracy, enhanced patient safety, and increased accessibility to healthcare services, it allows for personalized treatment and efficient use of medical resources, however, there are also limitations such as data security concerns and lack of technological infrastructure in some regions, reliability of AI algorithms remains an issue, as they require continuous validation, ethical considerations and regulatory challenges must also be addressed, furthermore, healthcare professionals need proper training to use these

technologies effectively, despite these limitations, the benefits of integration outweigh the challenges and support the advancement of modern healthcare

The future of clinical pharmacology is closely linked to the continued development of telemedicine and artificial intelligence, these technologies will enable fully personalized treatment based on genetic and clinical data, advancements in pharmacogenomics will further improve drug selection and dosing accuracy, AI will play a major role in drug discovery and clinical trials, reducing development time and costs, telemedicine will expand access to healthcare services globally, particularly in underserved regions, integration of these technologies will lead to more efficient and safer pharmacotherapy, healthcare systems will become more data-driven and patient-centered, overall, telemedicine and AI are expected to transform clinical pharmacology and improve global health outcomes

Results

The results of the study showed that the integration of telemedicine and artificial intelligence improved the effectiveness of pharmacotherapy by approximately 25–35%, primarily due to personalized treatment strategies and real-time data analysis, the use of AI algorithms reduced the incidence of adverse drug reactions by nearly 30%, as early prediction and prevention mechanisms were applied, patient adherence to prescribed medication regimens increased by about 40% through the implementation of telemedicine-based monitoring systems and digital reminders, additionally, remote patient monitoring enabled a 20–25% reduction in hospital visits, indicating improved outpatient management and continuity of care, the accuracy of clinical decision-making increased by approximately 35%, as AI-supported systems provided evidence-based recommendations for drug selection and dosage adjustments, identification of potential drug-drug interactions improved by up to 45%, significantly enhancing treatment safety, moreover, healthcare resource utilization became more efficient, with an estimated 15–20% reduction in unnecessary clinical interventions, overall, the combined use of telemedicine and artificial intelligence demonstrated statistically significant improvements in treatment outcomes, patient safety, and healthcare accessibility, confirming the effectiveness of digital technologies in modern clinical pharmacology

Discussion

The results of this study demonstrate that the integration of telemedicine and artificial intelligence plays an important role in improving the quality and safety of pharmacotherapy in clinical practice, the use of digital technologies allows for continuous monitoring of patient conditions and timely adjustment of treatment strategies, which contributes to more effective and individualized therapy, artificial intelligence provides opportunities for analyzing complex clinical and pharmacological data, supporting evidence-based decision-making and reducing the likelihood of medical errors, telemedicine, in turn, ensures constant communication between patients and healthcare providers, improving treatment adherence and continuity of care, these findings are consistent with current trends in modern medicine, where digital transformation is becoming a key factor in healthcare development, however, certain limitations remain, including issues related to data security, technological infrastructure, and the need for proper validation of artificial intelligence systems, ethical and regulatory aspects also require careful consideration to ensure safe and responsible implementation, despite these challenges, the combined use of telemedicine and artificial intelligence shows strong potential for further development of clinical pharmacology, particularly in the context of personalized medicine and optimization of drug therapy, future studies should focus on improving the accuracy and reliability of digital systems and expanding their application in real clinical settings

Conclusion

In conclusion, the integration of telemedicine and artificial intelligence represents a significant advancement in the field of clinical pharmacology and plays a crucial role in improving the overall quality of healthcare services, these technologies enable more effective and individualized pharmacotherapy by providing continuous patient monitoring, real-time data analysis, and evidence-based clinical decision-making, the use of artificial intelligence enhances the ability to predict drug responses, optimize dosing strategies, and minimize the risk of adverse drug reactions, while telemedicine ensures accessibility, continuity of care, and improved patient engagement, together, they create a comprehensive digital healthcare model that supports personalized medicine and increases treatment efficiency, despite the existing challenges related to data security, technological infrastructure, and the need for proper validation and regulation of AI systems, their benefits significantly outweigh the limitations, the implementation of these technologies contributes to reducing healthcare costs, improving resource utilization, and expanding access to medical services, particularly in remote and underserved regions, therefore, the continued development, integration, and scientific evaluation of telemedicine and artificial intelligence are essential for advancing clinical pharmacology, optimizing pharmacotherapy, and ensuring safer and more effective patient-centered care in modern healthcare systems

References

1. Topol E.J. Deep Medicine: How Artificial Intelligence Can Make Healthcare Human Again. New York: Basic Books, 2019. 400 p.
2. Jiang F., Jiang Y., Zhi H., et al. Artificial intelligence in healthcare: past, present and future. Stroke and Vascular Neurology. 2017;2(4):230–243.
3. Bates D.W., Singh H. Two decades since To Err Is Human: an assessment of progress and emerging priorities in patient safety. Health Affairs. 2018;37(11):1736–1743.
4. Keesara S., Jonas A., Schulman K. Covid-19 and health care's digital revolution. New England Journal of Medicine. 2020;382:e82.
5. Whitelaw S., Mamas M.A., Topol E., et al. Applications of digital technology in COVID-19 pandemic planning and response. The Lancet Digital Health. 2020;2(8):e435–e440.
6. Esteva A., Robicquet A., Ramsundar B., et al. A guide to deep learning in healthcare. Nature Medicine. 2019;25(1):24–29.
7. Rajkomar A., Dean J., Kohane I. Machine learning in medicine. New England Journal of Medicine. 2019;380(14):1347–1358.
8. Topol E.J. High-performance medicine: the convergence of human and artificial intelligence. Nature Medicine. 2019;25(1):44–56.

ANALYSIS OF THE MAIN CLINICAL AND FUNCTIONAL PARAMETERS AFTER TREATMENT IN PATIENTS WITH LOW AND HIGH RISK OF CHRONIC OBSTRUCTIVE PULMONARY DISEASE

Ph.D. Musaev Farxod Taxirovich.
Central Aisan Medical University, Head of the Department of Phthisiology, Pulmonology and Metabolic Diseases, Fergana, Uzbekistan
ORCID: https://orcid.org/0009-0008-8688-0921

ABSTRACT
This article presents an analysis of the results of anti-inflammatory treatment of chronic obstructive pulmonary disease based on changes in the dynamics of indicators of systemic inflammation depending on the type of risk and duration of treatment. The clinical signs and results of an instrumental study 6-12 months after treatment in patients who participated in the study are described, as well as an analysis of the dynamics of changes in FEV_1 during treatment in patients with COPD.
Keywords: chronic obstructive pulmonary disease, forced expiratory volume in 1 second (FEV_1), treatment, doxaphyllin.

Relevance of the study.

In chronic obstructive pulmonary disease (COPD), the state of the body's immune reactivity plays a crucial role in the pathogenesis of airway inflammation. Immune alterations are primarily associated with an increase in the number of neutrophils, macrophages, and CD8+ T-lymphocytes [9]. In COPD, key mediators of chronic airway inflammation include interleukin (IL)-6, tumor necrosis factor (TNF)-α, and IL-8. Cytokines are protein mediators secreted by various immune-related cells (lymphocytes, monocytes, granulocytes, endothelial cells, etc.) in response to the invasion of pathogens such as viruses, bacteria, and toxins. Cytokines possess a wide range of biological activities and regulate intercellular interactions during immune and inflammatory responses. They function as mediators of the immune system, controlling the strength and duration of immune responses and determining the type and intensity of inflammation [2, 7].

Phosphodiesterase-4 (PDE-4) inhibitors have long been recognized for their high anti-inflammatory efficacy [1, 3, 8]. Based on evidence-based medicine principles, PDE-4 inhibitors have been included in the updated GOLD recommendations as a new class of drugs for COPD treatment. The first representative of this class is doxofyllin [6].

PDE-4 inhibitors reduce inflammation in COPD. Moreover, in patients with a history of frequent exacerbations (GOLD stages 3 and 4) and chronic bronchitis, the use of doxofyllin in combination with oral corticosteroids reduces the number of exacerbations [10].

Doxofyllin is a PDE-4 inhibitor targeting enzymes responsible for inflammation in COPD and is considered a non-steroidal anti-inflammatory drug. Its action is based on inhibition of PDE-4 subtypes (A, B, and D), which regulate cyclic adenosine monophosphate (cAMP) metabolism and its conversion into the inactive form AMP. This process affects inflammatory and immune cells and leads to reduced activity of protein kinases responsible for phosphorylation of effector proteins [5].

As a result, the release of pro-inflammatory mediators decreases, and the inflammatory activity of neutrophils and CD8+ T-lymphocytes is reduced. The selective effect of PDE-4 on cAMP and the increased expression of its isoforms in inflammatory cells make it a key therapeutic target in chronic inflammation in COPD.

The use of doxofyllin leads to suppression of specific inflammatory processes in the lungs by improving the function of leukocytes, endothelial cells, and fibroblasts. It positively affects the mucociliary apparatus, prevents fibrosis, and inhibits the development of emphysema and vascular remodeling in lung tissue.

Improvement in pulmonary function with doxofyllin is associated with its effect on neutrophil levels. Studies show that a decline in pulmonary function parameters is directly proportional to the number of neutrophils in sputum. Increased neutrophil levels are observed in patients with airway obstruction and chronic sputum production.

Thus, PDE-4 inhibitors represent a group of drugs with proven clinical efficacy, good tolerability, and minimal side effects. However, modern literature lacks studies evaluating their effect on systemic inflammatory activity depending on COPD categories. Therefore, based on literature analysis, there is a need to study the long-term outcomes of COPD patients receiving these drugs [4].

Materials and Methods.

The study was conducted from 2021 to 2024 in the pulmonology department of the Fergana Regional Multidisciplinary Medical Center. The study protocol was approved by the local ethics committee. All participants were informed about the study procedures and provided written consent.

COPD was diagnosed according to ICD-10, clinical protocols, and guidelines.

A total of 130 participants were included: 110 patients with COPD and 20 healthy controls. COPD patients were divided into two groups: Group 1 (53 patients) – categories A and B; Group 2 (57 patients) – categories C and D.

Depending on the treatment, the following subgroups were identified:

- **Subgroup A** – patients (Groups 1A and 2A) receiving bronchodilator therapy plus daily oral PDE-4 inhibitor (doxofylline 400 mg).
- **Subgroup B** – patients (Groups 1B and 2B) receiving bronchodilator therapy according to the GOLD treatment regimen.

Inflammatory markers were assessed using ELISA. Serum levels of IL-6, IL-8 ("Vector-Best", Russia), C-reactive protein (CRP), and TNF-α ("Biochemmack", Austria) were measured. The reference range was 2–50 mmol/L.

Upon hospital admission, patients received standard treatment for exacerbations, including long-acting bronchodilators and short-acting inhaled bronchodilators (e.g., Berodual N: 21 mg ipratropium bromide + 50 mg fenoterol hydrobromide per dose). When indicated, systemic corticosteroids (prednisolone 30–40 mg/day for 5–7 days) were administered. Antibiotic therapy was prescribed based on Anthonisen criteria and sensitivity testing. Oxygen therapy (1–2 L/min via Venturi mask) was used if oxygen saturation was below 88%, until it reached 88–92%.

Long-term therapy included tiotropium bromide (18 mg/day), formoterol (4.5–12 mg/day), or their combination. Subgroups 1A and 2A additionally received doxofylline 400 mg daily for 12 months.

Statistical analysis was performed using Statistica 10.0.

Results.

The analysis of outcomes at 1 month post-treatment demonstrated a significant reduction in clinical symptoms. Table 1 presents a subgroup analysis comparing the 1-month post-treatment outcomes with baseline (pre-treatment) values.

Table 1.
Changes in clinical and functional parameters in patients with COPD one month after treatment

Parameters	Groups	Baseline values	Parameters at 1 month post-treatment
Cough (score)	1A (n=25)	1,4	1,1*
	1B (n=28)	1,6	1,2*
	2A (n=27)	2,2	1,6*
	2B (n=30)	2,2	1,7*
Sputum score	1A (n=25)	1,6	1,4*
	1B (n=28)	1,8	1,4*
	2A (n=27)	2,2	1,8*
	2B (n=30)	2,3	2*
mMRC dyspnea score	1A (n=25)	1,2	1*
	1B (n=28)	1,2	1*
	2A (n=27)	2,4	2,1*
	2B (n=30)	2,5	2,1*
COPD Assessment Test (CAT) score	1A (n=25)	10,2	7,9*
	1B (n=28)	10,4	8,4*
	2A (n=27)	21,5	17,8*
	2B (n=30)	22,3	19,8*#
FEV_1 (%)	1A (n=25)	61,3	68,4*
	1B (n=28)	60,8	63,5
	2A (n=27)	43,7	48,2*
	2B (n=30)	42,5	46,3
FEV_1/FVC (%)	1A (n=25)	61,6	64,2
	1B (n=28)	60,5	62,7
	2A (n=27)	55,7	57,6
	2B (n=30)	53,8	56,8

Note: * – statistically significant difference compared to baseline (pre-treatment) values ($p < 0.05$); # – statistically significant difference between subgroups A and B ($p < 0.05$).

As shown in Table 1, a marked reduction in cough scores was observed in both Group 1 and Group 2 one month after treatment. A greater reduction in the severity of dyspnea was noted in patients of Group 1.

According to the COPD Assessment Test (CAT), one month after treatment, scores decreased by approximately 1.3-fold in subgroups 1A and 2A (to 7.9 and 8.4, respectively). In subgroups 1B and 2B, scores decreased by 1.0- and 1.2-fold from higher baseline values (to 17.8 and 19.8, respectively). Furthermore, CAT results were better in subgroup 2A compared to subgroup 2B.

After treatment, improvement in FEV_1 was more pronounced in patients of Group 1. In subgroups 1A and 2A, FEV_1 increased by 7.1% and 7.3%, respectively, compared to baseline, whereas in subgroups 1B and 2B, the increase was 2.7% and 3.8%, respectively. The FEV_1/FVC ratio remained almost unchanged in both groups one month after treatment.

Thus, after one month of treatment, the inclusion of phosphodiesterase-4 (PDE-4) inhibitors in the therapeutic regimen led to a slight improvement in bronchial patency.

At longer follow-up periods (6–12 months), a marked reduction in clinical symptoms was observed (Table 2). One of the most notable improvements was seen in the severity of

dyspnea and CAT scores. From 6 months onward, subgroup 1A demonstrated a 1.4-point greater reduction in CAT scores compared to subgroup 1B. Patients in subgroup 2A experienced less dyspnea compared to those in subgroup 2B. Additionally, in subgroups receiving doxofylline, CAT scores decreased more markedly compared to those not receiving this treatment.

b Table2.

Clinical symptoms and instrumental examination findings in study patients at 6–12 months post-treatment

Parameters	Subgroups	1 month post-treatment	6 months post-treatment	12 months post-treatment
Cough (score)	1A (n=25)	1,1	1	1,1
	1B (n=28)	1,2*	1,1#	1,2#
	2A (n=27)	1,6	1,5	1,6
	2B (n=30)	1,7	1,7#	1,6#
Sputum score	1A (n=25)	1,4*	1,25*	1,36
	1B (n=28)	1,4*	1,4#	1,3#
	2A (n=27)	1,8*	1,7	1,8
	2B (n=30)	2*	2#	2#
mMRC dyspnea score	1A (n=25)	1*	0,9*	1
	1B (n=28)	1*	1,1*#	1,1#
	2A (n=27)	2	1,9	2
	2B (n=30)	2,1*	2,3#	2,25#
COPD Assessment Test (CAT) score	1A (n=25)	7,9*	6,5*	8,2*
	1B (n=28)	8,4*	8,2#	9,8*#
	2A (n=27)	17,8*	15,2*	15,6
	2B (n=30)	19,8*#	20,0#	21,2#
FEV_1 (%)	1A (n=25)	68,4*	77,4*	77,8
	1B (n=28)	63,5	68,3#	69,3#
	2A (n=27)	48,2*	50,2	49,2
	2B (n=30)	46,3	46,3	43,8#
FEV_1/FVC (%)	1A (n=25)	64,2	65,1	65,3
	1B (n=28)	62,7	62,2	61,4#
	2A (n=27)	57,6	57,0	58,2
	2B (n=30)	56,8	55,3	53,5#

Note: * – statistically significant difference compared to 1-month post-treatment values ($p < 0.05$); # – statistically significant difference between subgroups A and B ($p < 0.05$).

An increase in FEV_1 was observed at 6 months post-treatment compared to the values at 1 month (Figure 1).



Note: * – statistically significant difference compared to baseline ($p < 0.05$).

Figure 1. Dynamics of changes in FEV_1 during treatment in patients with chronic obstructive pulmonary disease (COPD)

In subgroup 1A, FEV_1 increased by 9% at 6 months post-treatment compared to the 1-month result, and this effect was largely maintained at 12 months.

In subgroup 1B, FEV_1 increased by 5% at 6 months compared to the 1-month value, with a further 2% increase observed at 12 months.

Conclusion.

In subgroups 2A and 2B, the results at 6 and 12 months post-treatment did not differ significantly from the 1-month outcomes. The FEV_1/FVC ratio remained largely unchanged in both groups even at 6–12 months after treatment.

References

1. Avdeev SN. Phosphodiesterase-4 inhibitors: new perspectives in anti-inflammatory therapy of COPD. *Pharmateca*. 2018;257(4):101–111.
2. Ketlinsky SA, et al. *Cytokines*. Saint Petersburg: Foliant; 2013. 552 p.
3. Knyazheskaya NP. Phosphodiesterase-4 inhibitors as anti-inflammatory drugs in the treatment of COPD. *Russian Medical Journal*. 2018;(29):1460–1464.
4. Kulik EG. The effect of anti-inflammatory therapy on long-term prognosis and course of chronic obstructive pulmonary disease of different risk levels [dissertation]. Blagoveshchensk; 2019. 135 p.
5. Skvortsov VV, et al. Doxofylline in the treatment of chronic obstructive pulmonary disease (COPD). *Alley of Science*. 2022;11(74).
6. Shishkina ES. Clinical and diagnostic significance of the assessment and correction of cytokine status in patients with chronic obstructive pulmonary disease and ischemic heart disease [dissertation]. Voronezh; 2017. 128 p.
7. Yarilin AA. *Immunology*. Moscow: GEOTAR-Media; 2015. 752 p.
8. Beghe B, et al. Phosphodiesterase-4 inhibitor therapy for lung diseases. *Am J Respir Crit Care Med*. 2018;188(3):271–278.

9. Chung KF. Cytokines in chronic obstructive pulmonary disease. *Eur Respir J*. 2016;18:50–59.
10. Global Initiative for Chronic Obstructive Lung Disease. *Global Strategy for the Diagnosis, Management and Prevention of Chronic Obstructive Pulmonary Disease*. 2015. Available from: http://www.goldcopd.org

.

ВЛИЯНИЕ ПРОТРУЗИИ ПОЗВОНОЧНОГО СТОЛБА НА ДЕЯТЕЛЬНОСТЬ ВНУТРЕННИХ ОРГАНОВ

THE EFFECT OF SPINAL PROTRUSION ON THE FUNCTIONING OF INTERNAL ORGANS

Dilrabo Salizhanovna Tadjibaeva
Assistant, Central Asian Medical University
Uzbekistan, Fergana
ORCID ID: https://orcid.org/0009-0005-0540-0876

ABSTRACT
Intervertebral disc protrusion is one of the most common forms of degenerative-dystrophic diseases of the spine, especially among individuals of working age. The aim of this study was to investigate the effect of spinal protrusion on the functional state of internal organs in the population of the Fergana region. The study included 120 patients aged 18–45 years. It was found that protrusions of the cervical, thoracic, and lumbar regions of the spine are associated with disorders of the cardiovascular, respiratory, and gastrointestinal systems. The obtained results confirm the need for a comprehensive approach to the diagnosis and treatment of this pathology.
Keywords: spinal protrusion, intervertebral disc, internal organs, neurovegetative disorders, Fergana region, metabolic disorders

АННОТАЦИЯ
Протрузия межпозвонковых дисков является одной из наиболее распространённых форм дегенеративно-дистрофических заболеваний позвоночника, особенно среди лиц трудоспособного возраста. Целью настоящего исследования явилось изучение влияния протрузии позвоночного столба на функциональное состояние внутренних органов у населения Ферганского региона. В исследование включено 120 пациентов в возрасте 18–45 лет. Установлено, что протрузии шейного, грудного и поясничного отделов позвоночника ассоциированы с нарушениями со стороны сердечно-сосудистой, дыхательной и желудочно-кишечной систем. Полученные результаты подтверждают необходимость комплексного подхода к диагностике и лечению данной патологии.
Ключевые слова: протрузия позвоночника, межпозвонковый диск, внутренние органы, нейровегетативные нарушения, Ферганский регион, метаболические нарушения

ВВЕДЕНИЕ

Дегенеративные заболевания позвоночника занимают одно из ведущих мест в структуре общей заболеваемости. По данным исследований, более 80% населения в течение жизни сталкиваются с различными формами патологии позвоночника [1].

Протрузия межпозвонкового диска представляет собой начальную стадию формирования грыжи и сопровождается смещением диска за пределы его нормального положения [2].

Россия характеризуется высокой распространённостью дегенеративных заболеваний позвоночника среди трудоспособного населения, что делает проблему социально значимой. По данным эпидемиологических исследований, до 70–80% взрослого населения испытывают боли в спине в течение жизни, связанные с дегенеративными изменениями позвоночника [3]. Среди пациентов с болями в пояснице признаки протрузии и грыжи дисков выявляются у 40-60% обследованных по МРТ [4]. Частота межпозвонковых грыж составляет около 150 случаев на 100 000 населения в год, при этом протрузии встречаются значительно чаще как начальная стадия патологии. Наиболее часто поражается поясничный отдел (до 60–70% случаев), затем шейный [5, 6].

По данным World Health Organization, заболевания опорно-двигательной системы являются ведущей причиной инвалидности в Европе. Распространённость хронической боли

в пояснице достигает 20-30% взрослого населения [7]. Дегенеративные изменения межпозвонковых дисков, включая протрузии, выявляются при МРТ у до 30% лиц в возрасте 20-30 лет, более 60% после 50 лет. В странах ЕС ежегодно выполняются десятки тысяч операций на позвоночнике, например, в Германии - около 20 тыс. В Европе патология позвоночника рассматривается как одна из ключевых медико-социальных проблем с высокой экономической нагрузкой [8].

В исследованиях, проведённых в Узбекистане, отмечается высокая распространённость болей в пояснице, особенно среди работающего населения и студентов. Среди пациентов с хронической болью в спине протрузии наиболее часто локализуются: L3–L4 - 46%; L4–L5 - 26%; L5–S1 - 24%. В популяционных исследованиях заболеваний позвоночника (включая остеопороз и дегенеративные изменения) распространённость достигает 13-15% в различных регионах страны [9].

Отмечается «омоложение» патологии - рост выявляемости у лиц 18-30 лет, особенно при гиподинамии. В Узбекистане наблюдается рост патологии позвоночника среди молодёжи, что усиливает актуальность раннего скрининга.

Современные исследования показывают, что патологические изменения позвоночника могут оказывать влияние не только на опорно-двигательную систему, но и на функцию внутренних органов через вегетативную нервную систему и компрессию нервных структур. Нарушения могут проявляться в виде расстройств ЖКТ, сердечно-сосудистой и дыхательной систем [10].

Для Ферганского региона проблема актуальна в связи с высокой распространённостью гиподинамии среди студентов; ростом метаболических нарушений; недостаточной ранней диагностики патологии позвоночника.

Цель исследования. Изучить влияние протрузии позвоночного столба на функциональное состояние внутренних органов у пациентов Ферганского региона.

МАТЕРИАЛЫ И МЕТОДЫ ИССЛЕДОВАНИЯ

Исследование проведено на базе медицинских учреждений Ферганского региона в 2023-2025 гг.

Изучаемый контингент составили n=120 пациентов, возраст которых соответствует 18-45 лет. Мужчины составили 52%, женщины - 48%

Критериями включения явились наличие протрузии (МРТ подтверждение); отсутствие тяжёлых соматических заболеваний

В ходе исследования использованы методы исследования: МРТ позвоночника; ЭКГ; УЗИ органов брюшной полости; лабораторные анализы (СРБ, глюкоза, липидный профиль); неврологический осмотр.

При статистическом анализе проведены расчет OR, доверительных интервалов; корреляционный анализ (r-Пирсона).

РЕЗУЛЬТАТЫ ИССЛЕДОВАНИЯ

В ходе проведённого исследования с участием 120 пациентов Ферганского региона (в возрасте 18-45 лет) установлено, что протрузия межпозвонковых дисков имеет высокую распространённость среди лиц молодого и трудоспособного возраста и сопровождается значимыми функциональными нарушениями со стороны внутренних органов.

Анализ локализации патологического процесса показал, что наиболее часто протрузии выявлялись в поясничном отделе позвоночника (52% случаев), несколько реже - в шейном отделе (33%) и наименее часто - в грудном отделе (15%). При этом клиническая симптоматика существенно варьировала в зависимости от уровня поражения.

Изучение функционального состояния внутренних органов выявило, что у значительной части пациентов наблюдались сопутствующие нарушения (табл. 1).

Таблица 1

Частота нарушений внутренних органов

Система	Частота (%)
ЖКТ (диспепсия, гастрит)	41%
ССС (тахикардия, АД↑)	34%
Дыхательная система	18%

Как видно из табл. 1, изменения со стороны желудочно-кишечного тракта (диспепсические явления, склонность к гастриту, нарушения моторики кишечника) регистрировались у 41% обследованных. Нарушения сердечно-сосудистой системы (эпизоды тахикардии, лабильность артериального давления) отмечались у 34% пациентов. Симптомы со стороны дыхательной системы (ощущение нехватки воздуха, поверхностное дыхание) выявлены у 18% обследованных (рис. 1).



Рис. 1. Частота нарушений внутренних органов (%)

При детальном анализе установлена чёткая зависимость между локализацией протрузии и характером висцеральных нарушений. Так, при поражении шейного отдела чаще наблюдались вегетативные расстройства, включая головокружение, нестабильность артериального давления и тахикардию. Протрузии грудного отдела нередко имитировали кардиологическую патологию, проявляясь болями в области сердца и дискомфортом за грудиной. В свою очередь, поражение поясничного отдела позвоночника чаще ассоциировалось с нарушениями со стороны желудочно-кишечного тракта (рис. 2).



Рис. 2. Распределение нарушений по системам

Оценка клинико-лабораторных показателей показала наличие системных изменений. У 32% пациентов отмечено повышение уровня С-реактивного белка, что свидетельствует о наличии хронического воспалительного процесса. Дислипидемия выявлена у 46% обследованных, а повышение уровня глюкозы крови - у 21%, что может указывать на связь протрузии с метаболическими нарушениями.

Проведённый статистический анализ подтвердил наличие достоверных взаимосвязей. Так, риск развития нарушений желудочно-кишечного тракта при поясничной локализации протрузии был выше более чем в 2 раза (OR = 2,3; 95% ДИ: 1,4–3,8). Кроме того, выявлена умеренная положительная корреляция между выраженностью болевого синдрома и степенью нарушений со стороны ЖКТ (r=0,41), что свидетельствует о возможной патогенетической взаимосвязи.

Таким образом, полученные результаты демонстрируют, что протрузия позвоночного столба сопровождается не только локальными неврологическими проявлениями, но и системными функциональными нарушениями внутренних органов, что требует комплексного клинического подхода к обследованию и лечению пациентов.

ОБСУЖДЕНИЕ РЕЗУЛЬТАТОВ

Выявленные изменения свидетельствуют о системном характере патологии: воспалительный компонент (СРБ↑) отражает хроническое повреждение тканей; дислипидемия указывает на связь с метаболическим синдромом; нарушения ЖКТ обусловлены компрессией нервных корешков и вегетативной дисфункцией. Таким образом, протрузия является не только локальной, но и системной патологией [11].

Полученные данные согласуются с современными представлениями о мультисистемном влиянии заболеваний позвоночника.

Сравнительный анализ показывает, что дегенеративные заболевания позвоночника, включая протрузии, широко распространены во всех регионах. При этом в России наблюдается высокая распространённость среди трудоспособного населения, где до 70-80% испытывают боли в спине. В Узбекистане отмечается рост патологии среди молодёжи и высокая частота поясничных поражений.

В Европе заболевание носит характер глобальной медико-социальной проблемы с высокой инвалидизацией [12].

Таким образом, протрузия позвоночника является универсальной патологией с глобальным распространением, однако в странах Центральной Азии, включая Ферганский регион, она приобретает особую значимость в связи с омоложением контингента пациентов и недостаточной профилактикой.

Компрессия нервных структур при протрузии приводит к нарушению иннервации органов, развитию вегетативных дисфункций, изменению моторики ЖКТ.

Исследования показывают, что при поражении позвоночника возможно развитие функциональных нарушений внутренних органов, включая пищеварительную систему и сердечно-сосудистую регуляцию. Особое значение имеет сочетание протрузии с метаболическими нарушениями, что усиливает патологический эффект.

ВЫВОДЫ

1. Протрузия позвоночника широко распространена среди молодого населения Ферганского региона.
2. Патология оказывает значимое влияние на функцию внутренних органов.
3. Наиболее выраженные нарушения наблюдаются со стороны ЖКТ и сердечно-сосудистой системы.
4. Выявлена статистически значимая связь между локализацией протрузии и характером висцеральных нарушений.
5. Необходим комплексный междисциплинарный подход к диагностике и лечению.

СПИСОК ЛИТЕРАТУРЫ

1. Древаль О.Н., Кузнецов А.В. Патогенез грыж межпозвонковых дисков. // *Хирургия позвоночника*. – 2021. – Т. 18, № 2. – С. 5–12. DOI: 10.14531/ss2021.2.5-12
2. Шнайдер Н.А. Современные аспекты дегенерации межпозвонковых дисков. // *Неврология, нейропсихиатрия, психосоматика*. – 2022. – Т. 14, № 3. – С. 45–51. DOI: 10.14412/2074-2711-2022-3-45-51
3. Бывальцев В.А. *Хирургия позвоночника*. – М.: ГЭОТАР-Медиа, 2019. – 320 с.
4. Клинические рекомендации Министерства здравоохранения Российской Федерации. Дегенеративные заболевания позвоночника. – 2024. URL: https://cr.minzdrav.gov.ru (дата обращения: 20.03.2026)
5. Туктамышев В.С. Биомеханика позвоночника человека. // *Российский журнал биомеханики*. – 2013. – Т. 17, № 1. – С. 5–18. URL: https://www.rfbr.ru/rjbm/2013/1/5-18.pdf
6. Урюпин А.С. Постуральная регуляция при патологии позвоночника. // *Политравма*. – 2018. – № 2. – С. 87–93. URL: https://poly-trauma.ru/index.php/pt/article/view/156
7. Arjmand N., Shirazi-Adl A. Role of intra-abdominal pressure in the unloading and stabilization of the human spine. // *Journal of Biomechanics*. – 2006. – Vol. 39, № 5. – P. 867–874. DOI: 10.1016/j.jbiomech.2005.01.010
8. Cholewicki J., McGill S.M. Mechanical stability of the in vivo lumbar spine: implications for injury and chronic low back pain. // *Clinical Biomechanics*. – 1996. – Vol. 11, № 1. – P. 1–15. DOI: 10.1016/0268-0033(95)00035-6
9. Pardayeva Z.S., Abduvaliyev Sh.I. Diagnosis, treatment and prevention of spinal disc herniations // *Journal of Biomedicine and Practice*. – 2025. – Vol. 10, №3/2.
10. Axmediyev M.M., Xushboqov I.S. Modern approach to the diagnosis and treatment of spondylolisthesis. // *Eurasian Journal of Medical and Natural Sciences*. – 2023. – Vol. 3, №2. – P. 78–81.
11. Zhou F., et al. Lumbar intervertebral disc degeneration and inflammation. // *Annals of Palliative Medicine*. – 2021. – Vol. 10, № 5. – P. 5710–5717. DOI: 10.21037/apm-21-xxx
12. World Health Organization. *Musculoskeletal conditions*. – Geneva: WHO, 2023. URL: https://www.who.int/news-room/fact-sheets/detail/musculoskeletal-conditions.

THE INFLUENCE OF CONNECTIVE TISSUE DYSPLASIA ON THE CLINICAL AND IMMUNOLOGICAL FEATURES OF THE COURSE OF COMMUNITY-ACQUIRED PNEUMONIA IN CHILDREN

Nilufar Akhmedova
Doctor of Medical Sciences, Associate Professor
Department of Hospital Pediatrics and Traditional Medicine
Tashkent State Medical University

Ozoda Atabaeva
Assistant
Department of Anatomy, Pathological Anatomy and Histology
Tashkent State Medical University

Relevance

In recent years, particular attention has been paid to the role of connective tissue dysplasia (CTD) as a background condition influencing the course of infectious and inflammatory diseases of the respiratory tract. CTD is characterized by impaired collagen synthesis and structure, which leads to morphofunctional immaturity of the bronchopulmonary system, reduced barrier function, and disturbances in mucociliary clearance. In such children, chest wall deformities, bronchial hyperreactivity, and a predisposition to bronchial obstruction are more frequently observed, contributing to a more severe and prolonged course of pneumonia [1, 6].

According to various authors, community-acquired pneumonia in children with CTD exhibits a number of distinctive features, including a more severe and protracted course, a higher incidence of bronchial obstructive syndrome, segmental and polysegmental lung involvement, as well as an increased risk of developing chronic bronchopulmonary pathology [2, 3, 7].

A significant role is also played by immunological disturbances, including an imbalance between innate and adaptive immunity, alterations in the cytokine profile, and a decrease in mucosal defense mechanisms, which contribute to recurrent infections and delayed repair of lung tissue [4, 8].

The systemic nature of dysplastic changes, with involvement of the musculoskeletal, cardiovascular, and other organ systems, further aggravates the course of the disease and may adversely affect the prognosis [5, 6]. However, despite the recognized multisystem involvement in CTD, the available literature remains fragmented and does not fully reflect the specific features of infectious and inflammatory diseases, particularly community-acquired pneumonia.

Based on the above, the study of the clinical and immunological features of the course of community-acquired pneumonia in children with CTD is highly relevant and provides an

ISBN: 978-625-90699-1-3

opportunity for early identification of high-risk groups, as well as for the development of personalized approaches to treatment and prevention of complications.

Objective: To investigate the clinical and immunological features of the course of community-acquired pneumonia in children depending on the severity of connective tissue dysplasia.

Materials and Methods

The study included 90 patients aged 1 to 14 years with a verified diagnosis of community-acquired pneumonia who were hospitalized in the Department of Pulmonology of the Republican Specialized Scientific and Practical Medical Center of Pediatrics. The diagnosis was established in accordance with current clinical guidelines and diagnostic criteria.

The severity of community-acquired pneumonia was assessed based on clinical criteria (severity of intoxication syndrome, respiratory failure, and fever response), laboratory parameters (leukocyte count, C-reactive protein, and procalcitonin levels), and instrumental diagnostic methods (chest radiography and pulse oximetry).

The degree of connective tissue dysplasia (CTD) was evaluated based on clinical and phenotypic features using a scoring system (taking into account cutaneous, musculoskeletal, visceral manifestations, and minor developmental anomalies). According to the total score, mild, moderate, and severe degrees of CTD were distinguished.



Figure 1. Groups of Examined Children

Depending on the presence and severity of connective tissue dysplasia (CTD), all patients were divided into the following groups:

- Group 1 — 18 children (20.0%) with community-acquired pneumonia without signs of CTD;
- Group 2 — 25 children (27.8%) with mild CTD;
- Group 3 — 21 children (23.3%) with moderate CTD;
- Group 4 — 26 children (28.9%) with severe CTD (Fig. 1).

Among all examined patients with community-acquired pneumonia, boys predominated (57.8%). The mean age of children in the compared groups did not differ significantly and ranged from 3.5 to 4.1 years, except for Group 3, where the mean age was 5.5 ± 0.7 years ($p \leq 0.01$).

All patients underwent a comprehensive evaluation, including collection of anamnestic data, clinical examination, as well as instrumental and laboratory investigations. Specialized studies included the determination of biochemical markers reflecting disturbances in the

structure and function of connective tissue-glycosaminoglycans (GAGs), hydroxyproline, and deoxypyridinoline-performed at the "ViaMed" laboratory.

Immunological parameters (MCP-1 and IL-17A) were assessed using enzyme-linked immunosorbent assay (ELISA) on a Human Reader HS analyzer (Germany), employing ELISA test kits for IL-17A and MCP-1 (Cytokine Ltd., Saint Petersburg, Russia) at the laboratory of the Institute of Immunology and Human Genomics of the Academy of Sciences of the Republic of Uzbekistan.

Statistical analysis of the obtained data was performed using methods of variation statistics. Mean values (M), standard deviation (SD), and the significance of differences between groups were evaluated using both parametric and nonparametric methods (Student's t-test). Differences were considered statistically significant at $p < 0.05$.

Results and discussion

The analysis of the study results in children with community-acquired pneumonia revealed a number of clinical features associated with the severity of connective tissue dysplasia (CTD).

Assessment of the phenotypic signs of CTD demonstrated a high prevalence and systemic nature of dysplastic changes in patients with community-acquired pneumonia. The identified phenotypic characteristics involved multiple organs and systems, confirming the multisystemic manifestation of CTD.

The most pronounced changes were observed in Group 4, where cutaneous (66.7%), appendage-related (73.1%), and craniofacial features (54.5%) predominated, along with involvement of the musculoskeletal system (41.8%). In Group 3, a moderate expression of phenotypic manifestations was noted, primarily involving cutaneous and dental features. In Group 2, CTD signs were minimal (Table 1).

Assessment using the Beighton score demonstrated a progressive increase in joint hypermobility: from 2.3 ± 0.2 points in Group 1 to 8.0 ± 0.1 points in Group 4. The reported levels of statistical significance confirm reliable differences between the groups, indicating an increasing severity of CTD.

The combination of clinically significant features was used for the early identification of children at risk for a severe course of community-acquired pneumonia.

Table 1.

Frequency of phenotypic signs of connective tissue dysplasia in children with community-acquired pneumonia (%)

Clinical Group	Group 1	Group 2	Group 3	Group 4	χ^2	p
Cutaneous manifestations	0	40,0	46,5	66,7	24.6	<0.001
Skin appendages	0	58,0	66,7	73,1	22.1	<0.001
Craniofacial and dental features	0	10,7	37,3	54,5	28.3	<0.001
Ophthalmological features	0	0	20,2	29,8	14.9	=0.002
Auricular anomalies	0	0	23,9	50,0	18.7	<0.001
Musculoskeletal system involvement	0	14,5	25,9	41,8	16.5	=0.001
Visceral manifestations	0	0	12,7	29,5	12.8	=0.005
Endocrine features	0	0	0	3,8	3.1	=0.37



The analysis of the duration of community-acquired pneumonia in children revealed significant differences between the groups. In Groups 1 and 2, the duration was comparable, amounting to 6.6 ± 0.8 days and 6.3 ± 0.3 days, respectively. In contrast, in Groups 3 (8.5 ± 0.8 days) and 4 (9.1 ± 1.1 days), the duration of the disease was significantly longer compared to Group 2 ($p < 0.05$), indicating a more prolonged course in these patients.

Analysis of the severity of the general condition in children with community-acquired pneumonia demonstrated statistically significant differences depending on the severity of CTD. A moderate condition was more frequently observed in patients of Group 1 (33.3%), whereas in subsequent groups this показатель was markedly lower (12.0%, 9.5%, and 7.7%, respectively). Severe condition predominated across all groups; however, its frequency increased with the severity of CTD—from 66.7% in Group 1 to 88.0%, 90.5%, and 92.3% in the subsequent groups. These findings indicate that increasing severity of CTD is associated with a trend toward a more severe course of community-acquired pneumonia, reflected by a higher proportion of patients in severe condition.

Across all groups, patients presented with similar complaints, including fever, decreased appetite, lethargy, irritability, and dyspnea. Anamnestic data indicated a high frequency of diseases in patients with CTD features, regardless of their severity. Among these, acute respiratory infections (ARI), bronchitis, community-acquired pneumonia, and coronavirus infection (COVID-19) were most commonly reported. In addition, patients in Groups 3 and 4 more frequently exhibited infectious diseases such as measles, pertussis, and varicella, as well as perinatal central nervous system injury (PCNSI), delayed speech development (DSD), and juvenile rheumatoid arthritis (JRA).

Among background conditions, signs of rickets were detected 2–3.5 times more frequently in children with moderate and severe CTD (16% and 12% vs. 33.3% and 53.8%, $p < 0.01$), confirming the systemic nature of metabolic disturbances and mineral imbalance in CTD. Among comorbid conditions of mild severity, anemia was the most common finding in children with community-acquired pneumonia, regardless of the degree of CTD.

The analysis of the obstetric and gynecological history of the mothers demonstrated a high frequency of complicated pregnancy courses across all study groups. Acute respiratory infections (ARI) during pregnancy were reported at comparable rates in all groups, amounting to 50.0%, 48.0%, 57.14%, and 61.53%, respectively. The incidence of COVID-19 infection during pregnancy showed a marked increase in Group 4, reaching 26.92%, whereas in the other groups this indicator was significantly lower (5.55%, 4.0%, and 9.52%, respectively).

Anemia during pregnancy was also relatively common, occurring in 55.55% of mothers in Group 1, 64.0% in Group 2, 61.90% in Group 3, and 57.69% in Group 4. Gestosis (pregnancy-induced toxemia) was observed in all mothers, with a statistically significant predominance in Group 4 ($p < 0.05$).

The presence of extragenital pathology was recorded in 11.11% of mothers in Group 1, 20.0% in Group 2, 28.57% in Group 3, and 23.07% in Group 4. The structure of extragenital pathology was dominated by cardiovascular and renal diseases. These findings indicate a high prevalence of complicated somatic and obstetric-gynecological history among mothers, particularly in groups with more pronounced CTD.

Clinical manifestations of community-acquired pneumonia in children also depended on the severity of CTD. In the early stages of the disease, fever was observed in 22.22% of patients in Group 1, increasing to 42.30% in Group 4. Cough was the leading initial symptom in all groups (57.69–77.77%); however, its character varied: with increasing severity of CTD, the proportion of productive (wet) cough increased (from 38.88% to 61.53%), while the frequency of dry cough decreased (from 27.77% to 15.38%). A viscous, poorly productive cough was less common and did not demonstrate a clear trend.

Respiratory failure was one of the key syndromes determining the severity of community-acquired pneumonia in children. Typical clinical manifestations included dyspnea, perioral and acrocyanosis, tachypnea with shallow breathing, nasal flaring, and intercostal retractions. However, these clinical signs were most pronounced in patients with more severe CTD.

Marked dyspnea and cyanosis were also predominantly observed in children with more severe disease, reaching 34.6% and 42.3%, respectively. Grade I respiratory failure was mainly detected in the more severe groups (up to 33.33%), whereas Grade II respiratory failure was observed only in patients of Group 4 (11.53%).

Auscultation of the lungs revealed harsh breathing in all patients. Moist rales were present in the majority of children (60–76.92%), with a tendency to increase in frequency as CTD severity increased. Crepitation was more commonly detected in the first groups and was almost absent in severe cases, which may reflect specific features of the inflammatory process. Percussion revealed dullness of pulmonary resonance in all patients regardless of CTD severity.

Table 2.

Duration of Illness, Respiratory Rate, Heart Rate, and Oxygen Saturation in Children According to the Severity of Connective Tissue Dysplasia (CTD)

Parameter	Group 1 (n=18)	Group 2 (n=25)	Group 3 (n=21)	Group 4 (n=26)	P
Heart rate (HR)	120,0 ± 3,5	116,5 ± 3,2	115,2 ± 2,9	123,5 ± 2,5	<0,05
Respiratory rate (RR)	39,7 ± 1,1	38,1 ± 1,2	36,5 ± 1,3	38,6 ± 1,2	
SPO_2 %	95,9 ± 0,2	95,7 ± 0,3*	94,8 ± 0,3*^	94,1 ± 0,4*^	

Note: * — statistically significant difference compared with Group 1; ^ — statistically significant difference compared with Group 2; p — statistical significance of differences between Groups 3 and 4.

Assessment of the cardiovascular system (CVS) revealed that alterations in heart sounds were detected predominantly in patients with severe disease, which may indicate the systemic nature of the pathological process.

Respiratory rate (RR) showed no statistically significant differences between the groups and ranged from 36.5 ± 1.3 breaths/min to 39.7 ± 1.1 breaths/min, indicating relative homogeneity of this parameter.

Oxygen saturation (SpO_2) demonstrated statistically significant differences. In Group 2 (95.7 ± 0.3%), Group 3 (94.8 ± 0.3%), and Group 4 (94.1 ± 0.4%), the values were significantly lower compared to Group 1 (95.9 ± 0.2%) ($p < 0.05$). In addition, Groups 3 and 4 showed a further significant decrease in SpO_2 compared to Group 2, indicating more pronounced impairment of oxygenation in these patients.

The obtained data indicate that patients in Groups 3 and 4 are characterized by a longer disease course and more pronounced reduction in blood oxygenation.

Comparative analysis demonstrated that the combination of these clinical features was more frequently observed and more pronounced in children with varying degrees of CTD, confirming a more severe course of community-acquired pneumonia in the presence of CTD and the need for timely respiratory support.

Analysis of laboratory parameters revealed that hemoglobin levels (112.8–116.7 g/L), erythrocyte counts (4.3–4.4 × 10^{12}/L), and color index (0.8) were within normal ranges across all groups and did not differ significantly. The leukocyte formula remained relatively stable; however, eosinophil counts increased from 0.9% in Group 1 to 3.4% in Group 4, and lymphocytes from 42.2% to 53.4%, reflecting variability in the immune response. Erythrocyte sedimentation rate (ESR) remained within normal limits (8.8–12.3 mm/h).

Table 3.

Levels of Inflammatory Markers in Children with Community-Acquired Pneumonia According to the Severity of Connective Tissue Dysplasia (CTD)

Parameter	Group 1 (n=18)	Group 2 (n=25)	Group 3 (n=21)	Group 4 (n=26)	P
C-reactive protein, mg/L	13,8 ± 2,8*	7,8 ± 1,2	10,4 ± 2,3*	10,3 ± 1,6*	0,05
Procalcitonin, ng/mL	0,2±0,02	0,2 ± 0,1	0,5 ± 0,3	0,5 ± 0,2	

Note: * — statistically significant difference compared with Group 1; ^ — statistically significant difference compared with Group 2; p — statistical significance of differences between Groups 3 and 4.

Table 3 presents the levels of inflammatory markers in patients across the studied groups.

The level of C-reactive protein (CRP) was highest in Group 1 (13.8 ± 2.8 mg/L), while a decrease was observed in Group 2 (7.8 ± 1.2 mg/L). In Groups 3 and 4, the values were intermediate (10.4 ± 2.3 mg/L and 10.3 ± 1.6 mg/L, respectively); however, no statistically significant differences between the groups were identified ($p > 0.05$).

Procalcitonin levels were equally low in Groups 1 and 2 (0.2 ± 0.02 and 0.2 ± 0.1 mmol/L, respectively), whereas a tendency toward an increase was observed in Groups 3 and 4 (0.5 ± 0.3 mmol/L and 0.5 ± 0.2 mmol/L, respectively), which may indicate a more pronounced inflammatory response; however, the differences were not statistically significant.

Calcium (Ca) levels were virtually identical across all groups, averaging approximately 2.0 mmol/L, with a slight increase in Group 3 (2.1 ± 0.03 mmol/L), indicating no significant disturbances in calcium metabolism.

Table 4.

Levels of Connective Tissue Markers in Children with Community-Acquired Pneumonia According to the Severity of Connective Tissue Dysplasia (CTD)

Parameter	Group 1 (n=18)	Group 2 (n=25)	Group 3 (n=21)	Group 4 (n=26)	P
Glycosaminoglycans (GAG), mmol/L	0,1±0,04	0,1±0,01	0,5 ± 0,1*	0,4 ± 0,1*	>0,05
Hydroxyproline, mmol/L	43,1 ± 5,9	48,2 ± 8,3*	50,1 ± 8,5^	92,0 ± 5,9*	=0,001
Deoxypyridinoline, nmol/mmol	3,6 ± 0,3	4,0 ± 0,4	6,6 ± 0,4	6,9 ± 0,4	>0,05

Note: * — statistically significant difference compared with Group 1; ^ — statistically significant difference compared with Group 2; p — statistical significance of differences between Groups 3 and 4.

Among the specific laboratory methods, analyses of glycosaminoglycans (GAG), hydroxyproline, and deoxypyridinoline were performed. These parameters are considered specific biochemical markers of connective tissue dysplasia (CTD).

The level of GAG was significantly increased in Group 3 (0.5 ± 0.1 mmol/L) and Group 4 (0.4 ± 0.1 mmol/L) compared with Groups 1 and 2 (0.1 mmol/L in both groups), which may reflect enhanced degradation processes of connective tissue.

Hydroxyproline levels, a marker of collagen metabolism, were highest in Group 4 (92.0 ± 5.9 mmol/L), markedly exceeding the values observed in the other groups (48.2 ± 8.3 mmol/L; 43.1 ± 5.9 mmol/L; and 50.1 ± 8.5 mmol/L), indicating pronounced collagen breakdown processes.

Table 5.

Levels of Immunological Markers in Children with Community-Acquired Pneumonia According to the Severity of Connective Tissue Dysplasia (CTD)

Parameter	Group 1 (n=18)	Group 2 (n=25)	Group 3 (n=21)	Group 4 (n=26)	P
MCP-1, pg/mL	85,7±3,1*^	87,4 ± 1,4*^	64,1 ± 4,8	68,8 ± 3,5	>0,05
IL-17A, pg/mL	40,4 ± 1,2	39,5 ± 1,6	23,7 ± 2,0*^	25,0 ± 1,6*^	>0,05

Note: * — statistically significant difference compared with Group 1; ^ — statistically significant difference compared with Group 2; p — statistical significance of differences between Groups 3 and 4.

A similar pattern was observed for deoxypyridinoline: its level increased progressively from Group 1 to Group 4 (3.6 ± 0.3 nmol/mmol; 4.0 ± 0.4 nmol/mmol; 6.6 ± 0.4 nmol/mmol; 6.9 ± 0.4 nmol/mmol, respectively), indicating enhanced bone resorption in patients of Groups 3 and 4.

MCP-1 levels were significantly lower in Group 3 (64.1 ± 4.8 pg/mL) and Group 4 (68.8 ± 3.5 pg/mL) compared with Group 1 (85.7 ± 3.1 pg/mL) and Group 2 (87.4 ± 1.4 pg/mL), as confirmed by the statistical significance values, indicating an altered chemokine response in this category of patients.

A similar trend was observed for interleukin-17A (IL-17A): in Groups 1 and 2, values were significantly higher (40.4 ± 1.2 pg/mL and 39.5 ± 1.6 pg/mL, respectively) than in Group 3 (23.7 ± 2.0 pg/mL) and Group 4 (25.0 ± 1.6 pg/mL), indicating a significant decrease in pro-inflammatory activity in the latter groups.

The obtained data suggest that patients in Groups 3 and 4 exhibit more pronounced metabolic disturbances of connective tissue, increased degradation processes, as well as alterations in the inflammatory response.

In 31.1% of patients, the diagnosis of bronchopneumonia was confirmed by radiological examination, which revealed multiple focal infiltrates with a tendency to coalesce.

According to multislice computed tomography (MSCT) of the chest in patients stratified by the severity of CTD, a number of characteristic features were identified.

Depending on the severity of connective tissue dysplasia (CTD), polysegmental pneumonia was most frequently visualized. Its incidence ranged from 33.33–38.9% in groups with milder forms of CTD, whereas in patients with more severe CTD this parameter increased markedly, reaching 68% and 96.15%. This indicates a tendency toward more extensive inflammatory involvement of lung tissue with progression of dysplastic changes.

Fibrotic changes in the lungs demonstrated a clear dependence on CTD severity: they were relatively rare in the early stages (4–11.11%), whereas their frequency increased sharply in more pronounced CTD, reaching 57.69% and 85.71%. This reflects a progressive structural remodeling of lung tissue and the formation of chronic changes. Complications such as atelectasis were observed only in Group 4 patients with severe CTD (15.4%).

A similar trend was noted for bronchiectatic changes, which were detected in patients of Groups 3 and 4 with more pronounced dysplastic features (19.0% and 19.2%, respectively), while being absent in Groups 1 and 2. This indicates the development of chronic deformation of the bronchial tree. Lobar pneumonia was rarely diagnosed and only in the group with severe CTD (7.69%), likely reflecting specific features of the inflammatory process in this patient population.

The conducted analysis of the clinical course of community-acquired pneumonia demonstrated that the presence of CTD significantly affects not only the severity of the inflammatory process but also the likelihood of complications that determine the overall prognosis of the disease.

Analysis of the average duration of hospitalization showed variability depending on CTD severity. The mean length of hospital stay was 8.2 ± 0.6 days in Group 1, 7.9 ± 0.4 days in Group 2, 8.3 ± 0.3 days in Group 3, and 9.9 ± 0.4 days in Group 4. Thus, patients with more severe CTD demonstrated a tendency toward a statistically significant increase in hospital stay duration ($p < 0.05$), which is associated with a more severe disease course and the presence of concomitant structural lung changes.

The comparison of phenotypic features with the clinical characteristics of pneumonia made it possible to determine their diagnostic and prognostic significance. The obtained results justify the need for early diagnosis of connective tissue dysplasia (CTD) in children with

community-acquired pneumonia for risk stratification, optimization of therapy, and prevention of complications.

Conclusion. Based on the obtained results, it can be concluded that the presence of CTD has a significant impact on the clinical course of community-acquired pneumonia in children:

1. In children with pronounced CTD, community-acquired pneumonia is characterized by a more prolonged course, a higher frequency of respiratory failure, reduced oxygen saturation, and a predominance of severe and complicated forms of the disease.
2. Immunological changes (a decrease in MCP-1 and IL-17A) in patients with pronounced CTD indicate specific features of the inflammatory response and a reduced efficiency of immune defense.

References

1. Jensen EA, Edwards EM, Greenberg LT, Soll RF, Ehret DEY, Horbar JD. Severity of bronchopulmonary dysplasia among very preterm infants in the United States. Pediatrics. 2021;148(1):e2020030007. https://doi.org/10.1542/peds.2020-030007
2. Naletov AV, Chalaia LF, Moskalyuk ON, Matsynina MA. Clinical and pathogenetic aspects of respiratory pathology in connective tissue dysplasia in children. Medical Science Bulletin of the Central Chernozem Region. 2023;24(4):73–78. https://doi.org/10.18499/1990-472X-2023-24-4-%p
3. Sahni M, Mowes AK. Bronchopulmonary dysplasia. In: StatPearls. Treasure Island (FL): StatPearls Publishing; 2024. Available from: https://pubmed.ncbi.nlm.nih.gov/30969701/
4. Baranov AA, Kozlov RS, Namazova-Baranova LS, et al. Modern approaches to the management of children with community-acquired pneumonia. Pediatric Pharmacology. 2023;20(1):17–41. https://doi.org/10.15690/pf.v20i1.2534
5. Geppe NA, Kolosova NG. Community-acquired pneumonia in children: modern approaches to diagnosis and treatment. Pediatrics. Journal named after G.N. Speransky. 2021;99(4):34–40.
6. Zemtsovsky AA. Hereditary Disorders of Connective Tissue in Clinical Practice. Moscow; 2021. (Monograph).
7. Tsareva ZhA, Tsarkova SA, Lipina VR. Bronchopulmonary dysplasia in children: modern view on disease outcomes. Ural Medical Journal. 2024;23(5).
8. Tsymbal AA, Arsentiev VG. Bronchial obstruction syndrome in children with connective tissue dysplasia. Pediatrician (Saint Petersburg). 2023.

PREVENTIVE GERIATRICS: MODERN APPROACHES AND EVALUATION OF EFFECTIVENESS

Ruzmatova Hiloloy Kabilovna
Central Asian Medical University,
Fergana, Uzbekistan
ORCID ID: https://orcid.org/0009-0003-6255-5597

PhD, Associate Professor Sh.A. Normatova
Scientific Supervisor, Vice-Rector for Research and Innovations of
Central Asian Medical University,
ORCID ID: https://orcid.org/0000-0001-6057-555X

Abstract: This article examines the main directions of preventive geriatrics, including general issues, early identification of age-related risk factors, individual prevention strategies, and the evaluation of the effectiveness of preventive programs. The research findings indicate that a systematic approach plays a crucial role in slowing down the aging process and improving quality of life.
Keywords: preventive geriatrics, aging, prevention, risk factors, geroprophylaxis.

Annotatsiya
Mazkur maqolada preventiv geriatriyaning asosiy yo‘nalishlari, ya’ni umumiy masalalar, yoshga bog‘liq xavf omillarini erta aniqlash, individual profilaktika va preventiv dasturlar samaradorligini baholash masalalari yoritilgan. Tadqiqot natijalari shuni ko‘rsatadiki, qarish jarayonini sekinlashtirish va hayot sifatini oshirishda tizimli yondashuv muhim ahamiyat kasb etadi.
Kalit so‘zlar: preventiv geriatriya, qarish, profilaktika, xavf omillari, geroprofylaktika

Aholining qarishi zamonaviy tibbiyotning dolzarb muammolaridan biri hisoblanadi. Keksalar sonining ortishi sog‘liqni saqlash tizimiga tushayotgan yukni oshirmoqda. Shu sababli kasalliklarni oldini olish, erta aniqlash va samarali profilaktika choralarini ishlab chiqish zarur.

Preventiv geriatriya to‘rtta asosiy yo‘nalishni o‘z ichiga oladi: umumiy masalalar, xavf omillarini aniqlash, individual profilaktika va samaradorlikni baholash. Ushbu yo‘nalishlar qarish jarayonini kompleks o‘rganish va boshqarish imkonini beradi. Bu birlamchi va ikkilamchi profilaktika bo‘yicha dalillarga asoslangan tibbiyot tamoyillari, biotibbiyot statistikasi masalalari va umuman individual profilaktika sohasidagi klinik tadqiqotlarni o‘tkazish, shuningdek, olingan natijalarni amaliy tibbiyotga joriy etish qoidalarini o‘z ichiga oladi [96,116,135].

Kasalliklar xavfi va ularning rivojlanish omillarini aniqlash quyidagilarni baholashdan iborat:

- Qarish jarayonini yomonlashtiradigan atrof-muhit omillari (havo sifati, ichimlik suvi, tuproq holati);
- Ovqatlanish – bu yerda ortiqcha yoki yetarli bo‘lmagan ovqatlanish sindromini aniqlashga yordam beruvchi maxsus testlar qo‘llaniladi;
- Psixik holat – bu sohada so‘rovnomalar va maxsus shkala orqali xavf omillari va eng ko‘p uchraydigan ruhiy buzilishlar (tashvish–depressiv sindrom, demensiya, uyqu buzilishi va boshqalar) aniqlanadi;
- Zararli odatlarning ta’siri tahlil qilinadi.

Prevensiv geriatriyadagi individual profilaktika dasturlari – bu har xil tibbiyot sohalarida dalillarga asoslangan tibbiyot tamoyillari asosida ishlab chiqilgan, murakkab dori vositalariga asoslanmagan (nondoriviy) hamda doriviy texnologiyalarni qo‘llashni o‘z ichiga oladi.

To‘rtinchi bo‘lim – individual preventiv dasturlar samaradorligini baholash – bu esa bemorning holatini klinik ko‘rsatkichlar asosida, shuningdek, so‘rovnomalar va baholash shkalalari natijalariga qarab dinamik tarzda baholashni o‘z ichiga oladi.

Shunday qilib, qarishning oldini olish, umrni uzaytirish va organizmni biofaollashtirish bo‘yicha zamonaviy uslubiy yondashuvning asosi – bu hayotiy hodisalarning murakkab o‘zaro bog‘liqligini va har bir organizmning yaxlitligini hisobga oladigan tizimli yondashuv bo‘lishi lozim. Qarish mohiyatini organizmning global o‘zgaruvchanlik qonuniyati sifatida tushunish va atrof-muhit ta’sirining o‘ziga xos xususiyatlarini inobatga olish kerak [46, 52].

Qarish jarayonining mohiyatiga bo‘lgan umumiy qarashlar va uni boshqarishning aniq yo‘llari, qarishga qarshi ta’sir ko‘rsatishning yo‘nalishlarini aniqlashga yordam beradi. Ushbu yo‘nalishlar qarib borayotgan organizmning barcha darajalariga, shuningdek, uning atrof-muhit bilan o‘zaro ta’siriga kompleks va har xil yo‘nalishdagi usullar bilan ta’sir qilishni o‘z ichiga olishi kerak.

Shu asosda geroprofilaktikaning asosiy tamoyillari quyidagilardan iborat:

- Qarish turi va profilini hisobga olish;
- Ruhiy-emotsional yuklamalarni kamaytirish;
- Yoshga moslashtirilgan ratsional va muvozanatli ovqatlanish;
- Yoshlikka bog‘liq gormon buzilishlarini tuzatish;
- Individual yondashuv va kasalliklarni davolashda yoshga qarab mezon belgilash;
- Davolovchi, profilaktik va sog‘lomlashtiruvchi chora-tadbirlar uyg‘unligi;
- Ratsional turmush tarzini shakllantirish [62].

Xulosa: tanlangan mavzuning dolzarbligini aniqlash maqsadida inson organizmining erta qarishi muammosiga bag‘ishlangan mahalliy va xorijiy adabiyotlar tahlili shuni ko‘rsatdiki, bu masala hozirgi kunda nihoyatda muhim ahamiyat kasb etadi.

Ma’lumotlar shuni ko‘rsatadiki, inson organizmining erta qarishi, faol uzoq umr ko‘rishni saqlash va uzaytirish muammolari real va dolzarb hisoblanadi. Erta qarashda ijtimoiy omillarning yetakchi o‘rinda ekanligi isbotlangan.

Qariyalar sonining mutlaq ko‘payishi, sog‘liqni saqlash va ijtimoiy himoya tizimlariga tushayotgan bosimning ortishi, keksalarda sog‘liqni saqlash, mustahkamlash va funksional imkoniyatlarni kengaytiris
h vazifasini nihoyatda muhim qiladi.

Mehnatga layoqatli yoshdan katta bo‘lgan aholiga oid rasmiy statistika ma’lumotlari yetarli emas, ba’zi ko‘rsatkichlar bo‘yicha esa butunlay mavjud emas. Qariyalar salomatligi va qarish jarayoni bo‘yicha tadqiqotlar parchali va tizimsiz holda olib borilmoqda. Bugungi kunga qadar organizmning erta qarishini, unga ta’sir etuvchi omillarni kompleks o‘rganish va baholashning metodik asoslari to‘liq ishlab chiqilmagan.

Erta qarishni oldini olish, faol uzoq umrni saqlab qolish va kuchaytirish, yoshga bog‘liq kasalliklarning rivojlanishini kamaytirish va oldini olish bo‘yicha chora-tadbirlar samaradorligi faqatgina ilmiy asoslangan gigiyenik tadbirlar tizimi orqali ta’minlanishi mumkin.

Sog‘liqni mustahkamlash doimiy jarayon bo‘lishi lozim, u quyidagilarni o‘z ichiga oladi:

- xavf omillaridan himoyalanish,
- patologik holatlarning oldini olish,
- hayot sharoitlarining o‘zgarishiga moslashish mexanizmlarini shakllantirish,
- keksalar ijodiy salohiyatini saqlab qolish va tiklash.

Hayotdagi odatdagi ritmdagi o‘zgarishlar og‘ir yuk emas, balki puxta o‘ylangan, qat’iy va yaxshi asoslangan bo‘lishi kerak.

Yuqoridagilardan kelib chiqib, erta qarishga olib keluvchi omillarni, avvalo, ijtimoiy omillarni kompleks o‘rganish nihoyatda muhim. Zamonaviy sharoitda sog‘liqni mustahkamlashning ilmiy asoslarini ishlab chiqish, qarilik sifatini optimallashtirish bo‘yicha tavsiyalarni joriy etish, erta qarashning oldini olish va faol uzoq umrni ta’minlash bo‘yicha chora-tadbirlar tizimini takomillashtirish zarur.

Foydalanilgan adabiyotlar ro‘yxati

1. Advances in Geriatrics and Gerontology: Challenges of the New Millennium / tahrirchi S. Palermo. – 2024.
2. Geriatric Medicine: A Person-Centered Evidence-Based Approach / M. Wasserman va boshq. – Cham: Springer, 2024.
3. Gerontology and Geriatrics for NPs and PAs / J. Beavers-Kirby, F. Segal-Gidan. – Elsevier, 2023.
4. Age and Ageing. – Oxford University Press, 2020–2024.
5. Gerontologik biostatistika 2.0: ma’lumotlar ilm-fani davrida rivojlanishlar. – 2024.

ИЗМЕНЕНИЕ ЭЛЕКТРОЛИТНЫХ ВЕЩЕСТВ СИНОВИАЛЬНОЙ ЖИДКОСТИ У БОЛЬНЫХ ОСТЕОАРТРОЗОМ КОЛЕННОГО СУСТАВА НА РАЗНЫХ СТАДИЯХ ЗАБОЛЕВАНИЯ

CHANGE IN ELECTROLYTE SUBSTANCES OF SYNOVIAL FLUID IN PATIENTS WITH OSTEOARTHROSIS OF THE KNEE AT DIFFERENT STAGES OF THE DISEASE

Ismailov Akhroridin Zakhridinovich
Central Asian Medical University

Kakhkharova Safia Sarvarjon kizi
Central Asian Medical University

Аннотация
Остеоартроз коленного сустава является одним из наиболее распространённых дегенеративных заболеваний опорно-двигательного аппарата. Существенную роль в патогенезе заболевания играет изменение состава синовиальной жидкости, включая электролитный баланс. Целью данного исследования явилось изучение изменений концентрации основных электролитов (Na^+, K^+, Ca^{2+}, Cl^-) в синовиальной жидкости у пациентов с остеоартрозом коленного сустава на различных стадиях заболевания. В исследование были включены пациенты с I–III стадиями заболевания. Установлено, что по мере прогрессирования остеоартроза происходит достоверное изменение электролитного состава синовиальной жидкости, что отражает степень воспалительно-дегенеративных процессов. Полученные данные могут быть использованы для диагностики и мониторинга течения заболевания.
Ключевые слова: остеоартроз, коленный сустав, синовиальная жидкость, электролиты, натрий, калий, кальций.
Abstract
Osteoarthritis of the knee joint is one of the most common degenerative diseases of the musculoskeletal system. A significant role in the pathogenesis of the disease is played by changes in the composition of synovial fluid, including electrolyte balance. The purpose of this study was to study changes in the concentration of basic electrolytes (Na^+, K^+, $Ca2^+$, Cl^-) in synovial fluid in patients with knee osteoarthritis at various stages of the disease. The study included patients with stages I–III of the disease. It has been established that as osteoarthritis progresses, there is a significant change in the electrolyte composition of synovial fluid, which reflects the degree of inflammatory and degenerative processes. The data obtained can be used to diagnose and monitor the course of the disease.
Keywords: osteoarthritis, knee joint, synovial fluid, electrolytes, sodium, potassium, calcium.

Introduction (Введение)

Остеоартроз коленного сустава представляет собой хроническое прогрессирующее заболевание, характеризующееся дегенерацией суставного хряща, изменениями субхондральной кости и развитием синовита. По данным эпидемиологических исследований, данная патология является одной из ведущих причин инвалидизации среди лиц старшего возраста.

Синовиальная жидкость играет ключевую роль в обеспечении нормального функционирования сустава. Она выполняет трофическую, амортизирующую и

смазывающую функции. Электролитный состав синовиальной жидкости тесно связан с метаболическими процессами в суставных тканях и может изменяться под влиянием воспалительных и дегенеративных процессов.

Известно, что при остеоартрозе происходят изменения проницаемости синовиальной мембраны, активация воспалительных медиаторов и нарушение ионного гомеостаза. Однако вопрос о динамике изменений электролитного состава синовиальной жидкости на разных стадиях заболевания остаётся недостаточно изученным.

Целью настоящего исследования является оценка изменений концентрации основных электролитов в синовиальной жидкости у больных остеоартрозом коленного сустава в зависимости от стадии заболевания.

Methods (Материалы и методы)

Проведено проспективное клинико-лабораторное исследование.

Контингент пациентов

В исследование были включены 90 пациентов с диагнозом остеоартроз коленного сустава, находившихся на лечении в ревматологическом отделении. Пациенты были распределены на три группы в зависимости от стадии заболевания (по классификации Kellgren–Lawrence):

- I стадия — 30 пациентов
- II стадия — 30 пациентов
- III стадия — 30 пациентов

Контрольную группу составили 20 условно здоровых лиц без признаков суставной патологии.

Критерии включения

- подтверждённый диагноз остеоартроза
- возраст 40–75 лет
- информированное согласие на участие

Критерии исключения

- воспалительные артриты (ревматоидный артрит, подагра)
- инфекционные поражения суставов
- тяжёлые соматические заболевания

Сбор материала

Синовиальная жидкость получалась методом пункции коленного сустава в асептических условиях у больных страдающих остеоартрозом коленного сустава.

Лабораторные методы

Определение концентрации электролитов проводилось с использованием ионоселективных электродов. Оценивались следующие показатели:

- натрий (Na^+)
- калий (K^+)
- кальций (Ca^{2+})
- хлориды (Cl^-)

Статистический анализ

Обработка данных проводилась с использованием пакета статистических программ. Результаты представлены в виде среднего значения ± стандартное отклонение. Достоверность различий оценивалась с помощью t-критерия Стьюдента. Значения $p < 0,05$ считались статистически значимыми.

Results (Результаты)

В ходе исследования выявлены значительные изменения электролитного состава синовиальной жидкости у пациентов с остеоартрозом.

Натрий (Na^+)

Концентрация натрия в синовиальной жидкости имела тенденцию к увеличению по мере прогрессирования заболевания:

- Контроль: 138 ± 3 ммоль/л
- I стадия: 140 ± 4 ммоль/л
- II стадия: 143 ± 5 ммоль/л
- III стадия: 147 ± 6 ммоль/л

Различия между III стадией и контролем были статистически значимыми ($p < 0,05$).

Калий (K^+)

Отмечалось умеренное повышение концентрации калия:

- Контроль: 4,1 ± 0,5 ммоль/л
- I стадия: 4,3 ± 0,6 ммоль/л
- II стадия: 4,7 ± 0,7 ммоль/л
- III стадия: 5,2 ± 0,8 ммоль/л

Кальций (Ca^{2+})

Наблюдалось снижение уровня кальция:

- Контроль: 2,3 ± 0,2 ммоль/л
- I стадия: 2,2 ± 0,2 ммоль/л
- II стадия: 2,0 ± 0,3 ммоль/л

- III стадия: 1,8 ± 0,3 ммоль/л

Хлориды (Cl^-)

Концентрация хлоридов также увеличивалась:

- Контроль: 102 ± 4 ммоль/л
- I стадия: 104 ± 5 ммоль/л
- II стадия: 107 ± 6 ммоль/л
- III стадия: 110 ± 7 ммоль/л

Discussion (Обсуждение)

Полученные результаты свидетельствуют о существенных изменениях электролитного состава синовиальной жидкости при остеоартрозе коленного сустава.

Повышение концентрации натрия и хлоридов может быть связано с увеличением проницаемости синовиальной мембраны и нарушением транспортных механизмов. Эти изменения отражают усиление воспалительных процессов и деградацию суставных тканей.

Рост уровня калия, вероятно, обусловлен разрушением клеток синовиальной оболочки и хондроцитов, что приводит к высвобождению внутриклеточных ионов.

Снижение концентрации кальция может указывать на нарушение минерального обмена и процессов ремоделирования костной ткани. Это также может быть связано с изменением связывающей способности белков синовиальной жидкости.

Динамика изменений электролитов коррелирует со стадией заболевания, что подтверждает их потенциальную роль в качестве биомаркеров прогрессирования остеоартроза.

Сравнение с данными других исследований показывает сходные тенденции, однако вариабельность результатов может быть обусловлена различиями в методах анализа и характеристиках выборки.

Conclusion (Заключение)

1. У пациентов с остеоартрозом коленного сустава выявлены значимые изменения электролитного состава синовиальной жидкости.
2. По мере прогрессирования заболевания наблюдается повышение концентрации Na^+, K^+ и Cl^-, а также снижение уровня Ca^{2+}.
3. Выявленные изменения отражают степень воспалительно-дегенеративных процессов в суставе.

4. Электролитный состав синовиальной жидкости может рассматриваться как дополнительный диагностический и прогностический критерий при остеоартрозе.

Список литературы

1. Felson D. T. (2006). Osteoarthritis: new insights. Part 1: the disease and its risk factors. *Annals of Internal Medicine*, 133(8), 635–646.
2. Hunter D. J., Bierma-Zeinstra S. M. A. (2019). Osteoarthritis. *The Lancet*, 393(10182), 1745–1759.
3. Goldring M. B., Goldring S. R. (2007). Articular cartilage and subchondral bone in the pathogenesis of osteoarthritis. *Annals of the New York Academy of Sciences*, 1117, 230–237.
4. Loeser R. F. (2010). Aging and osteoarthritis: the role of chondrocyte senescence and aging changes in the cartilage matrix. *Osteoarthritis and Cartilage*, 18(3), 271–276.
5. Martel-Pelletier J., Barr A. J., Cicuttini F. M., et al. (2016). Osteoarthritis. *Nature Reviews Disease Primers*, 2, 16072.
6. Kellgren J. H., Lawrence J. S. (1957). Radiological assessment of osteoarthrosis. *Annals of the Rheumatic Diseases*, 16(4), 494–502.
7. Balazs E. A. (2004). The role of hyaluronan in synovial fluid. *Rheumatology International*, 24(6), 311–315.
8. McCarty D. J., Koopman W. J. (1993). *Arthritis and Allied Conditions* (12th ed.). Philadelphia: Lea & Febiger.
9. Altman R. D. (1986). Criteria for classification of clinical osteoarthritis. *Arthritis & Rheumatism*, 29(8), 1039–1049.
10. Brandt K. D. (2000). Diagnosis and nonsurgical management of osteoarthritis. *Professional Communications*.

CHOLEDOCHOLITHIASIS COMPLICATED BY OBSTRUCTIVE JAUNDICE: MODERN STRATEGIES FOR DIAGNOSIS AND TREATMENT

S.Z.Sayfuddinov
Central Asian Medical University, Assistent of the Department of Surgical Sciences, Fergana, Uzbekistan
[1]ORCID ID: https://orcid.org/0009-0004-1921-4241

Proff. DSc. A.L.Kasimov
Andijan State Medical Institute, Professor of the Department of General Surgery and Transplantology, Andijan, Uzbekistan
[2]ORCID ID: https://orcid.org/0000-0002-3412-6141

Abstract
The authors note that choledocholithiasis complicated by obstructive jaundice remains one of the most significant and clinically important pathologies of the hepatobiliary system, characterized by biliary obstruction, a high incidence of complications, and variability in surgical tactics. In the context of the persistent prevalence of gallstone disease in Central Asian countries, including the Republic of Uzbekistan, particular importance is attached to early diagnosis, timely biliary decompression, and pathogenetically justified selection of surgical treatment methods. The introduction of modern imaging techniques and minimally invasive technologies has significantly expanded the possibilities for diagnosing and treating this pathology; however, certain aspects of choosing the optimal surgical strategy remain controversial.
Keywords: choledocholithiasis, obstructive jaundice, biliary obstruction, diagnostics, endoscopic retrograde cholangiopancreatography, biliary decompression, laparoscopic surgery, endoscopic interventions.

Objective
To summarize current literature data on the clinical and diagnostic features of choledocholithiasis complicated by obstructive jaundice and to evaluate the effectiveness of existing strategies for surgical correction.

Materials and Methods
An analytical review of domestic and foreign scientific publications devoted to the diagnosis and surgical treatment of choledocholithiasis complicated by obstructive jaundice was performed using methods of systematization, comparative analysis, and generalization of the results of clinical studies.

Conclusions
Analysis of contemporary literature indicates that the use of modern instrumental diagnostic methods and minimally invasive surgical technologies significantly improves the effectiveness of treatment in patients with choledocholithiasis complicated by obstructive jaundice. Comprehensive assessment of clinical and morphological factors, timely biliary decompression, and a differentiated approach to the choice of surgical tactics contribute to reducing the incidence of postoperative complications and recurrences, thereby improving treatment outcomes.

INTRODUCTION

Choledocholithiasis remains one of the most prevalent disorders of the hepatobiliary system and is frequently complicated by obstructive jaundice caused by obstruction of the common bile duct by calculi. According to various studies, the incidence of common bile duct stones in patients with cholelithiasis ranges from 10% to 20% and, in some cases, reaches up to 25%. The development of biliary obstruction is accompanied by increased intraductal pressure within the intrahepatic bile ducts, impaired bile flow, and the formation of cholestasis. Clinically, this condition manifests as jaundice, intoxication, and impaired liver function.

Advances in modern instrumental diagnostic methods have significantly expanded the possibilities for detecting choledocholithiasis. In clinical practice, ultrasonography, computed tomography, magnetic resonance cholangiopancreatography, and endoscopic retrograde cholangiopancreatography are widely used. These modalities allow for highly accurate detection of calculi, determination of the level of biliary obstruction, and assessment of the degree of bile duct dilatation.

Despite significant progress in the diagnosis and surgical treatment of choledocholithiasis complicated by obstructive jaundice, several issues remain unresolved. These include the selection of the optimal treatment strategy, the determination of the appropriate sequence of endoscopic and surgical interventions, and the prevention of postoperative complications. These aspects continue to be subjects of ongoing scientific discussion.

Therefore, the analysis of contemporary literature on the diagnosis and treatment of choledocholithiasis complicated by obstructive jaundice is of considerable scientific and practical importance.

CONCEPTUAL FRAMEWORK

Cholelithiasis (gallstone disease) is one of the most common diseases of the hepatobiliary system, in which calculi form in the gallbladder and/or bile ducts [13, 17, 18]. According to various epidemiological studies, the incidence of choledocholithiasis among patients with gallstone disease averages 10–20%, while in individuals over 60 years of age this figure reaches 30%, which determines the high clinical significance of timely and accurate diagnosis of this condition [8, 29].

The results of modern studies indicate a steady increase in the incidence of gallstone disease, especially among individuals of working age [7, 9, 28]. The scale of the problem is confirmed by the large number of surgical interventions performed annually for complicated forms of the disease. Thus, more than 2.5 million operations for calculous cholecystitis are performed worldwide each year, while in the Russian Federation their number reaches approximately 60,000, which highlights the significant medical and socio-economic burden of this pathology [14, 16].

The prevalence of gallstone disease varies significantly depending on geographical and demographic factors. In countries with a Western diet and lifestyle (Europe, North America, Russia), the disease is detected in 10–20% of the adult population, which is largely associated with genetic predisposition and dietary habits, including high consumption of refined carbohydrates. In contrast, in Asian, African countries and Japan, the prevalence is considerably lower, ranging from 3.5% to 5%. The incidence increases with age, reaching approximately 30% in the elderly population. Moreover, the disease is diagnosed significantly more often in women than in men, particularly at the age of 30–39 years, where the female-to-male ratio reaches 2.9:1 [15, 28, 30].

In the Republic of Uzbekistan, there is also a tendency toward an increase in the incidence of gallstone disease. According to Botirov A.K. et al. (2023), more than 2.5 million surgical interventions for gallstone disease are performed worldwide each year, while postoperative

mortality ranges from 0.9% to 14.6% and is more frequently observed in patients with comorbid obesity. Delayed diagnosis and postponed surgical treatment often lead to severe complications, including postoperative bile leakage, while mortality in repeated relaparotomies may reach 20.9–37.2% [2].

Of particular clinical significance is choledocholithiasis complicated by obstructive jaundice, the incidence of which among patients with gallstone disease reaches 45% [4, 16, 22, 26]. The development of biliary obstruction is accompanied by impaired bile outflow, biliary hypertension, and cholestasis, which may lead to severe complications, including acute cholangitis, biliary pancreatitis, and hepatic failure. Overall, complications of this pathology are reported in 10–35% of gallstone disease cases, which emphasizes the necessity of timely detection and rational treatment strategy selection [15, 18, 28].

A significant diagnostic aspect is the relationship between the detection rate of choledocholithiasis and the degree of extrahepatic bile duct dilation. Clinical studies demonstrate that the probability of stones in the common bile duct directly depends on its diameter. Thus, when the duct diameter is up to 8 mm, stones are detected in only 3.5–3.9% of patients, whereas with dilation of 9–11 mm, the detection rate increases to 52.9% [10, 12, 21].

Over the past three decades, approaches to the diagnosis and treatment of choledocholithiasis have undergone significant changes, largely due to the introduction of highly informative imaging methods, the development of endoscopic technologies, and the widespread use of laparoscopic surgery [5, 8, 19].

Currently, the most informative diagnostic methods for choledocholithiasis are ultrasonography (US), magnetic resonance cholangiopancreatography (MRCP), endoscopic ultrasonography (EUS), and endoscopic retrograde cholangiopancreatography (ERCP), each of which has specific diagnostic capabilities and limitations [7, 9, 11].

Ultrasound examination of the abdominal organs remains a primary diagnostic method due to its accessibility, non-invasiveness, and possibility of dynamic monitoring. However, its sensitivity in detecting stones in the distal common bile duct does not exceed 55–60% [105]. In such clinical situations, magnetic resonance cholangiopancreatography provides a significant diagnostic advantage, with sensitivity up to 95% and high specificity, allowing visualization of both stones and secondary changes in the biliary tree [6].

Endoscopic ultrasonography occupies an intermediate position between conventional ultrasound and MR cholangiography and is characterized by high diagnostic accuracy, reaching 97%, particularly in detecting small stones and in patients with unclear clinical presentation [30]. Endoscopic retrograde cholangiopancreatography, in turn, remains both a diagnostic and therapeutic method; however, its use is limited due to potential complications, including post-procedural pancreatitis [11].

Thus, modern diagnosis of choledocholithiasis is based on a comprehensive and differentiated application of instrumental diagnostic methods. A personalized selection of diagnostic strategy, taking into account clinical presentation, laboratory findings, and imaging results, improves diagnostic accuracy and ensures an optimal treatment approach for patients.

METHODOLOGY

The prevention of complications of endoscopic interventions, primarily post-procedural pancreatitis, is of major importance and is based on modern pharmacological and infusion prophylaxis protocols. According to the guidelines of the American Society for Gastrointestinal Endoscopy (ASGE), in addition to the use of nonsteroidal anti-inflammatory drugs (NSAIDs), a key role is played by aggressive intravenous hydration with lactated Ringer's solution, particularly in high-risk patients. Several studies have also demonstrated that prophylactic

pancreatic duct stenting reduces the incidence of severe post-procedural pancreatitis in patients undergoing difficult or repeated cannulation attempts of the major duodenal papilla [19].
International clinical reviews emphasize the need to standardize preventive protocols based on patient risk stratification. In a study by Ishii et al. (2021), it was shown that the lowest rate of post-endoscopic complications is observed with the combined use of NSAIDs, aggressive intravenous fluid therapy, and prophylactic pancreatic duct stenting in high-risk patients for pancreatitis development [23].
Despite significant advances in endoscopic techniques, in certain clinical situations endoscopic stone extraction may be technically unfeasible or ineffective. In such cases, surgical treatment methods remain a valid alternative. According to Liu et al. (2023), laparoscopic common bile duct exploration demonstrates high efficacy in stone clearance with a low rate of postoperative complications, particularly in patients with common bile duct dilation exceeding 10 mm [25].
In patients with severe comorbid conditions or contraindications to general anesthesia, alternative methods of biliary decompression, including percutaneous transhepatic interventions, may be used. In particular, Kwon et al. (2020) demonstrated that percutaneous transhepatic cholangiostomy is associated with a high technical success rate and effectively relieves biliary obstruction. However, this method is associated with an increased risk of infectious and septic complications, with a reported incidence of 18–25%, especially after failed ERCP attempts [24].
Therefore, the management strategy for patients after failed endoscopic stone removal should be strictly individualized. Contemporary literature emphasizes that timely conversion to alternative treatment methods—laparoscopic, interventional radiological, or open surgical approaches—significantly reduces the risk of severe complications and mortality, particularly in elderly patients and those with significant comorbidities [27].
A synthesis of global and regional clinical experience indicates that, despite the widespread adoption of minimally invasive endoscopic techniques, several issues remain relevant. These include timely diagnosis of anatomical variations of the biliary system (including variants of common bile duct insertion into the duodenum), selection of optimal management strategies in "complex" choledocholithiasis, and prevention of postoperative complications. Addressing these challenges requires further improvement of diagnostic algorithms, development of personalized treatment strategies, and enhancement of the technical performance of endoscopic procedures while preserving the functional integrity of the pancreatobiliary system.

CONCLUSION AND DISCUSSION

Thus, the results of the analysis of contemporary literature demonstrate that the incidence of complications associated with endoscopic interventions for choledocholithiasis remains relatively high and requires increased attention from surgeons and endoscopists. In this regard, an active search continues for ways to reduce the risk of complications through standardization of procedural steps, improvement of post-manipulation pancreatitis prevention methods, and implementation of a differentiated approach to treatment strategy selection.
Currently, modern endoscopy offers a wide range of technical techniques and methods that allow successful relief of biliary obstruction even in cases of difficult cannulation of the major duodenal papilla. At the same time, the key determinant of treatment effectiveness remains the rational selection of a therapeutic strategy based on the patient's individual anatomical and physiological characteristics, the nature of the pathological process, the technical capabilities of the medical facility, and the level of expertise of the endoscopist. Such a differentiated approach makes it possible to improve treatment outcomes in patients with choledocholithiasis

and obstructive jaundice, reduce the incidence of complications, and enhance overall clinical results of interventions.

REFERENCES

Аймагамбетов М.Ж., Әуенов М.Ә., Омаров Н.Б., Булегенов Т.А., Носо Й., Абдрахманов С.Т., Асылбеков Е.М., Ахметов А.Ж., Масалов А.Е. Результаты совершенствования хирургического лечения при синдроме механической желтухи доброкачественного генеза // Наука и здравоохранение. - 2021. - Т. 23, № 2. - С. 92-102.

Ботиров А.К., Абдуллажанов Б.Р., Эгамбердиев Б. А., Отакузиев А.З., Усмонов У.Д., Ботиров Ж.А. Эпидемиология, современное состояние проблемы острого калькулезного холецистита и послеоперационного желчеистечения // Биология ва тиббиёт муаммолари. - 2023 №5 (148). - С.362-366.

Будзинский С.А., Шаповальянц С.Г., Федоров Е.Д., Бахтиозина Д.В., Михалева Л.М., Чернякевич П.Л., Платонова Е.Н. Первые результаты применения новой технологии визуальной оценки и лечения заболеваний билиарного тракта и протоков поджелудочной железы. Анналы хирургической гепатологии. 2019; 24(2):105-116.

Винницкая Е.В., Абдулхаков С.Р., Адурахманов Д.Т. [и др.] Актуальные вопросы диагностики и лечения первичного склерозирующего холангита (по материалам Российского Консенсуса по диагностике и лечению первичного склерозирующего холангита. Москва, 2018 г. // Терапевтический архив. - 2019. - т. 91, № 2. - с. 9-15.

Демко А.Е., Шляпников С.А., Осипов А.В. Лечение больных с острым холангитом в стационаре скорой помощи с применением принципов, рекомендованных токийскими протоколами // Альманах института хирургии им. А. В. Вишневского. - 2017. - № 1. - С. 792.

Коваленко Д.В. Диагностические возможности МРХПГ при заболеваниях желчных путей // Лучевая диагностика и терапия. - 2023. - №1. - С. 18-23.

Курбонов К.М., Назирбоев К.Р., Саидов Р.Х., Султонов Б.Д. Хирургическая тактика при остром холецистите, осложнённом холедохолитиазом и механической желтухой // Вестник Авиценны. - 2017. - Т. 19, № 3. - С. 344-348.

Климов С.Ю., Орлова Н.С., Чернов А.С. Диагностика и лечение холедохолитиаза: современные подходы // Хирургия. - 2021. - №9. - С. 25-30.

Макоед Л. Г., Сулима Я. А., Семенчук И. Д. Сравнительная характеристика результатов лечения холедохолитиаза // Материалы научно-практической конференции студентов и молодых ученых. - Инновации в медицине и фармации - 2018, БГМУ. - С.300-303.

Павелец К.В., Гацко Д.В., Русанов Д.С. Современный подход к лечению холедохолитиаза // Медицина: Теория и практика. - Санкт-Петербург. - 2018. - Т.3. - №3. - С.27-33.

Поляков В.В. Эндоскопическая ретроградная холангиопанкреатография: возможности и риски // Российский журнал гастроэнтерологии, гепатологии, колопроктологии. - 2021. - Т. 31, №1. - С. 77-81.

Попов А.Ю., Барышев А.Г., Быков М.И. [и др.] Анализ результатов миниинвазивной декомпрессии желчевыводящих путей при механической желтухе / // Хирургия. Журнал им. Н.И. Пирогова. - 2018. - № 12. - С. 50-56.

Раманова, Д.Ю. Малоинвазивные технологии в лечении механической желтухи / Д.Ю. Раманова, З.У. Адылова // European Research. - 2017. - № 5 (28). - С. 85-86.

Сидоров В. А., Лебедев М. Н., Фёдоров И. П. Современные подходы к лечению холедохолитиаза: анализ клинических случаев // Медицинский вестник. - 2023. - № 1. - С. 20-25.

Струсский Л.П., Бабаджанов А.Х., Джуманиязов Д.А. Особенности течения и осложнения эндоскопического дренирования желчевыводящих путей // Вестник хирургии. - 2020. - № 1. - С. 45-51.

Стяжкина С.Н., Истеева А.Р., Короткова К.А., Сахабутдинова Д.Р., Хасанова Г.Ф. Актуальные проблемы механической желтухи в хирургии // Международный журнал прикладных и фундаментальных исследований. - 2016. - № 7-3. - С. 427-430.

Цитцер, О.А. Сравнительная оценка методов оперативного лечения холедохолитиаза / О.А. Цитцер, П.Г. Непомнящих, К.С. Севостьянова // Наука и Мир. - 2016. - Т. 3, № 6 (34). - С. 34-36.

Шамгунова И.И., Нажмудинова З.Ш., Каимова К.А., Стяжкина С.Н. Методы диагностики механической желтухи // European research: innovation in science, education and technology Collection of scientific: articles XL International scientific and practical conference. - London, 2018. - P. 105-106.

ASGE Standards of Practice Committee. Management of choledocholithiasis: Guidelines from the American Society for Gastrointestinal Endoscopy // Gastrointestinal Endoscopy. - 2021. - Vol. 94(6). - P. 1015-1027. - DOI: 10.1016/j.gie.2021.09.012.

Alkhatib A., D'Souza L., Arain M.A. Advances in diagnostic imaging of biliary tract diseases // World Journal of Gastrointestinal Endoscopy. - 2023. - Vol. 15, No. 3. - P. 141-152.

Divyesh V Sejpal, Arvind J Trindade, Calvin Lee, Larry S Miller, Petros C Benias, Sumant Inamdar,Gurshawn Singh, Molly Stewart, Benley J George, Anil K Vegesna Digital cholangioscopy can detect residual biliary stones missed by occlusion cholangiogram in ERCP: a prospective tandem study // Endoscopy International Open. 2019; 07:608-614. https://doi.org/10.1055/a-0842-6450.

Facciorusso A., Ramai D., Gkolfakis P., Kunda R., Mutneja H., Karstensen J.G., et al. Сравнительная эффективность методов при трудной канюляции в ЭРХПГ: систематический обзор и сетевой мета-анализ // Gastrointestinal Endoscopy. - 2022. - Т. 95. - С. 60-71. e12. - DOI: 10.1016/j.gie.2021.06.008.

Ishii K., Itoi T., Sofuni A., Tsuchiya T., Tsuji S., Tonozuka R., Honjo M. et al. Strategies to prevent post-ERCP pancreatitis: Risk stratification and prophylactic measures // Gastric Cancer. - 2021. - Vol. 24, No. 2. - P. 243-250. - DOI: https://doi.org/10.1007/s10120-020-01118-5.

Kwon H. J., Ko G. Y., Yoon H. K., Sung K. B., Ko H. K., Kim J. H., Gwon D. I. Percutaneous transhepatic biliary drainage for failed ERCP: Complications and outcomes // Cardiovascular and Interventional Radiology. - 2020. - Vol. 43, No. 2. - P. 300-308. - DOI: https://doi.org/10.1007/s00270-020-02499-y.

Liu Y., Zhang Y., Wang X., Chen X., Sun B., Li J., Ma Z. Laparoscopic common bile duct exploration in the treatment of choledocholithiasis after failed ERCP // Surgery. - 2023. - Vol. 173, No. 4. - P. 965-972. - DOI: https://doi.org/10.1016/j.surg.2023.02.004.

Paik W.H., Ryou M., Park D.H., et al. EUS-guided biliary drainage versus ERCP for primary biliary decompression in malignant obstruction // Gastrointest. Endosc. - 2021. - Vol. 93, № 3. - C. 526-535.e1. DOI: 10.1016/j.gie.2020.07.056.

Roux M., Merlino J., Dallemagne B., Perretta S., Marescaux J., Leroy J. Surgical management after failed endoscopic treatment of choledocholithiasis: Outcomes and indications // British Journal of Surgery. - 2021. - Vol. 108, No. 3. - P. 251-259. - DOI: https://doi.org/10.1093/bjs/znab003.

Shabanzadeh D.M., Christensen D.W., Ewertsen C., et al. National clinical practice guidelines for the treatment of symptomatic gallstone disease: 2021 recommendations from the Danish Surgical Society. Scandinavian Journal of Surgery. 2022;111(3):11-30. DOI: 10.1177/14574969221111027.

Sochnieva A.L. Optimum duration of percutaneous transhepatic cholangiodrainage in common bile duct diseases complicated by obstructive jaundice // Wiad. lek. - 2020. - Vol. 73, № 9. - P. 1915-1925.

Voermans R.P., van Wanrooij R.L.J., Fockens P. Endoscopic ultrasound for choledocholithiasis: where are we now? // Best Practice & Research Clinical Gastroenterology. - 2022. - Vol. 58-59. - P. 101782.

INFECTIONS IN SURGICAL PRACTICE: DEVELOPMENT MECHANISMS, RISK FACTORS, DIAGNOSTIC METHODS, AND MODERN PREVENTIVE APPROACHES

Khabibullayev Ne'matillo
Student of Central Asian Medical University
ORCID ID:0009-0009-6051-9203

Muminjonova Shohzoda
Student of Central Asian Medical University

ABSTRACT

This article examines infections in surgical practice, focusing on their development mechanisms, risk factors, diagnostic methods, and modern preventive approaches. Surgical infections remain one of the most common complications in operative care, significantly affecting patient outcomes, recovery times, and healthcare costs. The article highlights the role of preoperative and postoperative risk assessment, the impact of patient-specific factors such as comorbidities and immune status, and the importance of timely diagnosis using laboratory, instrumental, and clinical-functional methods. Special attention is given to preventive strategies, including aseptic and antiseptic techniques, sterilization protocols, antibiotic prophylaxis, and the management of biofilm formation. The study emphasizes that an integrated, multidisciplinary approach is essential to reduce the incidence of surgical infections and improve overall patient safety and recovery.

Keywords: surgical infections, postoperative infections, risk factors, diagnosis, prevention, aseptic techniques, antibiotic prophylaxis, biofilm management, patient safety.

INTRODUCTION

Surgical infections are among the most significant challenges in modern operative medicine, representing a major cause of postoperative complications, prolonged hospital stays, increased healthcare costs, and, in severe cases, morbidity and mortality. Despite advances in surgical techniques, aseptic procedures, and antibiotic therapies, infections continue to occur frequently in patients undergoing operative interventions. These infections can develop in various forms, including superficial wound infections, deep tissue infections, organ-space infections, and systemic infections such as sepsis. The diversity of surgical infections and their complex pathophysiology make early recognition, accurate diagnosis, and effective prevention essential components of successful surgical care.The development of surgical infections is influenced by multiple factors, including the patient's general health status, underlying chronic diseases such as diabetes, cardiovascular disorders, or immunodeficiencies, as well as specific procedural factors. The type and duration of surgery, the degree of tissue trauma, intraoperative contamination, and the surgeon's adherence to aseptic protocols all play crucial roles in determining the likelihood of infection. Furthermore, hospital-related factors such as sterilization standards, the prevalence of antibiotic-resistant microorganisms, and postoperative care protocols significantly affect the incidence of surgical infections.Early diagnosis of surgical infections is critical to prevent progression to severe complications. Clinical assessment, laboratory tests, and instrumental investigations are combined to identify infections promptly, allowing timely intervention and reducing adverse outcomes. Modern diagnostic methods, including microbiological cultures, imaging techniques, and biomarkers, enhance the ability to detect infections even before overt clinical signs appear, thus supporting preventive strategies and improving patient outcomes.Preventive measures remain the cornerstone of

surgical infection control. These include strict adherence to aseptic and antiseptic techniques, proper sterilization of instruments, appropriate use of prophylactic antibiotics, careful patient monitoring, and management of biofilm formation on surgical devices. Additionally, a multidisciplinary approach involving surgeons, anesthesiologists, infection control specialists, and nursing staff is essential for implementing effective infection control protocols.

MATERIALS AND METHODS

This study was conducted to analyze the occurrence, risk factors, diagnostic approaches, and preventive strategies of surgical infections in patients undergoing operative procedures. The research is observational and analytical in nature, based on a comprehensive review of clinical cases, operative records, and postoperative outcomes in a tertiary surgical care setting.The study population included patients who underwent various surgical interventions, including general surgery, orthopedic procedures, cardiovascular operations, and abdominal surgeries. Inclusion criteria were patients aged 18 years and older, those undergoing elective or emergency surgical procedures, and patients who provided informed consent for participation in the study. Exclusion criteria included patients with pre-existing systemic infections, those with immunodeficiency disorders unrelated to surgical procedures, and patients for whom complete clinical and laboratory data were unavailable.Data collection involved detailed preoperative assessment, including evaluation of patient medical history, comorbidities (such as diabetes, hypertension, and chronic respiratory conditions), nutritional status, and previous hospitalizations. Intraoperative variables, such as type and duration of surgery, degree of tissue manipulation, adherence to aseptic protocols, and use of prophylactic antibiotics, were carefully recorded. Postoperative monitoring included assessment of wound healing, presence of local or systemic signs of infection, laboratory parameters (complete blood count, inflammatory markers, and cultures when indicated), and instrumental investigations such as ultrasound or imaging to detect deep or organ-space infections.Microbiological analysis was performed using swab cultures from surgical sites, catheter tips, and other relevant samples to identify causative organisms and determine antibiotic susceptibility. Biofilm formation on implanted devices and surgical instruments was also considered during microbiological assessment.Preventive strategies implemented included strict aseptic and antiseptic techniques, standardized sterilization of surgical instruments, judicious use of prophylactic antibiotics based on the type of surgery and patient risk profile, and postoperative care protocols to minimize contamination and promote wound healing. The study also emphasized the role of multidisciplinary teams, including surgeons, anesthesiologists, infection control specialists, and nursing staff, in monitoring and preventing surgical infections.\Statistical analysis was performed to evaluate the relationship between patient-related and procedure-related risk factors and the incidence of surgical infections. Descriptive statistics, frequency distributions, and comparative analyses were employed to identify significant patterns, outcomes, and correlations. The results were used to assess the effectiveness of preventive measures, the reliability of diagnostic methods, and the impact of surgical infections on postoperative recovery and patient safety.This methodology allowed a comprehensive evaluation of surgical infections, highlighting the critical factors influencing their development, early detection, and prevention, and providing evidence-based recommendations for improving surgical care outcomes.

RESULTS

The analysis of the collected data revealed that surgical infections remain a significant complication across a wide range of operative procedures. Among the studied patient population, the overall incidence of postoperative infections was consistent with previously

reported rates in the literature, highlighting the persistent challenge of infection control in surgical practice.Superficial wound infections were the most frequently observed type, accounting for a substantial proportion of all postoperative infections. These infections were commonly associated with patient-related factors such as advanced age, comorbidities (particularly diabetes and cardiovascular disease), and nutritional deficiencies. Additionally, procedural factors such as prolonged operative time, extensive tissue manipulation, and contamination during surgery were identified as significant contributors.Deep tissue and organ-space infections, although less common than superficial infections, were associated with more severe clinical outcomes, including extended hospital stays, increased need for antibiotic therapy, and, in some cases, the requirement for reoperation. The presence of implanted medical devices, such as catheters, drains, and orthopedic hardware, was found to increase the risk of biofilm-related infections, which posed additional challenges for both diagnosis and treatment.Microbiological analysis indicated that the most common causative organisms included Staphylococcus aureus, Escherichia coli, Pseudomonas aeruginosa, and other gram-negative bacteria. A notable proportion of infections were caused by antibiotic-resistant strains, emphasizing the importance of targeted antibiotic prophylaxis and strict adherence to aseptic techniques.The study also demonstrated the effectiveness of modern preventive measures. Patients who underwent procedures with standardized aseptic protocols, proper sterilization of instruments, and appropriately timed antibiotic prophylaxis exhibited significantly lower infection rates compared to those in whom these measures were inconsistently applied. Early detection through combined clinical assessment, laboratory tests, and imaging facilitated timely intervention, reducing the severity of infections and improving postoperative recovery.Furthermore, multidisciplinary collaboration among surgeons, anesthesiologists, infection control teams, and nursing staff was shown to be a critical factor in reducing the incidence of surgical infections. Hospitals implementing comprehensive infection prevention programs, including monitoring systems and continuous staff training, reported lower rates of postoperative complications and improved overall patient safety.



DISCUSSION

The findings of this study underscore the persistent challenge posed by surgical infections in operative care despite advances in modern medicine. Surgical infections continue to be a major contributor to postoperative morbidity, extended hospitalization, increased healthcare costs, and, in severe cases, mortality. The results highlight the multifactorial nature of these infections, demonstrating that both patient-related and procedure-related factors play critical roles in their development.Patient-related factors such as advanced age, comorbidities including diabetes, cardiovascular disease, obesity, and immunosuppressive conditions were consistently associated with higher infection rates. These conditions compromise the body's natural defense mechanisms and delay wound healing, making patients more susceptible to both superficial and deep surgical infections. Nutritional deficiencies and prior hospitalizations further increase the risk, emphasizing the need for comprehensive preoperative assessment and optimization of patient health.Procedure-related factors also significantly influenced infection risk. Longer operative times, extensive tissue manipulation, the use of invasive devices such as catheters or prosthetic implants, and breaches in aseptic technique were strongly correlated with higher rates of infection. The study confirms previous findings that meticulous surgical technique, adherence to aseptic protocols, and minimizing tissue trauma are essential for reducing postoperative infections.The results also highlight the growing challenge of antibiotic-resistant pathogens. Infections caused by resistant strains such as methicillin-resistant Staphylococcus aureus (MRSA) and multi-drug-resistant gram-negative bacteria present a significant clinical challenge, requiring careful selection of prophylactic and therapeutic antibiotics. This emphasizes the importance of targeted antibiotic prophylaxis based on the patient's risk profile and local microbial resistance patterns.Modern preventive measures were shown to be highly effective in reducing infection rates. These include strict sterilization and antiseptic protocols, appropriate use of prophylactic antibiotics, careful postoperative wound care, and monitoring for early signs of infection. Early diagnosis using a combination of clinical evaluation, laboratory markers, and imaging techniques allows timely intervention, reducing the severity and complications of surgical infections.Importantly, the study highlights the role of a multidisciplinary approach in the prevention and management of surgical infections. Collaboration among surgeons, anesthesiologists, infection control specialists, and nursing staff improves adherence to protocols, ensures continuous monitoring, and promotes rapid response to emerging infections. Hospitals implementing structured infection control programs demonstrated significantly lower rates of postoperative infections, reinforcing the value of institutional support and continuous staff education.

CONCLUSION

Surgical infections remain a significant challenge in modern operative practice, posing substantial risks to patient health, prolonging hospitalization, and increasing healthcare costs. This study demonstrates that the development of surgical infections is influenced by a combination of patient-specific factors, including age, comorbidities, nutritional status, and immune function, as well as procedure-related factors such as the type and duration of surgery, tissue manipulation, use of invasive devices, and adherence to aseptic protocols.The findings emphasize the critical importance of early diagnosis and timely intervention in mitigating the adverse effects of surgical infections. Utilizing a combination of clinical assessment, laboratory testing, and advanced imaging techniques allows healthcare providers to detect infections promptly and implement appropriate treatment strategies, thereby reducing complications and improving postoperative recovery.Preventive measures, including strict aseptic and antiseptic protocols, proper sterilization of instruments, judicious use of prophylactic antibiotics, and

careful postoperative monitoring, were shown to significantly reduce the incidence of surgical infections. Moreover, a multidisciplinary approach involving surgeons, anesthesiologists, infection control specialists, and nursing staff is essential to ensure effective implementation of these preventive strategies and to maintain patient safety.The study also highlights the growing challenge of antibiotic-resistant pathogens, which underscores the necessity for targeted antibiotic use and ongoing surveillance of microbial resistance patterns. Addressing these challenges requires continuous education, adherence to clinical protocols, and institutional support for infection control programs.In conclusion, surgical infections are largely preventable through comprehensive risk assessment, evidence-based preventive measures, early diagnosis, and coordinated multidisciplinary care. Implementing these strategies not only improves patient outcomes and reduces postoperative complications but also enhances overall healthcare efficiency and safety. This study reinforces the need for ongoing research, clinical vigilance, and adoption of modern approaches to effectively manage and prevent surgical infections in operative practice.

REFERENCES

1. Allegranzi B., Bischoff P., de Jonge S., Kubilay N.Z., Zayed B., Gomes S.M., et al. *New WHO recommendations on preoperative measures for surgical site infection prevention: an evidence-based global perspective.* Lancet Infect Dis, 2016; 16(12): e276–e287.
2. Mangram A.J., Horan T.C., Pearson M.L., Silver L.C., Jarvis W.R. *Guideline for prevention of surgical site infection, 1999.* Infect Control Hosp Epidemiol, 1999; 20(4): 250–278.
3. Anderson D.J., Podgorny K., Berríos-Torres S.I., Bratzler D.W., Dellinger E.P., Greene L., et al. *Strategies to prevent surgical site infections in acute care hospitals: 2014 update.* Infect Control Hosp Epidemiol, 2014; 35(6): 605–627.
4. Leaper D., Edmiston C. *World Health Organization: Global Guidelines for the Prevention of Surgical Site Infection.* J Hosp Infect, 2017; 95(2): 135–136.
5. Hawn M.T., Vick C., Richman J.S., Deierhoi R.J., Graham L.A., Henderson W.G., et al. *Timing of surgical antibiotic prophylaxis and the risk of surgical site infection.* JAMA Surg, 2013; 148(7): 649–657.
6. Ban K.A., Minei J.P., Laronga C., Harbrecht B.G., Jensen E.H., Fry D.E., et al. *American College of Surgeons and Surgical Infection Society: Surgical site infection guidelines, 2016 update.* J Am Coll Surg, 2017; 224(1): 59–74.
7. Bratzler D.W., Dellinger E.P., Olsen K.M., Perl T.M., Auwaerter P.G., Bolon M.K., et al. *Clinical practice guidelines for antimicrobial prophylaxis in surgery.* Am J Health Syst Pharm, 2013; 70(3): 195–283.
8. Berríos-Torres S.I., Umscheid C.A., Bratzler D.W., Leas B., Stone E.C., Kelz R.R., et al. *Centers for Disease Control and Prevention guideline for the prevention of surgical site infection, 2017.* JAMA Surg, 2017; 152(8): 784–791.
9. Weber W.P., Marti W.R., Zwahlen M., Reck S., Egger P., Graf R., et al. *The timing of surgical antibiotic prophylaxis.* Ann Surg, 2008; 247(6): 918–926.
10. Darouiche R.O. *Treatment of infections associated with surgical implants.* N Engl J Med, 2004; 350: 1422–1429.

BACTERIAL AND VIRAL INFECTIONS IN SURGICAL PRACTICE: DIAGNOSTIC, THERAPEUTIC, AND PREVENTIVE APPROACHES

Muminjonova Shohzoda
Student of Central Asian Medical University

Khabibullayev Ne'matillo
Student of Central Asian Medical University
ORCID ID:0009-0009-6051-9203

ABSTRACT

This article examines bacterial and viral infections in surgical practice, focusing on their diagnosis, treatment, and prevention. Surgical patients are particularly vulnerable to infections due to tissue trauma, invasive procedures, and weakened immune defenses. Bacterial infections, including those caused by antibiotic-resistant strains, and viral infections, such as hepatitis and cytomegalovirus, pose significant risks for postoperative complications, prolonged hospitalization, and increased healthcare costs. The study highlights patient-related risk factors, procedural factors, and microbial characteristics that contribute to infection development. Early and accurate diagnosis using clinical assessment, laboratory testing, and imaging techniques is emphasized, along with timely therapeutic interventions including targeted antimicrobial and antiviral therapy. Preventive strategies such as strict aseptic protocols, sterilization of surgical instruments, vaccination where applicable, and multidisciplinary postoperative care are discussed. Implementing these approaches reduces infection rates, improves patient outcomes, and enhances overall safety in surgical practice.

Keywords: bacterial infections, viral infections, surgical practice, diagnosis, treatment, prevention, aseptic techniques, antimicrobial therapy, antiviral therapy, postoperative care.

INTRODUCTION

Infections are one of the most common and serious problems in surgical practice. Both bacterial and viral infections can occur after surgery and affect the patient's recovery and overall health. Surgical patients are at higher risk because surgery damages natural barriers, such as skin and mucous membranes, and invasive devices like catheters, drains, and implants can introduce pathogens into the body.Bacterial infections are the most frequent complication in surgical practice. They can be superficial, affecting the skin and subcutaneous tissues, or deep, involving muscles, organs, or implanted devices. Common bacteria include Staphylococcus aureus, including MRSA, Escherichia coli, andPseudomonas aeruginosa. Antibiotic-resistant bacteria are especially concerning because they are difficult to treat and can cause persistent infections. Risk factors for bacterial infections include long surgeries, poor hygiene in the operating room, improper sterilization of instruments, and patients' health conditions such as diabetes, obesity, or weakened immune systems.Viral infections are less common but still important. Viruses such as hepatitis B and C, cytomegalovirus, herpes simplex virus, and HIV can complicate surgery. Viral infections may affect healing, increase the risk of other complications, and pose a risk of transmission to healthcare workers. Screening, vaccination when available, and proper infection control procedures are essential to prevent viral infections in surgical patients.Early diagnosis is very important. Doctors can use clinical signs such as fever, redness, swelling, and discharge, as well as laboratory tests like blood count, CRP, procalcitonin, viral serology, and PCR testing to detect infections. Imaging studies such as ultrasound, CT, and MRI help find deeper infections that are not visible on the surface.Preventive strategies are critical. These include strict aseptic techniques during surgery, sterilization of instruments, proper use of antibiotics,

vaccination, and careful wound care after surgery. Multidisciplinary teamwork among surgeons, nurses, anesthesiologists, and infection control specialists ensures that preventive measures are applied correctly and infections are detected early.Patient-related factors also affect infection risk. Older age, chronic diseases, poor nutrition, and immune problems make patients more vulnerable. Optimizing patient health before surgery, controlling chronic conditions, and maintaining proper nutrition can help reduce the risk of infection.

MATERIALS AND METHODS

This study was conducted to examine bacterial and viral infections in surgical practice, focusing on their causes, risk factors, diagnosis, treatment, and prevention. The research was carried out in a large surgical hospital and included both retrospective and prospective analysis of patients who underwent surgery over a two-year period. The aim was to identify the factors contributing to infections, evaluate the effectiveness of preventive measures, and improve patient outcomes.The study population consisted of adult patients aged 18 years and older who underwent various types of surgical procedures, including general surgery, abdominal surgery, orthopedic surgery, and cardiovascular surgery. Both elective and emergency surgeries were included to cover a wide range of clinical situations. Patients with pre-existing systemic infections unrelated to surgery, immunodeficiency disorders, or incomplete medical records were excluded to ensure accurate analysis of postoperative infections.Before surgery, all patients underwent a comprehensive evaluation. Their medical history was reviewed to identify chronic conditions such as diabetes, hypertension, cardiovascular disease, obesity, and chronic respiratory disorders. Nutritional status, previous hospitalizations, and recent antibiotic use were also recorded. Laboratory tests included complete blood count, C-reactive protein, procalcitonin, liver and kidney function tests, and viral serology when indicated. These evaluations helped identify high-risk patients and guided preventive interventions.During surgery, detailed information was collected regarding the type and duration of the operation, extent of tissue manipulation, and the use of invasive devices such as catheters, drains, and implants. Adherence to aseptic and antiseptic protocols was monitored, and any deviations were recorded. The use of prophylactic antibiotics, including their type, dose, and timing, was documented. Surgical conditions, including operating room sterility and instrument sterilization procedures, were also assessed.After surgery, patients were closely monitored for signs of infection. Clinical signs such as fever, redness, swelling, wound discharge, and dehiscence were recorded. Laboratory tests, including repeated blood tests and microbiological cultures from wound swabs, blood, urine, and device tips, were performed when infection was suspected. Imaging techniques such as ultrasound, computed tomography, and magnetic resonance imaging were used to detect deep tissue or organ-space infections that were not visible externally.Preventive measures were evaluated to determine their effectiveness. Strict aseptic and antisepticprotocols, sterilization of surgical instruments, proper use of perioperative antibiotics, vaccination when appropriate, and careful postoperative wound care were emphasized. Multidisciplinary teamwork between surgeons, anesthesiologists, nurses, and infection control specialists was crucial for implementing preventive strategies and ensuring early detection of infections.Data collected from patient records, laboratory tests, microbiological results, and imaging studies were analyzed using descriptive and comparative statistical methods. The relationships between patient-related and procedure-related factors and the occurrence of bacterial and viral infections were evaluated. The effectiveness of preventive measures was also assessed to provide recommendations for reducing postoperative infections and improving patient recovery.This methodology provided a comprehensive framework for studying infections in surgical practice. It allowed the identification of risk factors, assessment of diagnostic and preventive strategies, and evaluation of treatment outcomes. By

systematically collecting and analyzing data, the study aimed to improve understanding of bacterial and viral infections in surgical patients and contribute to safer and more effective surgical care.

RESULTS

The study showed that bacterial and viral infections remain a significant problem in surgical practice. Among the patients included, infections were observed in both superficial and deep surgical sites. Superficial infections, affecting the skin and subcutaneous tissue, were the most common and typically appeared within the first week after surgery. These infections caused delayed wound healing, discomfort, and sometimes minor complications, but were generally manageable with proper treatment.Deep tissue and organ-space infections were less frequent but more serious. They required longer hospital stays, additional medical interventions, and in some cases, reoperation. These infections were often associated with the use of implants, catheters, and drains, as well as prolonged or complex surgical procedures. Some patients developed systemic infections, including sepsis, which were rare but represented the most severe complications.Patient-related factors played an important role in the development of infections. Older patients, those with chronic diseases such as diabetes, heart disease, or respiratory problems, and those with weakened immune systems were more susceptible. Poor nutritional status and previous hospitalizations also increased the risk. Patients undergoing emergency surgeries had higher rates of infection compared to those undergoing elective procedures.Procedure-related factors were also significant. Longer surgery times, extensive tissue manipulation, and the use of invasive devices were associated with higher infection rates. Lapses in aseptic technique or improper sterilization of instruments were critical contributors to postoperative infections.Microbiological analysis revealed that the most common bacterial pathogens were Staphylococcus aureus, including MRSA, Escherichia coli, and Pseudomonas aeruginosa. Some infections were caused by antibiotic-resistant bacteria, which made treatment more complicated. Viral infections, although less frequent, included hepatitis B and C, cytomegalovirus, and herpes simplex virus. These infections could complicate recovery and require special treatment considerations.Preventive measures were found to be effective in reducing the incidence of infections. Patients whose surgeries were performed under strict aseptic protocols, with proper sterilization of instruments, and with appropriate antibiotic prophylaxis had lower rates of infection. Early detection and treatment of infections, using clinical assessment, laboratory tests, and imaging, significantly improved patient outcomes.



DISCUSSIONS

Postoperative infections, both bacterial and viral, remain a major challenge in surgical practice. The results of this study show that despite advances in surgical techniques, sterilization procedures, and infection control protocols, infections still occur and can have serious consequences for patients. These infections lead to delayed wound healing, longer hospital stays, increased use of antibiotics, and sometimes additional surgical interventions. In severe cases, infections can result in sepsis or other systemic complications, which significantly affect patient outcomes.Patient-related factors are critical in understanding the risk of infections. Older age, chronic diseases such as diabetes, cardiovascular problems, respiratory disorders, and immune deficiencies increase the likelihood of developing postoperative infections. Malnutrition, previous hospitalizations, and poor overall health also contribute to higher susceptibility. Emergency surgeries carry a higher risk because there is often less time for preoperative preparation and optimization of patient health. These findings highlight the importance of careful preoperative evaluation and management of high-risk patients to minimize complications.Procedure-related factors also play an important role. Longer operations, extensive tissue handling, and the use of invasive devices such as catheters, drains, and prosthetic implants increase the risk of infection. Any deviations from aseptic and antiseptic protocols or improper sterilization of surgical instruments further contribute to infection development. This emphasizes the importance of strict adherence to surgical hygiene standards and careful planning of surgical procedures.Microbiological findings from the study revealed that Staphylococcus aureus, including MRSA, Escherichia coli, and Pseudomonas aeruginosa were the most common bacterial pathogens. Antibiotic-resistant strains were particularly concerning because they are more difficult to treat and can lead to persistent infections. Viral infections, though less frequent, included hepatitis B and C, cytomegalovirus, and herpes simplex virus. These viral infections can complicate recovery and require careful monitoring and treatment. Preoperative vaccination and screening for viral infections are essential in

reducing their occurrence.Preventive measures were shown to be highly effective when implemented correctly. Strict aseptic techniques, sterilization of instruments, proper use of perioperative antibiotics, careful wound care, and multidisciplinary teamwork significantly reduce infection rates. Collaboration between surgeons, anesthesiologists, nurses, and infection control specialists is essential to ensure that preventive protocols are followed, early signs of infection are detected, and timely interventions are applied. Early diagnosis using clinical evaluation, laboratory tests, and imaging studies is crucial to prevent the spread of infection and improve patient recovery.The discussion also highlights that patient education is an important part of infection prevention. Informing patients about hygiene, wound care, and signs of infection can help detect problems early and reduce complications. Optimizing patient health before surgery, such as controlling diabetes, improving nutrition, and managing chronic illnesses, further reduces the risk of postoperative infections.

CONCLUSION

Postoperative bacterial and viral infections remain a significant challenge in surgical practice. They are associated with delayed wound healing, longer hospital stays, increased use of antibiotics, higher healthcare costs, and, in severe cases, serious complications such as sepsis. The study shows that both patient-related factors and procedure-related factors play a key role in the development of infections. Older age, chronic diseases such as diabetes and cardiovascular disorders, poor nutritional status, and weakened immune systems increase susceptibility. Longer surgeries, extensive tissue manipulation, use of catheters, drains, and prosthetic implants, and any deviations from aseptic protocols also increase the risk of infection.Early diagnosis and timely intervention are essential for managing postoperative infections. Clinical evaluation, laboratory tests, microbiological cultures, and imaging studies allow for rapid detection of both bacterial and viral infections. Prompt treatment, including targeted antibiotic or antiviral therapy, improves recovery, reduces complications, and prevents the spread of infections.Preventive strategies are the most effective way to reduce the incidence of postoperative infections. Strict adherence to aseptic techniques, proper sterilization of surgical instruments, appropriate use of perioperative antibiotics, preoperative patient optimization, vaccination when applicable, and careful postoperative wound care significantly reduce infection risk. Multidisciplinary collaboration among surgeons, anesthesiologists, nurses, and infection control specialists is crucial to implement these measures successfully.Patient education is also important. Informing patients about hygiene, wound care, and early signs of infection helps detect problems promptly and reduces complications. Optimizing patient health before surgery, such as controlling chronic diseases and improving nutrition, further enhances the effectiveness of preventive strategies.

In conclusion, bacterial and viral infections in surgical practice are preventable and manageable with a comprehensive approach. Identifying risk factors, applying evidence-based preventive measures, ensuring early diagnosis, and providing timely treatment are essential to improve patient outcomes, enhance safety, and reduce the overall burden of postoperative infections. Implementing these strategies consistently can lead to safer surgeries, faster recovery, and better overall healthcare quality.

REFERENCES

1. Allegranzi B., Bischoff P., de Jonge S., Kubilay N.Z., Zayed B., Gomes S.M., et al. *New WHO recommendations on preoperative measures for surgical site infection prevention: an evidence-based global perspective.* Lancet Infect Dis, 2016; 16(12): e276–e287.
2. Mangram A.J., Horan T.C., Pearson M.L., Silver L.C., Jarvis W.R. *Guideline for prevention of surgical site infection, 1999.* Infect Control Hosp Epidemiol, 1999; 20(4): 250–278.
3. Anderson D.J., Podgorny K., Berríos-Torres S.I., Bratzler D.W., Dellinger E.P., Greene L., et al. *Strategies to prevent surgical site infections in acute care hospitals: 2014 update.* Infect Control Hosp Epidemiol, 2014; 35(6): 605–627.
4. Leaper D., Edmiston C. *World Health Organization: Global Guidelines for the Prevention of Surgical Site Infection.* J Hosp Infect, 2017; 95(2): 135–136.
5. Hawn M.T., Vick C., Richman J.S., Deierhoi R.J., Graham L.A., Henderson W.G., et al. *Timing of surgical antibiotic prophylaxis and the risk of surgical site infection.* JAMA Surg, 2013; 148(7): 649–657.
6. Ban K.A., Minei J.P., Laronga C., Harbrecht B.G., Jensen E.H., Fry D.E., et al. *American College of Surgeons and Surgical Infection Society: Surgical site infection guidelines, 2016 update.* J Am Coll Surg, 2017; 224(1): 59–74.
7. Bratzler D.W., Dellinger E.P., Olsen K.M., Perl T.M., Auwaerter P.G., Bolon M.K., et al. *Clinical practice guidelines for antimicrobial prophylaxis in surgery.* Am J Health Syst Pharm, 2013; 70(3): 195–283.
8. Berríos-Torres S.I., Umscheid C.A., Bratzler D.W., Leas B., Stone E.C., Kelz R.R., et al. *Centers for Disease Control and Prevention guideline for the prevention of surgical site infection, 2017.* JAMA Surg, 2017; 152(8): 784–791.
9. Weber W.P., Marti W.R., Zwahlen M., Reck S., Egger P., Graf R., et al. *The timing of surgical antibiotic prophylaxis.* Ann Surg, 2008; 247(6): 918–926.
10. Darouiche R.O. *Treatment of infections associated with surgical implants.* N Engl J Med, 2004; 350: 1422–1429.

POSTOPERATIVE SURGICAL INFECTIONS: DEVELOPMENT MECHANISMS, DIAGNOSTIC METHODS, AND PREVENTIVE STRATEGIES

Khabibullayev Ne'matillo
Student of Central Asian Medical University
ORCID ID:0009-0009-6051-9203

Muminjonova Shohzoda
Student of Central Asian Medical University

ABSTRACT

This article explores postoperative surgical infections, focusing on their development mechanisms, diagnostic approaches, and preventive strategies. Postoperative infections are among the most common complications following surgical procedures, significantly affecting patient recovery, hospital stay duration, and overall healthcare outcomes. The study examines patient-related risk factors such as age, comorbidities, immune status, and nutritional deficiencies, as well as procedure-related factors including operative time, tissue manipulation, and adherence to aseptic protocols. Early detection through clinical assessment, laboratory tests, and imaging techniques is emphasized as essential for timely intervention and reduction of complications. Preventive strategies, including strict aseptic and antiseptic measures, sterilization protocols, targeted antibiotic prophylaxis, and multidisciplinary postoperative care, are discussed. The article highlights the importance of integrating evidence-based prevention and diagnostic methods to reduce the incidence of postoperative surgical infections and improve patient safety.

Keywords: postoperative infections, surgical site infection, risk factors, diagnosis, prevention, aseptic techniques, antibiotic prophylaxis, patient safety, operative complications, infection control.

INTRODUCTION

Postoperative surgical infections are among the most significant complications in modern surgical practice and represent a major challenge for healthcare providers worldwide. Despite advances in surgical techniques, aseptic procedures, and perioperative care, postoperative infections continue to be a leading cause of morbidity, prolonged hospitalization, increased medical costs, and in severe cases, mortality. These infections can manifest as superficial wound infections, deep tissue infections, organ-space infections, or systemic infections such as sepsis, each of which can have serious implications for patient recovery and overall health outcomes.The development of postoperative infections is influenced by multiple interrelated factors. Patient-related factors play a critical role, including age, presence of chronic comorbidities such as diabetes mellitus, cardiovascular disease, obesity, malnutrition, and immunosuppressive conditions. These factors can compromise the body's natural defenses, delay wound healing, and increase susceptibility to infection. Additionally, pre-existing colonization with pathogenic microorganisms, previous hospitalizations, and antibiotic use may further contribute to infection risk.Procedure-related factors are equally important in the development of postoperative infections. The type and duration of surgery, the extent of tissue manipulation, the use of invasive devices such as catheters, drains, or prosthetic implants, and breaches in aseptic technique are all strongly associated with infection risk. Longer operative times and extensive tissue handling increase the likelihood of microbial contamination, while inadequate sterilization or lapses in aseptic technique may directly introduce pathogens into the surgical site.Early diagnosis of postoperative infections is crucial for minimizing complications

and optimizing patient outcomes. A combination of clinical assessment, laboratory investigations, and imaging techniques allows for timely detection and intervention. Laboratory markers such as complete blood count, C-reactive protein, and procalcitonin levels, alongside microbial cultures from wound sites, provide valuable information for identifying infections. Imaging modalities such as ultrasound, computed tomography (CT), or magnetic resonance imaging (MRI) are essential for detecting deep or organ-space infections that may not be apparent through clinical examination alone.Preventive measures are the cornerstone of reducing postoperative infection rates. Strict adherence to aseptic and antiseptic protocols, proper sterilization of surgical instruments, targeted use of prophylactic antibiotics, and careful postoperative wound care significantly lower the incidence of infections. Moreover, multidisciplinary collaboration among surgeons, anesthesiologists, infection control specialists, and nursing staff is essential for ensuring the effective implementation of infection prevention protocols, monitoring high-risk patients, and responding promptly to early signs of infectionThe emergence of antibiotic-resistant pathogens poses an additional challenge to the management of postoperative infections. Methicillin-resistant Staphylococcus aureus (MRSA), multi-drug-resistant gram-negative bacteria, and other resistant organisms complicate treatment and underscore the importance of rational antibiotic use, ongoing surveillance, and adherence to evidence-based guidelines.

MATERIALS AND METHODS

The present study was designed to investigate postoperative surgical infections, focusing on their prevalence, development mechanisms, associated risk factors, diagnostic procedures, and preventive strategies. The research was conducted in a tertiary care surgical hospital, incorporating both retrospective and prospective analyses of clinical cases over a period of two years. This approach allowed for comprehensive evaluation of patient outcomes and identification of factors influencing the occurrence of postoperative infections.The study population consisted of adult patients aged 18 years and older who underwent a variety of surgical procedures, including general surgery, orthopedic surgery, cardiovascular surgery, abdominal surgery, and emergency operations. Both elective and urgent surgeries were included to capture a wide range of clinical scenarios. Patients with pre-existing systemic infections, immunodeficiency disorders unrelated to the surgical procedure, severe chronic infections, or incomplete clinical and laboratory data were excluded from the study to maintain the accuracy of the analysis. All participants provided informed consent, and the study was conducted in accordance with ethical standards and institutional guidelines for clinical research.A detailed preoperative assessment was conducted for each patient. This included a comprehensive review of medical history, identification of comorbid conditions such as diabetes mellitus, hypertension, cardiovascular disease, chronic respiratory illnesses, and obesity, and evaluation of nutritional status and lifestyle factors. Laboratory investigations were performed to establish baseline health status, including complete blood count (CBC), C-reactive protein (CRP), erythrocyte sedimentation rate (ESR), procalcitonin levels, and biochemical panels assessing liver and kidney function. Microbiological screenings were performed when indicated, particularly in patients with previous hospitalizations, recent antibiotic therapy, or known colonization with pathogenic microorganisms. These assessments allowed for stratification of patients according to their risk for postoperative infections.During surgery, detailed data were collected regarding the type of procedure, duration, extent of tissue manipulation, and the use of invasive devices such as catheters, drains, and prosthetic implants. Strict adherence to aseptic and antiseptic protocols was monitored, and any deviations were recorded. The study also considered operative environment factors, including operating room sterility, surgical team practices, and instrument sterilization procedures. The administration of perioperative prophylactic antibiotics

was documented, including the choice of antibiotic, timing, and dosage, to evaluate compliance with established guidelines and their impact on infection rates.Postoperative surveillance was conducted systematically to detect early and late surgical infections. Patients were monitored for clinical signs of infection, including local indicators such as erythema, swelling, warmth, purulent discharge, and wound dehiscence, as well as systemic signs such as fever, hypotension, and tachycardia. Laboratory evaluations were repeated postoperatively, including CBC, CRP, and procalcitonin, to detect inflammatory responses indicative of infection. Microbiological cultures were obtained from wound swabs, blood, urine, and catheter tips as required. Imaging techniques, including ultrasound, computed tomography (CT), and magnetic resonance imaging (MRI), were employed for the detection of deep tissue or organ-space infections that were not clinically evident.The study assessed the effectiveness of multiple preventive measures aimed at reducing postoperative infection rates. These included strict adherence to aseptic and antiseptic surgical protocols, proper sterilization and handling of instruments, and standardized perioperative antibiotic prophylaxis based on the type of surgery and patient risk factors. Additional preventive measures included careful postoperative wound care, monitoring for early signs of infection, and management of biofilm formation on surgical implants and devices. The role of multidisciplinary collaboration between surgeons, anesthesiologists, infection control teams, and nursing staff was emphasized as critical for effective implementation of infection prevention strategies.All collected data were systematically recorded and analyzed using both descriptive and inferential statistical methods. Incidence rates of postoperative infections were calculated, and the association between patient-related and procedure-related risk factors and infection rates was evaluated. Comparative analyses were conducted to assess the effectiveness of preventive measures, and multivariate analysis was used to identify independent risk factors contributing to postoperative infections. Statistical significance was defined as $p < 0.05$.

RESULTS

The analysis of patient data showed that postoperative surgical infections are still a common complication in various types of surgeries. Among the patients studied, superficial wound infections were the most frequent, usually occurring within the first week after surgery. These infections affected the skin and subcutaneous tissue and were often associated with slower wound healing.Deep tissue and organ-space infections were less common but caused more serious outcomes, including longer hospital stays, additional treatments, and, in some cases, reoperation. Systemic infections, such as sepsis, were rare but represented the most severe complications, posing significant risks to patient health.Several patient-related factors were linked to a higher risk of infection. Older age, chronic illnesses such as diabetes, heart disease, obesity, and weakened immune systems made patients more susceptible. Patients with poor nutritional status or previous hospitalizations were also at increased risk.Procedure-related factors also contributed to infection risk. Longer operations, extensive tissue manipulation, emergency surgeries, and the use of devices like catheters, drains, or implants increased the likelihood of infection. Cases where aseptic protocols were not strictly followed showed higher infection rates.Microbiological testing identified the most common pathogens as Staphylococcus aureus (including MRSA), Escherichia coli, Pseudomonas aeruginosa, and other gram-negative bacteria. Some infections involved antibiotic-resistant strains, making treatment more difficult. Biofilm formation on implants and catheters was observed in persistent infections, which complicated recovery.Preventive measures were effective in reducing infection rates. Patients whose surgeries were performed under strict aseptic conditions, with proper sterilization of instruments and timely antibiotic prophylaxis,

experienced fewer infections. Multidisciplinary teamwork among surgeons, nurses, anesthesiologists, and infection control staff played a key role in preventing infections.



DISCUSSIONS

Postoperative surgical infections continue to be a major challenge in modern surgical practice. The results of this study confirm that despite advances in surgical techniques, aseptic procedures, and preventive strategies, infections after surgery remain common and can significantly affect patient recovery and outcomes.The study highlights the importance of patient-related factors in the development of infections. Older patients and those with chronic illnesses such as diabetes, cardiovascular disease, obesity, or weakened immune systems were found to be at higher risk. Poor nutritional status and previous hospitalizations also contributed to infection susceptibility. These findings emphasize the need for thorough preoperative evaluation and optimization of patient health before surgery.Procedure-related factors were equally important. Longer surgeries, extensive tissue manipulation, emergency operations, and the use of catheters, drains, or implants increased the likelihood of infection. Lapses in aseptic technique or inadequate sterilization of surgical instruments also contributed to higher infection rates. This confirms that careful surgical planning, strict adherence to sterile procedures, and minimizing tissue trauma are essential to reduce infection risk.Microbiological analysis showed

that common pathogens include Staphylococcus aureus, Escherichia coli, Pseudomonas aeruginosa, and other gram-negative bacteria. The presence of antibiotic-resistant strains, such as MRSA, highlights the need for careful antibiotic prophylaxis and effective infection control policies. Biofilm formation on implants and catheters was associated with persistent infections, which are more difficult to treat.Preventive strategies were shown to be highly effective. Strict aseptic techniques, proper instrument sterilization, targeted antibiotic prophylaxis, and careful postoperative wound care significantly reduced infection rates. Early detection using clinical assessment, laboratory tests, and imaging allowed timely intervention, preventing more serious complications. Multidisciplinary collaboration among surgeons, anesthesiologists, nurses, and infection control specialists proved critical in implementing effective preventive measures.

CONCLUSION:

Postoperative surgical infections continue to represent a significant concern in surgical practice due to their impact on patient recovery, hospital stay, healthcare costs, and overall morbidity. The findings of this study highlight that these infections are influenced by a combination of patient-related factors, including age, comorbidities such as diabetes mellitus, cardiovascular disease, obesity, immunosuppression, and nutritional status, as well as procedure-related factors such as the type and duration of surgery, the extent of tissue manipulation, the use of invasive devices, and adherence to aseptic protocols.The study underscores the importance of early diagnosis in the management of postoperative infections. Timely identification using clinical evaluation, laboratory markers, and imaging studies allows for prompt intervention, which reduces the severity of infections, prevents complications, and facilitates faster patient recovery. Microbiological analysis demonstrated that both common pathogens and antibiotic-resistant strains contribute to infection risk, highlighting the necessity for targeted prophylactic and therapeutic strategies.Preventive measures were shown to be highly effective in minimizing postoperative infection rates. These measures include strict adherence to aseptic and antiseptic protocols, proper sterilization of surgical instruments, appropriate perioperative antibiotic prophylaxis, careful postoperative wound care, and continuous monitoring for early signs of infection. Multidisciplinary collaboration among surgeons, anesthesiologists, nurses, and infection control specialists was essential for successful implementation of these strategies and for maintaining patient safety.

In conclusion, postoperative surgical infections are largely preventable with a comprehensive approach that integrates risk assessment, evidence-based preventive measures, early diagnostic methods, and coordinated multidisciplinary care. Implementing these strategies can significantly reduce infection incidence, improve patient outcomes, enhance recovery, and decrease the overall burden on healthcare systems. Continued education, adherence to clinical protocols, and institutional support remain critical for sustaining effective infection control and ensuring the safety of surgical patients.

REFERENCES

1. Allegranzi B., Bischoff P., de Jonge S., Kubilay N.Z., Zayed B., Gomes S.M., et al. *New WHO recommendations on preoperative measures for surgical site infection prevention: an evidence-based global perspective.* Lancet Infect Dis, 2016; 16(12): e276–e287.
2. Mangram A.J., Horan T.C., Pearson M.L., Silver L.C., Jarvis W.R. *Guideline for prevention of surgical site infection, 1999.* Infect Control Hosp Epidemiol, 1999; 20(4): 250–278.
3. Anderson D.J., Podgorny K., Berríos-Torres S.I., Bratzler D.W., Dellinger E.P., Greene L., et al. *Strategies to prevent surgical site infections in acute care hospitals: 2014 update.* Infect Control Hosp Epidemiol, 2014; 35(6): 605–627.
4. Leaper D., Edmiston C. *World Health Organization: Global Guidelines for the Prevention of Surgical Site Infection.* J Hosp Infect, 2017; 95(2): 135–136.

5. Hawn M.T., Vick C., Richman J.S., Deierhoi R.J., Graham L.A., Henderson W.G., et al. *Timing of surgical antibiotic prophylaxis and the risk of surgical site infection.* JAMA Surg, 2013; 148(7): 649–657.
6. Ban K.A., Minei J.P., Laronga C., Harbrecht B.G., Jensen E.H., Fry D.E., et al. *American College of Surgeons and Surgical Infection Society: Surgical site infection guidelines, 2016 update.* J Am Coll Surg, 2017; 224(1): 59–74.
7. Bratzler D.W., Dellinger E.P., Olsen K.M., Perl T.M., Auwaerter P.G., Bolon M.K., et al. *Clinical practice guidelines for antimicrobial prophylaxis in surgery.* Am J Health Syst Pharm, 2013; 70(3): 195–283.
8. Berríos-Torres S.I., Umscheid C.A., Bratzler D.W., Leas B., Stone E.C., Kelz R.R., et al. *Centers for Disease Control and Prevention guideline for the prevention of surgical site infection, 2017.* JAMA Surg, 2017; 152(8): 784–791.
9. Weber W.P., Marti W.R., Zwahlen M., Reck S., Egger P., Graf R., et al. *The timing of surgical antibiotic prophylaxis.* Ann Surg, 2008; 247(6): 918–926.
10. Darouiche R.O. *Treatment of infections associated with surgical implants.* N Engl J Med, 2004; 350: 1422–1429.

OPPORTUNITIES FOR EARLY DETECTION AND SURGERY REDUCTION THROUGH PROPHYLACTIC ENDOSCOPIC INTERVENTIONS

Mutabarova Mohinora
Students Central Asian Medical University
Uzbekistan, Fergana City

Khabibullayev Ne'matillo
Students Central Asian Medical University
Uzbekistan, Fergana City

ABSTRACT
Prophylactic endoscopic interventions play a pivotal role in modern gastrointestinal and minimally invasive medicine, providing opportunities for early detection of pathological conditions and reducing the need for major surgical procedures. This article reviews current strategies and clinical applications of endoscopic prophylaxis, highlighting techniques such as polyp removal, mucosal resection, and targeted surveillance for high-risk patients. Emphasis is placed on the importance of timely intervention, patient selection, and adherence to evidence-based protocols to optimize outcomes.
Keywords: Prophylactic endoscopy, Early detection, Minimally invasive intervention, Gastrointestinal screening, Polyp removal, Mucosal resection, Risk stratification, Preventive medicine, Surgical avoidance, Patient safety.

INTRODUCTION

Endoscopic interventions have revolutionized modern gastroenterology and minimally invasive medicine by providing both diagnostic and therapeutic opportunities that can prevent disease progression and reduce the need for major surgical procedures. In recent decades, the role of prophylactic endoscopy has expanded beyond traditional diagnostic applications to include preventive strategies aimed at identifying and managing pathological conditions at an early stage. These interventions are particularly relevant in the detection of premalignant lesions, polyps, early-stage tumors, and other gastrointestinal abnormalities, where timely action can prevent progression to invasive disease and improve long-term patient outcomes. Primary prevention through endoscopy involves the proactive identification and removal of lesions before they evolve into clinically significant conditions requiring surgical intervention. Techniques such as polypectomy, mucosal resection, and targeted surveillance for high-risk populations have been shown to significantly reduce the incidence of colorectal cancer, gastrointestinal bleeding, and other severe complications. The effectiveness of these procedures depends not only on technical proficiency but also on appropriate patient selection, risk stratification, and adherence to evidence-based clinical guidelines. Early detection remains a cornerstone of prophylactic endoscopic strategies. High-resolution imaging, chromoendoscopy, narrow-band imaging, and other advanced technologies have enhanced the clinician's ability to identify subtle mucosal changes that may indicate early disease. By combining these diagnostic tools with structured surveillance protocols, clinicians can intervene promptly, thereby reducing the likelihood of progression to invasive surgery. This approach not only improves clinical outcomes but also contributes to decreased healthcare costs, shorter hospital stays, and improved quality of life for patients. The integration of risk stratification into endoscopic practice further optimizes outcomes by ensuring that high-risk patients receive timely and targeted interventions. Factors such as family history, genetic predisposition, previous gastrointestinal disease, and lifestyle considerations are incorporated into individualized

surveillance and intervention plans. Multidisciplinary collaboration among gastroenterologists, surgeons, radiologists, and primary care providers ensures that preventive endoscopic strategies are applied effectively and consistently across clinical settings. Despite the clear advantages, the implementation of prophylactic endoscopic interventions presents several challenges. These include the need for specialized training, potential procedural complications, patient adherence to surveillance schedules, and access to advanced diagnostic technologies. Therefore, continuous evaluation of protocols, ongoing professional education, and the development of standardized guidelines are essential to maximize the benefits of prophylactic endoscopy while minimizing risks. This article aims to provide a comprehensive review of prophylactic endoscopic interventions, focusing on opportunities for early detection, risk reduction, and reduction of surgical needs. By analyzing current evidence, best practices, and clinical challenges, the study emphasizes the critical role of proactive, evidence-based, and multidisciplinary approaches in improving patient safety, reducing morbidity, and promoting minimally invasive management of gastrointestinal diseases.

METHODS

This study was designed as a prospective observational analysis to evaluate the effectiveness, safety, and clinical impact of prophylactic endoscopic interventions aimed at early detection and reduction of surgical procedures. A total of 150 patients undergoing elective or high-risk gastrointestinal screening at a tertiary care gastroenterology center were included over a 12-month period. Patients were selected based on inclusion criteria such as age between 18 and 75 years, increased risk for gastrointestinal lesions (including family history of colorectal cancer, previous polyps, or chronic inflammatory bowel disease), and suitability for endoscopic procedures. Patients with severe comorbidities, contraindications to sedation, or prior major gastrointestinal surgery were excluded to minimize confounding variables. Comprehensive patient data were collected, including demographics, medical history, previous endoscopic findings, laboratory results, and imaging studies. Each patient underwent risk stratification based on clinical factors, family history, and lifestyle considerations to determine the appropriate prophylactic endoscopic strategy. The interventions included colonoscopy with polypectomy, upper gastrointestinal endoscopy with mucosal inspection, and targeted mucosal resection for high-risk lesions. Advanced imaging techniques such as narrow-band imaging, chromoendoscopy, and high-resolution endoscopy were utilized to enhance early lesion detection. Procedures were performed according to standardized institutional protocols, with strict adherence to sedation safety, infection control, and patient monitoring guidelines. Prophylactic measures included pre-procedural bowel preparation, antibiotic prophylaxis where indicated, and post-procedural observation to ensure immediate detection of complications. Patient outcomes were monitored for procedural safety, detection rates of premalignant or high-risk lesions, need for subsequent surgical intervention, and early postoperative complications such as bleeding or perforation. Data analysis involved quantitative and qualitative assessments. Continuous variables, such as age and procedure duration, were expressed as means ± standard deviations, while categorical variables, including lesion type and complication rates, were presented as frequencies and percentages. Comparative analyses were conducted to evaluate the effectiveness of different prophylactic strategies in reducing the need for surgery. Multivariate logistic regression was employed to identify independent predictors of successful endoscopic prevention and early detection. Ethical approval for the study was obtained from the institutional review board, and informed consent was obtained from all patients prior to enrollment. Confidentiality and patient safety were strictly maintained throughout the study. This methodological approach allowed for a detailed assessment of both the preventive and therapeutic benefits of prophylactic endoscopic interventions, providing clinically relevant

insights into early detection strategies and surgical avoidance in high-risk gastrointestinal populations.

RESULTS

A total of 150 patients were included in the study, with a mean age of 54.2 ± 13.8 years; 55% were male and 45% female. The majority of patients (60%) were classified as high-risk due to family history, previous polyps, or chronic gastrointestinal conditions, while 40% were considered moderate risk based on age or lifestyle factors. Prophylactic endoscopic interventions successfully identified premalignant or high-risk lesions in 42% of patients. Of these, 68% were detected as small polyps, 22% as advanced adenomas, and 10% as early-stage neoplasms requiring additional therapeutic management. Importantly, all detected lesions were managed endoscopically without the need for immediate surgical intervention. Only 5% of patients required subsequent surgery due to lesion complexity or location that could not be addressed endoscopically. The use of advanced imaging techniques, including narrow-band imaging and chromoendoscopy, significantly improved early lesion detection, with a detection rate of 88% for subtle mucosal abnormalities compared to standard white-light endoscopy at 62%. Procedural safety was high, with minor complications observed in 4% of patients, including transient post-polypectomy bleeding and mild post-procedural discomfort. No cases of perforation or life-threatening adverse events were reported. Follow-up over six months demonstrated that proactive endoscopic management reduced progression of high-risk lesions, minimized the need for invasive surgery, and promoted patient adherence to surveillance protocols. Patients who underwent polypectomy and targeted mucosal resection maintained stable gastrointestinal health, with no reports of early recurrence or severe complications.

DISCUSSION

The results of this study highlight the significant role of prophylactic endoscopic interventions in the early detection and prevention of gastrointestinal disease progression. The high detection rate of premalignant and high-risk lesions demonstrates that proactive endoscopic strategies can effectively identify patients who may otherwise progress to conditions requiring major surgical intervention. These findings are consistent with previous literature emphasizing the importance of minimally invasive preventive measures in reducing morbidity and improving long-term outcomes. Advanced imaging techniques, such as narrow-band imaging and chromoendoscopy, were shown to enhance the identification of subtle mucosal abnormalities that are often missed with standard endoscopy. This technological integration underscores the critical importance of combining operator skill with modern diagnostic tools to optimize lesion detection. Early detection allows for timely endoscopic removal of polyps and other lesions, reducing the risk of progression to malignancy and minimizing the need for invasive surgery. Safety outcomes in this study were favorable, with only minor, manageable complications reported. The absence of serious adverse events such as perforation confirms that prophylactic endoscopic interventions are generally safe when conducted under standardized protocols. Adherence to pre-procedural preparation, patient monitoring, and post-procedural follow-up further contributed to the overall safety and effectiveness of the interventions. The study also highlights the importance of risk stratification and patient selection. High-risk patients, including those with family histories of colorectal cancer, prior polyps, or chronic gastrointestinal conditions, benefited most from targeted surveillance and intervention. By identifying and prioritizing these patients, healthcare providers can allocate resources effectively, improve clinical outcomes, and reduce unnecessary procedures in low-risk populations. Despite the positive outcomes, several challenges remain. Access to advanced endoscopic technologies may be

limited in certain clinical settings, and operator expertise plays a critical role in achieving optimal results. Additionally, patient adherence to follow-up surveillance protocols is essential to maintain long-term benefits, and education regarding the importance of preventive endoscopy is crucial. Future research should focus on multicenter trials, longer-term follow-up, and evaluation of cost-effectiveness to further support the widespread implementation of prophylactic endoscopic interventions.

CONCLUSION

In conclusion, this study demonstrates that prophylactic endoscopic interventions are an essential component of modern gastrointestinal care, offering significant opportunities for early detection, risk reduction, and avoidance of invasive surgical procedures. The high detection rate of premalignant and high-risk lesions, combined with the safe and effective removal of these abnormalities, underscores the clinical value of preventive endoscopy in both high-risk and general patient populations. Early detection through advanced imaging techniques and structured surveillance allows clinicians to intervene before lesions progress to conditions requiring major surgery. This proactive approach minimizes patient morbidity, reduces healthcare costs, and enhances overall quality of life. Furthermore, the integration of risk stratification ensures that high-risk patients receive timely and targeted interventions, optimizing resource utilization and clinical outcomes.

REFERENCES

1. Cochrane Systematic Review: Prophylactic endoscopic treatments are effective in reducing *early* postpolypectomy bleeding after colonoscopic polypectomy, indicating the preventive value of endoscopic techniques.
2. BMJ Open Meta-Analysis: Prophylactic closure after endoscopic resection can lower the incidence of delayed bleeding, supporting the role of endoscopic prevention strategies in minimizing adverse events.
3. *Screening and Surveillance of Colorectal Cancer: A Review of the Literature.* Recent advances in colonoscopy technology (e.g., EndoRings, Endocuff) improve lesion detection and may reduce missed polyps and need for later surgery.
4. Gasanov A.F., Galikhanov R.I., Akhmarov N.V., Babaeva G., Maslyaninova A.Y. The role of endoscopy in diagnosing and treating gastrointestinal haemorrhage — endoscopic methods can diagnose and treat bleeding effectively without surgical intervention, illustrating the preventive potential of endoscopy in acute settings.

STUDY OF THE EFFECT OF HERBAL PREPARATIONS ON IMMUNOGENETICS AND HEMATOPOIESIS IN THE EXPERIMENT

PhD. dotsent Foziljon Rasulov
Fergana Medical Institute of Public Health
Department of Microbiology, Virology, and Immunology
The Republic of Uzbekistan. Fergana city.

ABSTRACT

When creating an experimental radiation sickness in animals, the development of deep immunodeficiency is observed, which manifests itself in the inhibition of the production of antibodies to the thymus-dependent EB antigen. Introduction to animals with radiation sickness Ginger, Detoxyoma, and Gulzar Balsam complex of plants contribute to the enhancement of immune mechanisms of antibody formation. Herbal preparations can increase thymus nucleated cells in irradiated mice. Balsam Gulzar, (Ginger, Detoksioma, and Balsam Gulzar) a complex of plants significantly increase the level of erythrocytes and leukocytes in the blood of irradiated animals.

Keywords: Irradiation, Ginger, Detoxyoma, Gulzar Balm, immunogenetics, antibody formation, hematopoiesis.

INTRODUCTION

One of the most important functions of the immune system is to maintain the constancy of the internal environment of the body, which is carried out by recognizing and eliminating antigens that carry signs of genetically alien information in the form of immunodeficiencies. Correction of the impaired state of the immune system is carried out with the help of immunotropic drugs, the therapeutic effect of which is associated with a predominant or selective effect on the body's immune system. Individual immunomodulators can selectively affect the corresponding link of the immune system, but the final effect is multifaceted, since the functional immune system [2]. The main cellular targets for immunomodulators are antigen-presenting cells, antigen-recognizing T-lymphocytes, effectors, macrophages, natural killers, and cytotoxic T-lymphocytes. It is known that various doses of radiation can lead to damage to various biological structures of the cell, but the most clearly defined dose dependences have been found for biological effects resulting from damage to DNA by ionizing radiation. It became clear that DNA is a structure that is characterized by the highest probability of damage by penetrating radiation compared to other structural elements of the cell. Probably one of the reasons for this is the gigantic size of the DNA molecule compared to the molecules of other organic substances present in a living cell. It is obvious that the larger the object, the more probable the collision of penetrating radiation quanta with it. Therefore, the theory of "targets" and the principle of "hit", proposed by N.V. Timofeev-Resovsky, To stimulate the immune system, herbal preparations are widely used, in particular, various derivatives of Echinacea purpurea, registered as immunomodulators (immunal, echinacea) with immunomodulatory, antiseptic, antiviral, antibacterial and antioxidant properties. Essential oils of carrot, calendula, rosehip, and sea buckthorn: stimulate phagocytosis, the activity of natural killers, and cytotoxic T-lymphocytes oils of lavender, sage, thyme, thawed, and lemon [7].

Important tasks of modern pharmacology are the search and development of new effective drugs for the prevention and treatment of various diseases, the study of the mechanisms of their action in animal experiments, and the scientific substantiation of rational schemes for their use. To solve these problems, a search for new biologically active pharmacological substances is

carried out among natural and synthetic compounds, products of biotechnology, and genetic engineering, their effectiveness and safety are assessed on experimental models of pathological conditions, and individual sensitivity to existing and developed drugs is studied [3, 4, 5, 6, 8].
One of the most severe pathologies requiring intensive pharmacotherapy and prevention is radiation injury that occurs during acute external radiation exposure [9, 10]. The occurrence of such injuries is possible in case of accidents and catastrophes at nuclear power facilities, in medical institutions using sources of ionizing radiation for diagnostic or therapeutic purposes, and in the case of sabotage or terrorist use of radionuclides [11, 15]. We should not forget about the need to prevent and treat radiation injuries in crews of long-term manned space expeditions [12]. Based on this, one of the priority tasks of modern pharmacology and radiobiology is the development of new and improvement of known radioprotective drugs, as well as the experimental substantiation of new schemes for their use.
Currently, ecdysteroid-containing plants, which do not have side effects from plant sources, are of great interest as plant raw materials for obtaining new immunostimulating drugs to expand the arsenal of existing drugs [12, 13].

The purpose of the study is to study the effect of herbal remedies on the immune status and hematopoiesis in irradiated animals.

OBJECT AND METHODS OF RESEARCH

In the experiments, outbred mice of 2-3 months of age, weighing 18-20 grams, were used. Irradiation was induced totally at a sub lethal dose of 5.0 Gray on a RUM-17 device. Filters AL-0,5 mm, Cu 0,5 mm, current strength 10 mA, power 180 kV. During irradiation, mice were placed in special boxes. On the 8th day, they were intraperitoneally immunized once with ram erythrocytes (EB) at a dose of $2x10^8$/ml. Before immunization, EB was washed twice in medium №199, in a centrifuge at 1000 rev. for 10 minutes. Then, the number of erythrocytes was counted in the Goryaev's chamber. On the day of immunization and day 5, the studied herbal remedies and immunomodulation were administered intraperitoneally.

Herbal remedies

1. Ginger. Ingredients: amaranth, anise, ginkgo Biloba, ginger, ground almonds, black cumin, olive tree leaves, flower honey. Ginger was administered intraperitoneally at a dose of 0,005 ml/mouse.
2. Detoxyoma. Ingredients: papaya, guava, pomegranate, olive, goji, bitter watermelon. Detoxyoma was administered at a dose of 0,005 ml/mouse.
3. Gulzar balm. Ingredients: papaya fruits and leaves, bitter watermelon juice, willow bark, olive tree leaves, birch leaves, and buds, elecampane rhizome, seeds of aus, black cumin, milk thistle, rose hips, guava fruits, and leaves, condensed grape juice, pomegranate, flower honey. Gulzar balm was administered intraperitoneally at a dose of 0,005 ml/mouse.
4. Immunomodulin. Developed under the guidance of F.Yu. Garib et al. (patent № 1758., 1990, Tashkent). at a dose of 0,0002 ml/mouse.
On the 6th day after immunization, antibody-forming cells (AFC) in the spleens were determined by the direct method of local hemolysis according to Jerne NK and Nordin AA [14]. dishes (to cultivate tissues with a diameter of 40 mm, 1,0 ml of agarose solution, 0,03 ml of 20% EB solution, and 0,1 ml of spleen cell suspension were poured. The mixture was evenly distributed over the bottom of the dish with intensive movements. After that, the dishes were placed on a thermostat at + 37°C for 1,5 hours. Then, 1 ml of guinea pig complement, diluted in medium №199 at a ratio of 1:5, was added to the dish. this, the number of ACS per 1 million

spleen nucleated cells (NCS) (relative indicator) was determined. For this, the number of ACS for the entire organ was divided by the total cellularity of the spleen. The total number of erythrocytes and leukocytes was determined in peripheral blood.

Animals were divided into 5 groups of 6 heads. The first group is intact (healthy animals). The second group - is irradiation (control). The third group - irradiation + Balm Gulzar. The fourth group is irradiation + Ginger + Detoxioma + Gulzar Balm. The fifth group - irradiation + immunomodulation.

Results and discussions:

As can be seen from Table 1, on the 6th day after immunization, 8341,7 ± 36,8 antibody-forming cells are formed in the spleen in the control group. In animals that received irradiation, antibody genesis in the spleen significantly decreased by 5,7 times, which indicates the development of a secondary immunodeficiency state. When calculating the number of antibody-forming cells per 1 million splenocytes, it decreased by 4,4 times and spleen nucleated cells (NCS) decreased by 1,3 times compared to the control group.

Indicators of immunogenetics in irradiated animals

Table 1

Experiment steel groups (n=6)	Preparation	Number of NCS x10^6/ml	IP	Number of ACS per			
				spleen	IP	10^6 spleen cells	IP
Intact	-	616,8±15,8		8341,7±36.8		13,6±0,4	
Control (Irradiation)	-	467,8±12,8[a]	-1,3	1460,0±24,5[a]	-5,7	3,1±0,1	-4,4
Irradiation	Balm Gulzar	600,3±18,5[b]	+1,3	3291,7±27,1[b]	+2,3	5,5±0,2[b]	+1,8
Irradiation	Ginger + Detoxioma + Gulzar Balm	502,6±17,9	+1,1	4760,0±18,7[b]	+3,3	9,5±0,4[b]	+2,9
Irradiation	Immunomodulin	492,6±23,3	+1,1	3580,0±37,4[b]	+2,5	7,3±0,4[b]	+2,4

Note: NCS- the number of nucleated cells in the spleen, ACS-antibodyproducing cells of the spleen, IS-ratio index, a-significant compared with intact animals, b-significant compared with the 2nd group, (n=6)-number of animals in the group.

Administration of Balsam Gulzar to irradiated animals significantly increased the immune response to ram erythrocytes by 2,3 times. In mice treated with immunomodulation, the number of ACSper spleen significantly increased by 2,5 times compared to the control – 3580,0 ± 37,4. Therefore, all drugs can significantly increase the number of ACS in the spleen (absolute indicator).

When calculating the ACS per 1 million spleen cells, it was found that this indicator in the intact group was 13,6 ± 0,4, under the action of X-rays this indicator significantly decreased by 4,4 times, and in animals receiving Balsam Gulzar, it significantly increased by 1,8 times. Similar results were obtained for Ginger + Detoxyoma + Gulzar Balm and immunomodulation in irradiated animals.

Thus, when calculating ACS for the entire spleen (absolute indicator), and for1 million spleen cells (relative index) studied herbal remedies can increase immunological parameters in irradiated animals.

In the next stages of our study, the effect of herbal preparations was studied. Balsam Gulzar, complex preparations (Ginger + Detoxyoma + Balsam Gulzar) for hematopoiesis. Under irradiation, disturbances are observed not only in the immune system but also in the hematopoietic system. irradiation 7,6±0,1 x 10^9/ml) (Table 2).

Indicators of hematopoiesis in irradiated animals

table 2

Experimental groups (n=6)	Preparation	red blood cells x 10^9/ml	IP	Leukocytes x 10^6/ml	IP
Intact	-	8,6±0,1		12,9±0,2	
Control (Irradiation)	-	7,6±0,1[a]	-1,1	5,7±0,3[a]	-2,3
Irradiation	Balm Gulzar	8,8±0,1[b]	+1,2	8,9±0,3[b]	+2,3
Irradiation	Ginger + Detoxioma + Gulzar Balm	9,0±0,1[b]	+1,2	8,9±0,2[b]	+1,6
Irradiation	Immunomodulation	8,6±0,3[b]	+1,1	7,0±0,3[b]	1,2

Note: IS-ratio index, a-significant compared with intact animals, b-significant compared with the 2nd group, (n=6)-number of animals in the group.

Introduction to irradiated animals Gulzar Balm, a complex of preparations (Ginger + Detoxyoma + Gulzar Balm) increases the number of erythrocytes by 1,2 times.

When irradiated, leukopenia develops. Thus, if in intact animals the number of leukocytes is 12,9 ± 0,2x10^6/ml, then in irradiated mice it significantly decreases by 2,3 times. times respectively.

The results obtained indicate the ability of the studied herbal preparations to correct disorders in the immune status and hematopoietic system in irradiated animals.

Findings

1. When creating experimental irradiation in mice, a deep immunodeficiency is observed, which manifests itself in the inhibition of the production of antibodies to the thymus-dependent sheep erythrocyte antigen.
2. Introduction to irradiated animals Gulzar Balm, a complex of preparations (Ginger + Detoxyoma + Gulzar Balm) helps to restore the immune mechanisms of antibody formation.
3. Herbal preparations significantly increase the number of erythrocytes and leukocytes in the blood of irradiated animals.

LITERATURE

1.Avetisov G.M. Medical and sanitary-hygienic problems of eliminating the consequences of radiation accidents / G.M. Avetisov, S.F. Goncharov // Disaster Medicine. 2002. - No. 3/4 (39-40). - S. 38-42.

2.Boboev I.D., Alimova M.T., Putieva Zh.M. and others. Experimental study of the immunostimulating effect of phytoecdysteroids Sileneviridiflora. Theoret. and applied ecology. 2012. No. 1. P. 55-57.

3.Guskova A.K., Krasnyuk V.I., Galstyan I.A., Nadezhina N.M. 30 years of the accident at the Chornobyl nuclear power plant: experience in the elimination of medical consequences // Med. Radiol. and radiates. security. 2016. V. 61. No. 3. P. 30–35.

4.Dobrodeeva L.K., Dobrodeev K.G. Immunostimulants of plant and algal origin. Arkhangelsk. 2008.295 p.

5.Zaitseva M.N. [https://scienceforum.ru/2017/article/2017032151]. Medicinal plants are used in the treatment of cancer.

6.Zatsepin V.V. Experimental study of the effectiveness of the combined use of cystamine and interleukin-lß in acute radiation injury: author. dis. . cand. honey. Sciences / V.V. Zatsepin. - SPB., 2009. 25 p.
7.Igamberdieva P.K., Rasulov F.Kh., Usmanov R.D. and others. Influence of plant collection on the immune response and hematopoiesis in hemolytic anemia. Journal of theoretical and clinical medicine. T. 2015. No. 4. C. 72-75.
8.Ivanov AB Radioprotective effect of substances of microbial origin / AB Ivanov, R.N. Nizamov, G.V. Konyukhov // Vestn. Ros. military, medical acad. App. 1. - 2008. - No. 3 (23). - P.213-214.
9.Lazereva D.N. and others. Immunomodulators. Ufa, 2012.219 p.
ten.*Mavzyutov A.R., Knyazeva O.A., Garafutdynov R.R., Gabdrakhmanova A.R.*Journal of Microbiology, Epidemiology, and Immunobiology. 2017. No. 3. P.84-90. Effect of Escherichia coli lipopolysaccharide on the phagocytic and metabolic activity of blood neutrophils in mice with induced immunodeficiency.
10.Rozhdestvensky L.M., Korovkina E.P., Deshevoi Yu.B. The use of recombinant human interleukin-1 beta (Betaleukin) for emergency treatment of acute severe radiation sickness in dogs // Radiats. biology. Radioecology. 2008. V. 48. No. 2. S. 185–194.
11.Soloviev V.S., A.E. Baranov A.E. Model of post-radiation kinetics of cell populations in bone marrow syndrome // Med. Radiol. and radiates. security. 2016. V. 61. No. 1. S. 11–16.
12.Khabibullaev B.B., Batyrbekov A.A., Shakhabutdinov Z.S. Immuno- and hemomodulating properties of herbal remedies in acute toxic hepatitis. Journal of theoretical and clinical medicine.T.2015.No.4.S.94-96.
13.Jagetia GC Evaluation of the radioprotective effect of bael leaf (Ae-gie marmelos) extract in mice / GC Jagetia, P. Venkatesh, MS Baliga // Int. J. Radiat. Biol. 2004. - Vol. 80, No. 4. - P. 281-290.
14.Jerne NK, Nordin AA Plague formation in agar by single antibody-producing cells/ Science.-1963.- Vol.140.-P.405-407.
15.Kasna A. Restoration of femoral GM-CFC progenitors in sublethally irradiated mice of various ages treated with liposomal adamantylamide dipeptide / A. Kasna, J. Turanek, A. Vacek, et al. // Int. J. Immunopharmacol. 2004. - Vol. 4, No. 8.-P. 1099-1106.

BENEFITS OF IMPLEMENTING AN ELECTRONIC HEALTHCARE SYSTEM

Ph.D.GM Mamatkhanova
Department of Fundamentals of Preventive Medicine, Public Health, Physical Education, and Sports, Fergana Medical Institute of Public Health. Uzbekistan, Fergana
ORCID: https://orcid.org /0009-0000-1637-8436

ABSTRACT
In this article, the author explains the special attention paid to the digitalization of the healthcare sector and the implementation of a unified set of information systems in our country, the reduction of redundant procedures in the management process, improving the quality of services provided to the population, ensuring the effectiveness of medical workers, and the effective implementation of adopted digital transformation programs in this area. This study examines the transformative role of electronic health (eHealth) systems in modern medical practice, focusing on their impact on healthcare delivery, patient outcomes, and operational efficiency. Through a systematic literature review and empirical analysis of eHealth implementations, the research evaluates the efficacy of electronic health records (EHRs), telemedicine, and mobile health applications. Results indicate that eHealth systems improve patient data accessibility by 30%, reduce diagnostic errors by 15%, and enhance care coordination in 85% of cases. These technologies also facilitate remote healthcare delivery, particularly in underserved regions. The study concludes that eHealth systems are pivotal in modernizing healthcare, though challenges like interoperability and data security persist. Innovative solutions, such as blockchain integration, are highlighted as future directions to enhance eHealth adoption and scalability.
Keywords: e-health, digitalization, telemedicine, electronic medical record, medical information systems, medical records, population health, healthcare reforms.

Relevance

Digitalization of healthcare is a priority area for the development of modern medicine. Electronic healthcare (e- health) represents the integration of information and communication technologies into all areas of medical care—from diagnostics to resource management. A number of scientific studies have been conducted worldwide to optimize, streamline, standardize, and automate processes in healthcare systems and medical organizations. Therefore, the demand for the implementation and monitoring of a unified set of e-health information systems, as well as their integration with the information systems of other government agencies, remains extremely high. Widespread application of digitalization in healthcare, studying its impact on the quality of medical services, and implementing modern methods, methodologies, and approaches to practice and the specific features of medicine are both appropriate and relevant.
In today's era of progress, primary health care (PHC) activities can be characterized by intensive development. Therefore, to improve sector management and make sound decisions, it is necessary to meaningfully enrich specialized reporting documents, improve information support, and refine methods for analyzing PHC activities based on a systems approach and final results. This is because management decisions depend on the reliability and completeness of the information received in the information support and on high-quality analysis at every stage. All this requires the creation of an effective cost management system and ensuring control over the rational use of available resources.

Scientists worldwide are conducting research into optimizing medical records in primary healthcare. Therefore, research aimed at analyzing the operations of family clinics, assessing the performance, quality, and effectiveness of general practitioners, the reliability and quality of existing primary records in family clinics, the reliability of receiving and processing statistical reports, and the development and implementation of an electronic program for family clinics designed to improve and optimize the process of maintaining primary records on paper and the transition to an electronic system are becoming increasingly important.

Objective

To optimize medical and statistical accounting and reporting documents maintained in outpatient and polyclinic institutions, as well as to develop an electronic program aimed at improving their efficiency and information content.

Material and methods

Epidemiological, social-hygienic, sociological, time-keeping, and sanitary-statistical methods were used to study record-keeping in family clinics and the activities of general practitioners (GPs).

The practical results of the study are as follows:

- As a result of a comprehensive study and structural and functional analysis of the current system for registering medical accounting and reporting documentation, the unofficial forms of numerous medical records and registration journals used in primary health care institutions of the urban district, as well as the duplication of identical information in other documents, were assessed.
- The requirements for the correct registration of medical accounting and reporting documentation were assessed, eliminating the deficiencies identified during the structural and functional analysis of medical accounting documentation in primary health care institutions.
- A modernized package of basic medical documents for registering medical accounting and reporting documentation in primary health care institutions has reduced the total number of medical records, registration logs, and forms.

The scientific significance of the study's results is explained by the fact that requirements for the proper registration of medical accounting and reporting documentation made it possible to eliminate deficiencies through structural and functional analysis, while a modernized package of essential medical documents in primary care facilities resulted in a reduction in the total number of charts, journals, and forms. By implementing the developed electronic program for optimizing medical accounting in primary care, the number of paper documents completed by medical workers and the associated costs were reduced by 1.4 times.

Based on the data obtained, a program was developed to optimize accounting documentation and automate the report on Form No. 12 “Report on diseases registered among the population living in the service area of a medical and preventive institution” (Fig. 1).



Fig. 1. Algorithm of the automated electronic program for reporting diseases registered among the population living in the service area of a medical and preventive institution

Sequence of work:

1. The patient comes to see the doctor, the doctor registers him in the “Journal of daily doctor’s appointments” (form 024-1 y), after which the data is processed, and the entry about the visit is automatically transferred to the “Record of visits to doctors in outpatient and polyclinic institutions and at home” (form 039 y).
2. Then the doctor opens the electronic outpatient card “Outpatient Medical Card” (form 025 y) and begins to record the reason for the visit.
3. When a patient presents with a medical condition, the physician records the diagnosis on the "Registration Sheets of Final (Revised) Diagnoses," adding the ICD-11 disease code and a "+" or "-" sign. After saving this information, the record is automatically transferred to the "Statistical Form for Registration of Final (Revised) Diagnoses" (form 025-1 y).
4. After this, this registration is processed, and the outpatient observation log is filled automatically. Registration data is automatically entered into this log.
5. At the next stage, registration is automatically transferred to the “Control card for dispensary observation” (form 030 y), where, in accordance with the diagnosis of the patient taken under dispensary observation, the time of the next examination is determined.

As a result, doctors using this program save time on filling out paper forms and can devote more time to examining patients and asking about the causes of their illness. Ultimately, each doctor receives a complete visit report and Form No. 12 (morbidity report).

Implementation of the developed program improves the efficiency, informativeness, and reliability of statistical reports. The following results have been achieved: improved medical records; increased physician efficiency; increased reliability of statistical reports; and a reduction in the time spent completing medical records. Using the computer program, general practitioners are able to fully analyze data and improve the treatment and diagnostic process.

CONCLUSION

The implementation of e-health is not just a technical upgrade, but a fundamental shift toward patient-centered medicine. Despite the challenges associated with cybersecurity and the need for staff training, the benefits in terms of saved lives and cost-effectiveness make this path unbeatable for modern society.

Automating accounting and reporting in primary healthcare is crucial because accounting and reporting constitute a crucial part of a healthcare organization's operations, both in terms of achieving organizational goals and in terms of labor costs. The rapid processing of primary medical documentation using information technology enables the acquisition of reliable information, which, in turn, helps reduce the extensive workload associated with completing documents and the generation of records and reports on completed examinations and tests. The cost-effective implementation of computer programs has significantly increased patient interaction time, ensured a more efficient physician workflow, and produced reliable statistical reports. At the same time, paperwork costs have been reduced, and wait times in doctor's appointments have been shortened.

References

1. Ashurova M.Zh. , Mamatkhanova G.M., Ashurova M.D. New prospects for maintaining accounting and reporting medical documentation in primary health care institutions. "Biomedicine va amaliyot " magazines 4 sleep, 5 veins B.262-267
2. Ismailov S.I., Mamatkhanova G.M. Electronic document management as a key factor in improving the efficiency of healthcare management. Eurasian journal of medical and natural sciences, 2(8), pp. 38–45. https://doi.org/10.5281/zenodo.6976038 (14.00.00. No.)
3. Mamatkhanova G.M., Ashurova M.Zh. , Ashurova M.D. Comprehensive assessment of the current electronic database of primary accounting and reporting medical documentation in primary health care institutions. Economy and Society. 2020, No. 2 (69), pp. 506–512
4. Mamatxanova GM, Ismailov SI Optimization of medical records and implementation of electronic systems in healthcare . The American Journal of Medical Sciences and Pharmaceutical Research , 2021/3 , P.193-198 .
5. Mamatkhanova G.M., Ismailov S.I. Analysis of the quality of medical records and ways to improve them in primary health care institutions. Bulletin of the Association of Doctors of Uzbekistan scientific and practical medical journal 2021/4. From 17-20
6. Mamatkhanova G.M. Optimization of medical records and implementation of electronic systems in healthcare. Jamiyat va innovationalar – Society And Innovations – Society and Innovations Journal home page: https://inscience.uz/index.php/socinov/index 2021. 15.09. pp . 61-67
7. G. M. Mamatkhanova. Current issues of creating an automated system in primary health care institutions. Journal of clinical and preventive medicine. No. 3, 2023, ISSN 2181-3531, www.fjsti.uz P.79-87 .
8. Mamatxanova GM Hujjatlarni legal avtomatlashtirilgan tizimlari va ularni amaliyotga joriy ethish . O ' Zbekiston harbiy tibbiyoti (5) 2024 Yil 227-233 b.
9. Mamathanova Gulnora Maxmudovna . Aholi salomatligini baholashda statistik usullarni takomillashtirishning ahamiyati . ActaCAMU . №2 (10.1) 2025 . 305-309 b.
10. Mamatkhanova GM Current issues of creation automated system for completion of reporting based on accounting medical documentation. Journal of clinical and preventive medicine . №3 2025 y ISSN 2181-3531 www.fjsti.uz 62-65 p.

ACUTE CARDIOVASCULAR DISEASES IN THE PRIMARY HEALTH CARE SYSTEM

Siddikov Bobirjon Tursinali ugli
Fergana Regional Health Department

ABSTRACT

Cardiovascular diseases (CVD) are one of the leading causes of death worldwide, and according to WHO estimates, 31% of deaths worldwide are due to CVD. 85% of these deaths are due to myocardial infarction (MI) and stroke. In the Republic of Uzbekistan, this problem is even more acute, and according to the 2022 report of the Ministry of Health, 55.4% of all deaths are associated with acute CVD.

Unfortunately, acute respiratory infections are becoming increasingly common and are becoming the leading cause of death for people.

Myocardial infarction and acute cerebrovascular disease (stroke) are the leading causes of disability among the working-age population. Research shows that more than 80% of these diseases are caused by modifiable risk factors, making primary prevention a priority (Roth et al, 2020; WHO, 2023).

Family doctors and medical workers working in the primary health care (PHC) system are the most effective team for early identification of risk factors, prevention of disease progression, and orientation of the population towards a healthy lifestyle (Mirzayev et al., 2021).

Keywords: Sharp cardiovascular diseases, prevention, screening, SCORE2, family doctor, primary health care, arterial hypertension.

Introduction

Although premature mortality from cardiovascular diseases remains high in low-and middle-income countries, these diseases can be prevented and their complications significantly reduced through healthy lifestyles, physical activity, and early identification and management of modifiable risk factors (obesity, smoking, hypertension).

Research objective: To systematize evidence-based information and develop practical recommendations for improving acute UTI prevention in the primary health care system based on the results of a systematic meta-analysis of literature published in 2015-2024.

Research method. A systematic review was conducted based on scientific sources published between 2015 and 2024. The search was conducted in PubMed, Scopus, Web of Science, Cochrane Library, and ZiyoNET databases.

The selection criteria included RCTs, meta-analyses, systematic reviews, and national and international clinical guidelines (in English, Russian, and Uzbek). Animal studies, old studies, or studies for which the full text was not available were excluded.

Initially, 84 sources were identified, and after screening, the remaining 62 were analyzed in depth. The level of evidence was assessed based on the GRADE criteria.

Table 1.

Acute HIV prophylaxis in the NHS : evidence summary

Prevention of the direction	Main literary source/evidence	Risk reduction
Arterial hypertension screening and treatment	SPRINT, 2015; ESC/ESH, 2023; Ettehad et al., 2016	Stroke: −35–40%; KYuC: −20–25%
Lipid prophylaxis with statins	CTT Collaboration, 2019; NICE, 2023; Grundy, 2019	Major events: -22–35%
Smoking cessation program	Cahill et al., 2021; WHO 2023	Risk of KYuK: -50%
Mediterranean diet and nutrition	PREDIMED, 2018; Piepoli et al., 2016	Cardiovascular events: −30%
Physical activity (150 min/week)	Arnett et al., 2019; AHA 2022	Overall risk: -20–30%
Diabetes screening and control	ADA 2023; ACCORD 2021	Microvascular complications: −25%
Multidisciplinary team model	Bodenheimer et al., 2019; Shi, 2023	Program efficiency: +35–40%
EHR/CDSS implementation	Shojania et al., 2020; Tucker, 2021	Protocol compliance: +30–40%

Uzbekistan (Egamberdieva et al, 2020) revealed the following indicators: **Arterial hypertension:** 41.8%; **Smoking (among men):** 31.6%; **Overweight:** 38.4%; **Physical inactivity:** 34.2%

These identified indicators are significantly higher than the average for European countries, which proves the need to further strengthen preventive measures.

Arterial hypertension screening from 18 years old all patients for periodic accordingly blood pressure size the most economic fruit method (NNT = 150 patients over 5 years) observation, 1 stroke prevent ESC / ESH (2023) guidelines ≥140 / 90 mmHg mezzanine defines.

Lipid profile screening is recommended every 5 years starting at age 40. Integrating the SCORE2 calculator into BTSYoga increases the likelihood of correctly identifying the risk level by 2.5 times (Bjarnadottir et al, 2020).

Diabetes screening should be performed by HbA1c or fasting glucose in all overweight individuals over the age of 45, according to the ADA (2023).

is recommended in NICE (2023) and ESC (2021) guidelines.

Antihypertensive therapy. The SPRINT trial (2015) showed that lowering systolic blood pressure to <120 mmHg can reduce the risk of cardiovascular events by 25% . However, the ESC/ESH (2023) recommends a target of <130/80 mmHg as a balanced option.

Aspirin. The ASPREE (2018), ARRIVE (2018), and ASCEND (2018) trials did not show that the benefits of aspirin in primary prevention outweigh the risk of bleeding. The AHA/ACC (2019) recommends considering it only in high-risk, low-risk patients under 70 years of age

Tobacco cessation: Integrated smoking cessation clinics + pharmacotherapy (varenicline) increase 1-year abstinence rates by 25–35% (Cahill, 2021). This is the most cost-effective approach.

Strict adherence to a balanced diet: The PREDIMED study (2018) demonstrated that a Mediterranean diet can reduce cardiovascular events by 30%. The importance of recommendations adapted to local dietary traditions is emphasized.

Physical activity: ≥150 minutes of moderate-intensity physical activity per week reduces overall cardiovascular risk by 20–30% (Arnett et al, 2019).

A multidisciplinary team (family doctor + nurse + pharmacist + psychologist) increases efficiency by 35–40%. Electronic medical records (EHR) and clinical decision support systems (CDSS) improve compliance with protocols by 30–40%. Remote monitoring (telemonitoring) increases adherence to antihypertensive therapy (taking medications on time, following the diet prescribed by the doctor, coming to examinations on time) by 25–30%.

Successful international experiences show that the North Karelia Project in Finland has reduced the mortality rate from acute UTI by 82% over 30 years. The Quality and Outcomes Framework system has also significantly improved the effectiveness of clinical practice by increasing financial support for preventive indicators in the UK.

Conclusion

It is an urgent task to significantly reduce the complications of acute cardiovascular diseases and increase the effectiveness of prevention through early identification of risk factors, expansion of screening programs, and the use of modern technologies in primary health care.

LITERATURE

1. Global Burden of Cardiovascular Diseases and Risk Factors, 1990–2019: JACC Scientific Expert Panel / GA Roth [et al.] // Journal of the American College of Cardiology. - 2020. - Vol. 76, No. 25. – P. 2982–3021.

2. Cardiovascular diseases (CVDs): Fact sheet [Electronic resource] / World Health Organization. – 2023. – URL: https://www.who.int/news-room/fact sheets/detail/cardiovascular-diseases-(cvds (date of access: 04/13/2026).

3. 2021 ESC Guidelines on cardiovascular disease prevention in clinical practice / FL Visseren [et al.] // European Heart Journal. - 2021. - Vol. 42, No. 34. – P. 3227–3337.

4. 2023 ESH Guidelines for the management of arterial hypertension / G. Mancia [et al.] // Journal of Hypertension. - 2023. - Vol. 41, No. 12. – P. 1874–2071.

5. A Randomized Trial of Intensive versus Standard Blood-Pressure Control / SPRINT Research Group // The New England Journal of Medicine. - 2015. - Vol. 373, No. 22. – P. 2103–2116.

6. Efficacy and safety of statin therapy in older people: a meta-analysis of individual participant data from 28 randomized controlled trials / Cholesterol Treatment Trialists' Collaboration // The Lancet. - 2019. - Vol. 393, No. 10170. – P. 407–415.

7. Primary Prevention of Cardiovascular Disease: A Review / DK Arnett [et al.] // JAMA. - 2019. - Vol. 322, No. 8. - P. 756–767. (Note: this is an updated reference to the primary ACC/AHA recommendations).

8. Standards of Care in Diabetes—2023 / American Diabetes Association // Diabetes Care. - 2023. - Vol. 46, Suppl. 1. – P. S1–S291.

9. Primary Prevention of Cardiovascular Disease in Primary Care: A Systematic Review / Y. Shi [et al.] // Preventive Medicine. - 2023. - Vol. 168. – Art. 107409.

10. Effect of Aspirin on Cardiovascular Events and Bleeding in the Healthy Elderly / JJ McNeil [et al.] // The New England Journal of Medicine. - 2018. - Vol. 379, No. 16. – P. 1509–1518.

11. Mirzayev, OA Improving the indicators of the UQTC and the BTSY in Uzbekistan / OA Mirzayev, JH Abdullayev // Uzbekistan Medical Journal. – 2021. – No. 4. – P. 45–52.

12. Egamberdieva, DM STEPS study: prevalence of risk factors in Uzbekistan / DM Egamberdieva // Uzbekistan Medical Journal. – 2020. – No. 2. – P. 14–22.

ORGAN-PRESERVING STRATEGY AND PREVENTION OF HYSTERECTOMY IN PLACENTA PREVIA IN WOMEN WITH A UTERINE SCAR

Yulduz Djabbarova Kasimovna
Doctor of Medical Sciences, Academician of the Academy of Medical and Technical Sciences of the Russian Federation, , Professor of the Department of Obstetrics and Gynecology, Reproductology of the Tashkent State Medical University, Professor of the Department of Obstetrics and Gynecology,Neonatology of the Central Asian Medical University, Scientific Supervisor:
[1]ORCID ID:https://orcid.org/0000-0002-3456-2381

Maqsudova Barno Jaxongirovna
Master's degree student of the Department of Obstetrics and Gynecology, International Medical University, Central Asian Medical University, Student
[2]ORCID ID:https://orcid.org/0009-0005-8888-4190

ABSTRACT
Placenta previa in women with a uterine scar remains a significant obstetric challenge due to the high risk of massive hemorrhage and the frequent need for hysterectomy. This article explores organ-preserving strategies aimed at reducing maternal morbidity and preserving reproductive potential. The study analyzes modern diagnostic approaches, including prenatal imaging, as well as multidisciplinary management tactics during delivery. Special attention is given to surgical and non-surgical techniques such as uterine-sparing procedures, hemostatic interventions, and individualized patient care. Preventive measures that minimize the likelihood of hysterectomy are also discussed. The findings highlight that timely diagnosis, careful planning, and the use of advanced medical technologies can significantly improve maternal outcomes and reduce the necessity for radical surgical interventions.
Keywords: placenta previa, uterine scar, organ-preserving strategy, hysterectomy prevention, obstetric hemorrhage, maternal outcomes, uterine-sparing techniques, prenatal diagnosis.

АННОТАЦИЯ
Предлежание плаценты у женщин с рубцом на матке представляет собой серьёзную акушерскую проблему из-за высокого риска массивного кровотечения и частой необходимости проведения гистерэктомии. В данной статье рассматриваются органосохраняющие стратегии, направленные на снижение материнской заболеваемости и сохранение репродуктивной функции. Анализируются современные методы диагностики, включая пренатальную визуализацию, а также междисциплинарные подходы к ведению родов. Особое внимание уделяется хирургическим и нехирургическим методам, таким как органосохраняющие операции на матке, гемостатические вмешательства и индивидуализированное ведение пациенток. Также рассматриваются профилактические меры, позволяющие снизить вероятность проведения гистерэктомии. Полученные результаты показывают, что своевременная диагностика, тщательное планирование и использование современных медицинских технологий способны значительно улучшить материнские исходы и снизить необходимость радикальных хирургических вмешательств.
Ключевые слова: предлежание плаценты, рубец на матке, органосохраняющая тактика, профилактика гистерэктомии, акушерское кровотечение, материнские исходы, органосохраняющие методы, пренатальная диагностика.

ВВЕДЕНИЕ

Предлежание плаценты является одной из наиболее актуальных и сложных проблем современной акушерской практики, поскольку напрямую связано с высоким риском материнской и перинатальной заболеваемости и смертности. Особую клиническую значимость данная патология приобретает у женщин с рубцом на матке, сформированным в результате предыдущих кесаревых сечений или других хирургических вмешательств. В последние десятилетия наблюдается устойчивая тенденция к увеличению частоты операций кесарева сечения, что, в свою очередь, приводит к росту числа пациенток с рубцовыми изменениями миометрия. Это обстоятельство существенно повышает вероятность аномалий плацентации, включая предлежание плаценты и патологическое врастание плаценты, такие как placenta accreta spectrum.Предлежание плаценты у данной категории женщин нередко сопровождается массивными акушерскими кровотечениями, представляющими непосредственную угрозу для жизни матери. Традиционной тактикой в условиях неконтролируемого кровотечения зачастую становится гистерэктомия, которая, несмотря на свою эффективность в плане остановки кровопотери, приводит к полной утрате репродуктивной функции и может оказывать значительное психологическое и социальное воздействие на женщину. В связи с этим особую актуальность приобретает разработка и внедрение органосохраняющих подходов, направленных на минимизацию хирургической агрессии и сохранение матки.Современные достижения медицины позволяют пересмотреть традиционные подходы к ведению беременных с предлежанием плаценты и рубцом на матке. Важную роль играет ранняя и точная диагностика с использованием ультразвукового исследования и магнитно-резонансной томографии, что дает возможность своевременно выявлять степень инвазии плаценты и планировать оптимальную тактику родоразрешения. Комплексный междисциплинарный подход с участием акушеров-гинекологов, анестезиологов-реаниматологов и хирургов позволяет существенно повысить безопасность родоразрешения и снизить риск осложнений. Особое внимание в последние годы уделяется разработке органосохраняющих хирургических методов, таких как компрессионные швы, перевязка маточных и внутренних подвздошных артерий, эмболизация сосудов, а также использование современных гемостатических средств. Эти методы направлены на эффективный контроль кровотечения при сохранении анатомической и функциональной целостности матки. Кроме того, важную роль играют профилактические мероприятия, включая рациональное ведение предыдущих родов, снижение необоснованного числа кесаревых сечений и тщательное наблюдение за беременными из группы высокого риска.Таким образом, проблема органосохраняющей тактики при предлежании плаценты у женщин с рубцом на матке является многогранной и требует дальнейшего изучения. Разработка и внедрение эффективных профилактических и лечебных стратегий позволит не только снизить частоту гистерэктомий, но и улучшить качество жизни пациенток, сохранив их репродуктивный потенциал.

МАТЕРИАЛЫ И МЕТОДЫ

Настоящее исследование выполнено с целью оценки эффективности органосохраняющей тактики и профилактики гистерэктомии при предлежании плаценты у женщин с рубцом на матке. Работа носит клинико-аналитический характер и основана на комплексном изучении течения беременности, родоразрешения и послеродового периода у пациенток данной группы риска.В исследование были включены женщины с диагностированным предлежанием плаценты, имеющие в анамнезе один или более

рубцов на матке после кесарева сечения или других оперативных вмешательств. Критериями включения являлись: подтверждённое ультразвуковыми методами предлежание плаценты, наличие рубца на матке, одноплодная беременность, а также информированное согласие пациенток на участие в исследовании. Критериями исключения служили тяжелые сопутствующие соматические заболевания, врождённые аномалии развития плода, а также случаи, требующие экстренного оперативного вмешательства без возможности применения органосохраняющей тактики.Обследование пациенток проводилось в несколько этапов и включало клинические, лабораторные и инструментальные методы исследования. На этапе пренатальной диагностики всем женщинам выполнялось ультразвуковое исследование с допплерометрией для оценки локализации плаценты, степени её инвазии в миометрий и состояния маточно-плацентарного кровотока. При подозрении на патологическое врастание плаценты дополнительно применялась магнитно-резонансная томография, позволяющая уточнить глубину и распространённость инвазии.Особое внимание уделялось сбору анамнеза, включая количество предыдущих беременностей, родов, операций кесарева сечения, интервал между беременностями, а также наличие осложнений в предыдущих гестациях. Оценивались клинические показатели состояния пациенток, включая уровень гемоглобина, показатели свертывающей системы крови, а также наличие признаков анемии и других сопутствующих состояний.Тактика ведения беременности и родоразрешения разрабатывалась индивидуально для каждой пациентки с учётом степени риска кровотечения, локализации плаценты и предполагаемой степени её врастания. В плановом порядке родоразрешение осуществлялось путём кесарева сечения в условиях специализированного стационара, оснащённого необходимым оборудованием и возможностями для оказания экстренной помощи. В операционной бригаде участвовали акушеры-гинекологи, анестезиологи-реаниматологи и, при необходимости, сосудистые хирурги.В рамках органосохраняющей тактики применялись различные методы контроля кровотечения. К ним относились наложение компрессионных швов на матку, перевязка маточных артерий, временная окклюзия сосудов, а также использование современных гемостатических препаратов. В ряде случаев применялась эндоваскулярная эмболизация маточных артерий как до, так и после оперативного вмешательства. Объём кровопотери оценивался интраоперационно и в раннем послеоперационном периоде.Эффективность проведённых мероприятий оценивалась по ряду критериев, включая объём кровопотери, необходимость переливания компонентов крови, частоту выполнения гистерэктомии, а также показатели послеоперационного восстановления. Дополнительно анализировались длительность госпитализации, частота осложнений и общее состояние пациенток в послеродовом периоде.Статистическая обработка полученных данных проводилась с использованием общепринятых методов медицинской статистики. Рассчитывались средние значения показателей, стандартные отклонения, а также проводился сравнительный анализ результатов в зависимости от применяемой тактики лечения. Достоверность различий оценивалась с использованием соответствующих критериев, что позволило сделать обоснованные выводы об эффективности органосохраняющего подхода.

РЕЗУЛЬТАТЫ

В ходе проведённого исследования были получены данные, свидетельствующие о высокой клинической значимости органосохраняющей тактики при ведении пациенток с предлежанием плаценты и рубцом на матке. Анализ течения беременности и исходов родоразрешения показал, что применение комплексного индивидуализированного

подхода позволяет существенно снизить частоту радикальных хирургических вмешательств, в частности гистерэктомии.У большинства пациенток предлежание плаценты было диагностировано на ранних сроках беременности благодаря своевременному проведению ультразвукового исследования. Это позволило заранее определить группу высокого риска и разработать оптимальную тактику ведения беременности и родоразрешения. В ряде случаев было выявлено сочетание предлежания плаценты с признаками её патологического врастания, что требовало более тщательной подготовки и участия мультидисциплинарной команды специалистов.Применение органосохраняющих методов в процессе кесарева сечения продемонстрировало высокую эффективность в контроле акушерского кровотечения. Использование компрессионных швов, перевязки маточных артерий и современных гемостатических средств позволило значительно уменьшить объём интраоперационной кровопотери по сравнению с традиционными подходами. У значительной части пациенток удалось избежать массивных кровотечений, требующих экстренного удаления матки.Частота выполнения гистерэктомии в исследуемой группе была существенно ниже по сравнению с ожидаемыми показателями у данной категории пациенток. В тех случаях, когда органосохраняющая тактика оказывалась недостаточно эффективной и сохранялась угроза жизни пациентки, своевременно принималось решение о выполнении гистерэктомии как жизнеспасающей меры. Однако общее количество таких случаев было ограниченным. Дополнительно было отмечено снижение потребности в массивной гемотрансфузии. Уровень послеоперационной анемии у пациенток, которым были применены органосохраняющие методы, был менее выражен, что способствовало более быстрому восстановлению в послеродовом периоде. Средняя продолжительность госпитализации также была сокращена, а частота послеоперационных осложнений, включая инфекционные и тромбоэмболические, оставалась на относительно низком уровне.



ОБСУЖДЕНИЕ

Полученные результаты подтверждают актуальность и клиническую значимость применения органосохраняющей тактики у пациенток с предлежанием плаценты на фоне рубца на матке. Данная категория беременных традиционно рассматривается как группа высокого риска развития массивных акушерских кровотечений, что нередко приводит к необходимости выполнения гистерэктомии как жизнеспасающего вмешательства. Однако современные подходы к диагностике и лечению позволяют существенно пересмотреть эту стратегию в сторону сохранения репродуктивного органа.Снижение частоты гистерэктомии, выявленное в ходе исследования, свидетельствует о высокой эффективности комплексного подхода, включающего раннюю диагностику, тщательное планирование родоразрешения и использование современных хирургических и нехирургических методов гемостаза. Особое значение имеет пренатальная идентификация пациенток с высоким риском патологического врастания плаценты, что позволяет заранее подготовить необходимую тактику ведения и минимизировать интраоперационные осложнения.Сравнение полученных данных с результатами других исследований показывает, что внедрение органосохраняющих технологий, таких как компрессионные швы, перевязка сосудов и эндоваскулярные методы, способствует значительному снижению объёма кровопотери и уменьшению необходимости в массивных трансфузиях. Это, в свою очередь, положительно сказывается на послеоперационном восстановлении и снижает риск развития осложнений.Следует отметить, что успех органосохраняющей тактики во многом зависит от уровня подготовки медицинского персонала и технической оснащённости лечебного учреждения. Проведение подобных операций требует высокой квалификации хирургов, наличия современного оборудования и возможности привлечения специалистов смежных профилей, включая анестезиологов-реаниматологов и сосудистых хирургов. Междисциплинарный подход является ключевым фактором в обеспечении безопасности пациенток.

ВЫВОДЫ

Проведённое исследование позволяет сделать вывод о том, что предлежание плаценты у женщин с рубцом на матке остаётся серьёзной акушерской проблемой, сопряжённой с высоким риском массивных кровотечений и неблагоприятных материнских исходов. Однако внедрение современных органосохраняющих подходов существенно расширяет возможности безопасного ведения данной категории пациенток.Установлено, что ранняя и точная пренатальная диагностика играет ключевую роль в снижении риска осложнений, позволяя своевременно выявлять степень аномалии плацентации и планировать оптимальную тактику родоразрешения. Индивидуализированный подход к каждой пациентке с учётом клинических факторов риска способствует повышению эффективности лечения и снижению частоты экстренных вмешательств.Применение органосохраняющих методов, включая компрессионные швы, перевязку сосудов и использование современных гемостатических средств, показало высокую эффективность в контроле акушерских кровотечений. Это позволило значительно снизить частоту выполнения гистерэктомии и сохранить репродуктивную функцию у большинства пациенток.Важным фактором успешной реализации органосохраняющей тактики является мультидисциплинарный подход, включающий участие квалифицированных специалистов и проведение родоразрешения в условиях специализированных медицинских учреждений. Наличие необходимого оборудования и

готовность к оказанию экстренной помощи существенно повышают безопасность пациенток.

ИСПОЛЬЗОВАННАЯ ЛИТЕРАТУРА

1. Jauniaux E., Chantraine F., Silver R.M., Langhoff-Roos J. *FIGO classification for the clinical diagnosis of placenta accreta spectrum disorders.* International Journal of Gynecology & Obstetrics, 2019; 146: 20–24.
2. Silver R.M., Landon M.B., Rouse D.J., Leveno K.J., Spong C.Y., Thom E.A., et al. *Maternal morbidity associated with multiple repeat cesarean deliveries.* Obstetrics & Gynecology, 2006; 107(6): 1226–1232.
3. Eller A.G., Porter T.F., Soisson P., Silver R.M. *Optimal management strategies for placenta accreta.* BJOG, 2009; 116: 648–654.
4. Sentilhes L., Kayem G., Chandraharan E., Palacios-Jaraquemada J., Jauniaux E. *Conservative management of placenta accreta spectrum.* International Journal of Gynecology & Obstetrics, 2018; 140: 37–44.
5. Jauniaux E., Collins S., Burton G.J. *Placenta accreta spectrum: Pathophysiology and evidence-based clinical management.* Best Practice & Research Clinical Obstetrics & Gynaecology, 2017; 38: 1–13.
6. Silver R.M. *Abnormal Placentation: Placenta Previa, Placenta Accreta, and Vasa Previa.* Obstetrics & Gynecology, 2015; 126: 654–668.
7. Eller A.G., Bennett M., Sharshiner M., Soisson P., Treadwell M.C., Porter T.F., Silver R.M. *Maternal morbidity in cases of placenta accreta managed conservatively.* American Journal of Obstetrics & Gynecology, 2011; 204(1): 63.e1–63.e5.
8. Chandraharan E., Arulkumaran S. *Conservative management of placenta accreta: Surgical techniques and outcomes.* Best Practice & Research Clinical Obstetrics & Gynaecology, 2012; 26: 685–701.
9. Royal College of Obstetricians and Gynaecologists (RCOG). *Placenta praevia and placenta accreta: Diagnosis and management.* Green-top Guideline No. 27, 2018.
10. Timor-Tritsch I.E., Monteagudo A., Cali G., Elimian A., Palacios-Jaraquemada J., Kaelin Agten A. *Prenatal identification and imaging of placenta accreta spectrum.* American Journal of Obstetrics & Gynecology, 2014; 211(2): 145–157.

ASSESSMENT OF THE ANTIOXIDANT ACTIVITY OF THE DRUG "RUTAN" IN ACUTE TOXIC HEPATITIS AT THE PREPUBERTAL AGE

Senior assistant, Fatima Adilbaevna Khalmuratova[1]
[1]Karakalpakstan Medical Institute, Nukus, Karakalpakstan, Uzbekistan

Doctor of Medical Sciences, Professor Ziyaviddin Zainutdinovich Hakimov
[2]Department of Pharmacology, Tashkent State Medical University , Tashkent, Uzbekistan
[2]ORCID ID: https://orcid.org/0000-0002-7404-3447

Doctor of Medical Sciences, Professor Alisher Khudayberdievich Rakhmanov[3]
[3]Institute of Medical, Scientific and Innovative Research at the Tashkent State Medical University, Tashkent, Uzbekistan
[3]ORCID ID: https://orcid.org/0000-0002-7404-3447

Relevance

Toxic liver injuries occupy a significant place in the structure of diseases in children and adolescents. During puberty, the incidence of toxic hepatitis increases due to expanded exposure to xenobiotics, including medications (NSAIDs and antibiotics), dietary supplements, industrial toxins, and alcohol. Functional instability of the antioxidant system (AOS) in adolescence is associated with neuroendocrine changes, possible micronutrient deficiency, and increased energy demands. This determines the need to search for effective drugs with antioxidant properties for inclusion in the comprehensive therapy of toxic hepatitis in adolescents.

Objective

To comparatively evaluate the effects of the drugs "Rutan" and "Karsil" on lipid peroxidation (LPO) processes and the activity of antioxidant system enzymes in acute toxic hepatitis (ATH) in prepubertal animals.

Materials and Methods

The experiment was performed on 30 one-month-old white rats. The ATH model was induced by intragastric administration of a 50% oil solution of carbon tetrachloride (CCl_4) at a dose of 0.2 ml/100 g body weight for 4 days. One day after the last administration, the animals were divided into groups: the first received "Rutan" (25 mg/kg), the second — "Karsil" (40 mg/kg), and the third served as an untreated control. The drugs were administered once daily for 6 days. In the microsomal-cytosolic liver fraction, the levels of LPO products (acyl hydroperoxides — AHP and malondialdehyde — MDA) and the activity of AOS enzymes (catalase — CAT and superoxide dismutase — SOD) were determined.

Results

The formation of free radicals that damage biological membranes, including hepatocytes, is a key pathogenetic mechanism of hepatotoxic injury. Therefore, the levels of AHP and MDA were studied in prepubertal animals with ATH treated with Rutan and Karsil. In animals with ATH, a marked intensification of LPO processes was observed: AHP increased more than

4.5-fold and MDA 3.7-fold compared with intact animals. Experimental therapy with "Rutan" (25 mg/kg) reduced AHP by 68.6% and MDA by 64.0% relative to untreated animals. Administration of "Karsil" also led to a pronounced decrease in LPO parameters, comparable to the effect of the reference drug "Rutan". The content of LPO products depends on the activity of AOS enzymes. The development of ATH was accompanied by a decrease in AOS enzyme activity: after 6 days, CAT and SOD activities decreased by 64.9% and 60.7%, respectively, compared with healthy animals. Against the background of "Rutan" therapy, a statistically significant increase in CAT (by 131.9%) and SOD (by 122.1%) activity was observed compared with untreated animals. "Karsil" showed a similar effect: CAT activity increased by 140.4% and SOD by 111.6%.

Conclusions

Experimental therapy of ATH with "Rutan" effectively reduces the intensity of free-radical lipid oxidation and increases the activity of liver antioxidant enzymes. In terms of the severity of its antioxidant effect, "Rutan" is not inferior to "Karsil" and can be considered a promising agent in the comprehensive therapy of toxic hepatitis in adolescents.

CONTEMPORARY APPROACHES TO THE LABORATORY ETIOLOGICAL DIAGNOSIS OF FOODBORNE POISONING AND ISSUES OF THEIR EPIDEMIOLOGICAL SURVEILLANCE IN UZBEKISTAN

[1]Atabekov N.S., Doctor of Medical Sciences;
https://orcid.org/0009-0008-5914-5200

[2]Shadmanova N.A., Doctor of Medical Sciences,
https://orcid.org/0009-0005-2610-4021
[1,2]Committee for Sanitary-Epidemiological Welfare and Public Health of the Republic of Uzbekistan.

Associate Professor, [3]Iskandarova G.T., Doctor of Medical Sciences,
https://ORCID.org/0009-0000-0537-0818

Professor; [4]Asatova N.B., Candidate of Medical Sciences, Associate Professor;
https://orcid.org/0009-0000-7154-752X

[5]Sapaeva F.R., Doctor of Biological Sciences, Associate Professor
https://orcid.org/my-orcid?orcid=0000-0002-9772-2020

[3,4,5]Center for Professional Development of Medical Workers under the Ministry of Health of the Republic of Uzbekistan.

Abstract
This article analyzes bacterial pathogens responsible for food poisoning and acute intestinal diseases to assess the reliability of microbiological investigations. The authors emphasize the importance of advancing molecular biology and actively implementing rapid methods of bacteriological analysis for effective epidemic control. The importance of developing regulatory standards for microbiological diagnostics, establishing protocols for clinical trials, and introducing modern diagnostic methods in local microbiological laboratories is also highlighted.
Keywords: food poisoning, toxins, genotype, retrospective analysis.

Acute infectious intestinal diseases, foodborne infections, and food poisoning are currently recognized as significant global public health challenges. According to the World Health Organization, approximately 600 million people worldwide—equivalent to one in every 10–12 individuals—seek medical attention each year due to various clinical manifestations resulting from the consumption of contaminated food products. Furthermore, these conditions are associated with an estimated 420,000 deaths annually.

At the same time, it is noted that among these cases, approximately 125,000 occur in children under the age of five, with the development of severe diarrheal syndrome accounting for about 43%. In most instances, contamination of food products arises due to biological factors, with approximately 70% of cases associated with various microorganisms and, in part, their exo- or endotoxins [4;6].

It is evident that significant and responsible tasks lie ahead for the healthcare system, particularly for sanitary and epidemiological services, in preventing and at least partially reducing such cases.

The investigation of food poisoning cases requires the application of modern laboratory methods in practice, enabling the monitoring of particularly hazardous variants (genotypes) of microorganisms that act as etiological agents of the disease, and necessitating the integration of epidemiological investigations with laboratory findings.

The above-mentioned issues remain highly relevant in the context of Uzbekistan as well. From this perspective, the speed of obtaining results in classical microbiological (bacteriological) diagnostic methods widely used in routine laboratory practice—particularly the duration of analysis—is of great importance for epidemiologists and clinicians. In practice, there is often a need to accelerate the diagnostic process in order to ensure timely diagnosis and to determine appropriate treatment strategies.

Objective of the study: To assess the effectiveness of microbiological investigations aimed at identifying the etiological agents of food poisoning and acute infectious intestinal diseases.

Materials and Methods

A retrospective analysis was conducted based on annual reports of the Epidemiology Department and the Bacteriology Laboratory of the Committee for Sanitary and Epidemiological Well-being and Public Health under the Ministry of Health. The study also included the examination of pure isolates of bacterial pathogens preserved in the bacteriological laboratory museum. These investigations involved the assessment of main biological characteristics using biochemical “color” test panels, as well as serological analysis using agglutinating sera. In addition, mass spectrometry analysis (MALDI-TOF MS) was applied. The study was carried out in accordance with the Ministry of Health of the Republic of Uzbekistan Order No. 177 (2015), WHO protocols, EUCAST guidelines (v.13.0; 2023), and other relevant standards.

Results

As of 2024, there are 213 bacteriological laboratories within the territorial departments of the Committee for Sanitary and Epidemiological Well-being and Public Health, of which 210 (98.6%) are operational. Microbiological/bacteriological diagnosis of patients with acute infectious intestinal diseases and food poisoning is primarily conducted in the laboratories of the Committee’s regional departments, while 17.4% of examinations are carried out in bacteriological laboratories within the regional branches (44 units) of the Republican Specialized Scientific-Practical Medical Center for Epidemiology, Microbiology, Infectious and Parasitic Diseases.

Patients with acute infectious intestinal diseases and food poisoning most often seek care at territorial treatment and prevention institutions, as well as infectious disease hospitals of the regional branches of the above-mentioned center. Biological samples are analyzed in the laboratories of these institutions, and the results of microbiological investigations serve as the basis for diagnosis. Bacterial pathogens isolated during analysis are identified as etiological agents of the disease and subsequently submitted to the regional bacteriology laboratories of the Committee for confirmation.

The first stage of this study was devoted to retrospective analysis. According to annual reports of the sanitary-epidemiological authorities, food poisoning cases occur throughout the year and demonstrate seasonal patterns, specific contributing factors, associated clinical symptoms, outbreak potential, and other characteristics. In particular, analysis of local data for 2021–2023 indicates a partial increase in the incidence of acute infectious intestinal

diseases and food poisoning in recent years (Figure 1). However, a significant decrease in mortality among patients suggests that timely and rapid response measures implemented by the Committee for Sanitary and Epidemiological Well-being and Public Health and the regional branches of the specialized center have been effective.



Figure 1. Food poisoning cases registered in the Republic of Uzbekistan during 2021–2023.

At the same time, based on the results of microbiological investigations of foodborne poisoning cases over the past three years, "botulism diagnosis" and "foodborne toxic infections" were identified as the main biological factors. In 2021, 71 cases were registered (176 affected individuals), in 2022 — 36 cases (82 individuals), and in 2023 — 39 cases (145 individuals). All recorded fatal outcomes were associated with a diagnosis of botulism, while the presented reports did not fully reflect comprehensive microbiological analyses of foodborne toxic infections.

As the second part of the retrospective analysis, annual reports from the bacteriology laboratories of the Committee for Sanitary and Epidemiological Welfare and Public Health were reviewed. In particular, it was found that a total of 283,505 microbiological examinations for acute intestinal infections were conducted during 2021–2023, with the majority of isolated microorganisms belonging to Gram-negative bacteria of the Enterobacteriaceae family. At the same time, analysis of year-by-year data demonstrated that most laboratories primarily focused on officially recognized pathogenic enterobacteria (Shigella spp., Salmonella spp., Escherichia coli, as well as conditionally pathogenic Enterobacteria (CPE)). (Figure 2.) Data concerning Gram-positive microorganisms were limited mainly to the isolation of pathogenic staphylococci, specifically *Staphylococcus aureus*. However, according to recent scientific literature, there are seven recognized species of coagulase-positive pathogenic staphylococci, including *S. aureus*, *S. delphini*, *S. hyicus*, *S. intermedius*, *S. lutrae*, *S. pseudintermedius*, and *S. schleiferi* subsp. *coagulans*. Among these microorganisms, particular attention is given to species capable of producing enterotoxins—key virulence factors—such as *S. hyicus*, *S. pseudintermedius*, *S. chromogenes*, *S. cohnii*, *S. epidermidis*, *S. lentus*, *S. lugdunensis*, *S. saprophyticus*, *S. sciuri*, *S. warneri*, and *S. xylosus*.



Figure 2. Etiological analysis of acute intestinal infectious diseases (2021–2023).

The proportion of positive results for pathogenic staphylococci across the years remained extremely low, accounting for 9.6% in 2021, 9.8% in 2022, and 9.6% in 2023. A similar pattern was observed for members of the Enterobacteriaceae family, with a clear trend toward overdiagnosis of conditionally pathogenic enterobacteria (CPE) (Table 1).

Table 1.
Etiological analysis of acute intestinal infectious diseases caused by members of the Enterobacteriaceae family (2021–2023).

Years	Total number of examinations	Number of isolated strains	Members of the Enterobacteriaceae family			
			Shigella spp.,	Salmonella spp.,	E. coli	CPE
2021	82296	5290	423	452	2	4413
2022	103172	6951	663	563	-	5656
2023	98037	5708	334	468	-	4906
Total	**283505**	**17949**	**1420**	**1483**	**2**	**14975**

The results of microbiological investigations of acute intestinal infectious disease (AIID) cases caused by Gram-negative intestinal pathogens over the past three years demonstrated an extremely low rate of etiological verification. Specifically, the detection rates were 6.4% in 2021, 6.7% in 2022, and 5.8% in 2023, with the majority of biological samples yielding only conditionally pathogenic enterobacteria (CPE). Errors were observed during laboratory investigations, particularly at the pre-analytical and analytical stages. In particular, regional bacteriological laboratories involved in AIID diagnostics are inadequately equipped

in terms of material and technical resources and lack the capacity to perform modern laboratory investigations. Weak control over the collection and transportation of biological samples, sampling after the initiation of antibiotic therapy, and failure to adhere to required sampling volumes and intervals have negatively affected the reliability of laboratory results.

It is well established that, in the study of pathogens such as *Escherichia coli* and *Staphylococcus aureus*, it is essential to determine the presence of toxins and toxin-associated genes. These indicators are critical for confirming the etiological role of the isolated pathogen in acute intestinal infections. However, despite the availability of more than 170 PCR (polymerase chain reaction) instruments within the system of the Committee for Sanitary and Epidemiological Welfare and Public Health, their utilization in practice remains limited, indicating that unresolved issues in this field continue to be highly relevant.

It should be noted that sufficient conditions have been established for the microbiological analysis of pathogens such as *Shigella* spp. and *Salmonella* spp., including the availability of specific diagnostic reagents such as agglutinating sera. However, comprehensive characterization of isolated strains—including their biological properties, antimicrobial susceptibility, and genetic features—is not routinely performed. This not only limits the generation of complete and reliable data on local strains but also often results in superficial rather than in-depth and comprehensive epidemiological analyses.

In 2023, a total of 92 *Salmonella* strains submitted to the bacteriology laboratory of the Committee for confirmation were studied in terms of their biochemical characteristics, serotypes, and antimicrobial susceptibility. Of these, 85 strains were confirmed as belonging to the genus *Salmonella*, comprising 12 serovars. Among them, 34 strains (40%) were identified as belonging to the *S. Enteritidis* subtype. This finding indicates that this particular serovar is widely распространён in the territory of the Republic of Uzbekistan, consistent with global trends (Figure 3).



Figure 3. Results of serotyping analysis of *Salmonella* spp. isolates obtained from acute intestinal infectious disease (AIID) cases in 2023.

At the next stage of the study, resistance of *Salmonella* spp. strains to 18 antimicrobial agents was investigated [1;2;3;5]. The obtained results demonstrated that local isolates were generally susceptible to β-lactam antibiotics, particularly cephalosporins, while exhibiting resistance to fluoroquinolones. This finding indicates the emergence of specific, epidemiologically significant, and potentially hazardous variants of *Salmonella* circulating

within the regions of the republic. At the same time, this situation has highlighted the need to establish a systematized database within the framework of the Committee for Sanitary and Epidemiological Welfare and Public Health and has provided a scientific and practical basis for further work in this direction (Figures 4 and 5).



Figure 4. Susceptibility of *Salmonella* spp. strains to cephalosporins.



Figure 5. Susceptibility of *Salmonella* spp. strains to fluoroquinolones.

It should be noted that conventional microbiological (bacteriological) investigation of acute intestinal infectious diseases, including the sequential steps of bacterial pathogen analysis and interpretation of obtained results, requires from three days to one week, and in some cases even longer. Undoubtedly, results obtained after such prolonged investigations may become of limited relevance for epidemiologists conducting outbreak investigations.

Accordingly, in emergency situations, even preliminary and presumptive information regarding the nature of the causative pathogen is of critical importance for field

epidemiologists involved in outbreak response activities. In recent years, the development of molecular biology and the emergence of modern approaches to epidemic control have necessitated reconsideration of strategies aimed at accelerating bacteriological diagnostics. In particular, significant advances have been achieved in recent years in microbiological methods for determining the etiological agents of acute intestinal infections and foodborne outbreaks. Modern molecular genetic techniques are available for the detection and monitoring of pathogenic and opportunistic microorganisms, as well as for the assessment of their antimicrobial susceptibility. Therefore, it is necessary to revise the current laboratory workflow for microbiological investigation of acute intestinal infections and other infectious diseases in the future and to develop updated diagnostic protocols incorporating modern molecular approaches.

References:

1. Crump J.A., Sjölund-Karlsson M., Gordon M.A., Parry C.M. Epidemiology, clinical presen- tation, laboratory diagnosis, antimicrobial resistance, and antimicrobial management of inva- sive *Salmonella* infections. *Clinical Microbiology Reviews.* 2015; 28(4): 901-37.
2. CDC. National Antimicrobial Resistance Monitoring System (NARMS) Now: Human Data. Atlanta, Georgia: U.S. Department of Health and Human Services, CDC. Available at: https:// www.cdc. gov/narmsnow.
3. EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Preven- tion and Control), 2019. The European Union summary report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2018. *EFSA Journal*. 2019; 17(2): 5598.
4. European Food Safety Authority, European Centre for Disease Prevention and Control, 2021. The European Union One Health 2019 zoonoses report. EFSA J 19:e06406.
5. Park JY, Park K, Ok G, Chang HJ, Park TJ, Choi SW, Lim MC. Detection of *Escherichia coli* O157:H7 Using Automated Immunomagnetic Separation and Enzyme-Based Colo- rimetric Assay. Sensors (Basel). 2020 Mar 4;20(5):1395. doi: 10.3390/s20051395. PMID: 32143335; PMCID: PMC7085514.
6. World Health Organization. Informe Estimación de la carga mundial de las enfermedades de transmisión alimentaria. Ginebra [Internet]. 2015 [Updated: 2017]. Available from: http:// www.who.int/mediacentre/news/releases/2015/foodborne-disease-estimates/es/ [Accessed: February 2017].

ONCOLOGIC OUTCOMES OF HYPERTHERMIC INTRAPERITONEAL CHEMOTHERAPY IN THE TREATMENT OF ASCITES-TYPE OVARIAN CANCER (ATOC): RESIDUAL DISEASE AND METASTASIS

Sulaymanov Dilshod Abdurakhimovich[1]
PhD, MD Head of department of oncology and radiology,
Central Asian Medical University, Fergana, Uzbekistan.
ORCID: 0009-0008-7314-7712

ABSTRACT

Ascites-type ovarian cancer (ATOC) is characterized by advanced disease, poor prognosis, and frequent peritoneal spread. This study evaluated outcomes of cytoreductive surgery with hyperthermic intraperitoneal chemotherapy (HIPEC) versus surgery with standard systemic chemotherapy in 70 patients aged 14–75 years. Patients were divided into a main group (n=34) receiving cytoreductive surgery plus HIPEC and adjuvant polychemotherapy, and a control group (n=36) receiving surgery with systemic chemotherapy alone. HIPEC was performed using a closed perfusion system at 42–43°C. Patients received 6 chemotherapy cycles every 3 weeks. The HIPEC group demonstrated improved outcomes, including a 5-year survival rate of 10.1 ± 1.7%, longer disease-free intervals, and lower recurrence and metastasis rates compared with the control group, in which no 5-year survivors were observed. Distant metastases occurred less frequently in the HIPEC group (8.8% vs 19.4%). These findings suggest that HIPEC combined with cytoreductive surgery may improve survival and reduce disease progression in ATOC patients.

Keywords: Ascites-type ovarian cancer (ATOC); HIPEC; cytoreductive surgery; peritoneal metastasis; ovarian malignancy; peritoneal dialysis; closed-looped system.

Material and Methods

Research involved 70 patients who was ill with ascites type of ovarian cancer and from 14 to 75 years old patients. The oncologic anamnesis continued from 6 months to 9 months. Operation was performed in all patients. Epithelial, germinogen, stromal type of tumors were found in histology. Ascites was ascertained all of the patients [1]. According to treatment, patients are divided into two groups: the 1st group (main) - 34 patients were used cytoreductive operation with HIC then adjuvant PTT was carried out according to the following scheme (CAP, CP, CC); the 2nd group (control) - 36 patients were treated according to the following scheme (CAP, CP, CC), which was used for cytoreductive operation and adjuvant chemotherapy. Chemotherapy cycles were repeated every 3 weeks. There were 6 chemotherapy

cycles. The cycles were conducted with the noted treatment, general condition of patients [2]. HIC was done by following methods: drainage tube installed to patients in a main group for using hyperthermic intraperitoneal chemotherapy in abdominal cavity regardless of size of operation. "Peritoneal dialysis" type drainage system (4 catheters> F-20). The upper drainage tubes were placed on the sidewall of the abdominal cavity and the lower drainage tubes were placed on the small pelvis. HIC method which we created is projected for using in closed contour. It gives an opportunity for treatment before operation and also after operation. It is unmatched Germany, and it is possible to simplify the operation of the operation at the same time. HIC device was created for this methodology [patent for device 19.09.2007 year FAP 00399]. This device involves thermostatic bath, a pump with roller, a magistral line, a thermos control line, exchange of heat and a dish which uses for filtering evacuation fluid. Electron device has relay to manage temperature of additional fluid (picture 1) [3].



Picture 1. Schematics appearance of device for hyperthermic intraperitoneal chemotherapy [Patent No. FAP 00399, 19.09.2007].

Device is used by following way: hyperthermic fluid pour in abdominal cavity by special placed upper drainage. Thermostatic bath warms the infusion fluid to a certain temperature and controls the temperature in the set temperature setting in the heat exchange temperature to maintain the intra-abdominal fluid temperature (42-43°C). When using the control circuitry, the temperature is automatically maintains (45-46° C), which, in turn, ensures constant temperature [4]. In the dynamics, hemodynamic and thermometric parameters of the patient are monitored

continuously. One thermometer is in the rectum, the other in the axillary, and the third one is set to the external acoustic meatus. The main and essential results of our study and the section of results these are the study of the effects of various treatment modalities on survival in patients with ATOC in these research groups [5].



Picture 2. The effect of the type of the treatment of ascites of ovarian cancer to the survival of patients.

When the survival lines are compared (picture 2), the high values in both groups continue to the last 12 months. In the next episode, the difference between the indicators of the groups are evident, and at the end of the observation, there are indicators that have a strong different outcome. As a result, none of patient in the control group had reached the 5-year period. The 5-year survival rate was found to be 10.1 + 1.7% for the patients of main group. In the future, this figure may be higher, as most of the patients of main group have survived and have not reached the 5-year period [6]. In conclusion, it can be said that, as a result of the above-mentioned results, not only hyperthermic intraperitoneal chemotherapy affect the ages of patients, but also it found major changes in the lives of patients. The following our measures are better than literatures.

Relapse and metastases: In 17 of the patients with ascites of the ovarian cancer is found relapse, it is 24.2%. The duration of the non-relapse cycle was determined from the onset of the treatment to the development of the process.



Picture 3. Indicators of disease progression in combination therapy.

As you can see from the picture above, the first relapse of the patients in the main group was detected after surgery in 14 months, and the relapses in 5 patients of the control group were detected 6 months later in 12 cases. The distant metastasis was found in liver and lung tissue, which was observed in 10 patients and was 14.2%. Metastases were found to be 3 (8.8%) patients in the main group and 7 (19.4%) in the control group. Continued growth is less rare in the relapse and metastasis, 7 patients in the control group were found to be 19.4%, and in the main group this figure was 5 patient and 14.7%. Thus, the non-relapse and continuous growth period in the HIC group was longer than the control group [7,8]. In patients with localized OC, primary conclusions can be made about the use of this method:

- The use of this method is relatively safe and effective than systematic therapy;
- There is no strong hematological effect when compared to systemic chemotherapy;
- It extends the non-relapse period in these patients and improves the quality of life.

After a large number of excellent examinations, this method may be recommended for patients with OC. The initial obtained results do not conflict with the knowledge of HIC method of literatures in other pathologies [9].

Discussion of results

The results showed that the spread of ovarian cancer had a positive effect on HIC. This method is valid. Hyperthermia is a key factor in the three-way treatment in a complex treatment:

1. It has its own cytotoxic effect on tumor tissue.
2. It provides deeper penetration of chemotherapeutic agents into tissues and increases the absorption of the drug into the tumor.
3. It increases the cytotoxicity of chemotherapeutic agents [10].

Using a closed-looped system by us has its advantages: There are all conditions to control and maintain the adequate temperature of the perfume, the efficient use of fluids and cytostatic and their long-term circulation [11]. The chemotherapeutic agents applied to this method is for a long time in the abdominal cavity, which is associated with a plasma barrier, in result increases the concentration of the chemotherapy included in the abdominal cavity and it is maintained throughout the therapy period. The thermal factor is also important, which allows the drug to penetrate into the tissue and also has its own cytotoxic effect and enhances the cytotoxic effects of chemicals [12].

Conclusions

Thus, based on the results of this study, it can be concluded that the hyperthermic intraperitoneal chemotherapy (HIC/HIPEC) method contributes to improving the overall treatment outcomes in patients with ascites associated with both localized and disseminated ovarian cancer. The obtained data indicate that the inclusion of HIPEC in the combined treatment approach enhances oncologic effectiveness compared with standard treatment consisting of cytoreductive surgery followed by systemic chemotherapy alone. Importantly, the analysis of perioperative and postoperative outcomes demonstrated that the application of HIPEC does not significantly increase the rate of complications or negatively affect patient safety when compared with conventional treatment methods. The procedure was generally well tolerated, and no substantial increase in treatment-related morbidity was observed, which supports its feasibility in clinical practice. In addition, HIPEC was associated with improved disease control, including longer disease-free intervals and reduced rates of recurrence and metastatic progression. These findings suggest a potential benefit of locoregional hyperthermic chemotherapy in eliminating residual microscopic tumor cells within the peritoneal cavity, which are often responsible for disease relapse. Based on these results, it is advisable to consider HIPEC as a valuable component of multimodal treatment in ovarian cancer with ascites. Furthermore, continuation

of systemic chemotherapy after HIPEC is recommended as an essential step for prophylaxis and further reduction of the risk of disease progression at the subsequent stages of treatment

References:

1. Abdikhakimov A.N., Kasymov A.A., Abdukhakimov U.N., Sulamanov D.A. Immediate results of postoperative hyperthermic intraperitoneal chemoperfusion. In: Proceedings of the 3rd Congress of Oncologists of Uzbekistan (May 14–16, 2015, Tashkent). Tashkent; 2015. p. 21.
2. Akhmedov O.M., Kurbanov D.D., Tillashaykhova M.Kh., Ishchenko I.V. Efficacy of neoadjuvant chemotherapy in ovarian cancer. News of Dermatovenereology and Reproductive Health. 2015;(2):155–156.
3. Baklanova N.S., Kolomiets L.A. Ultrasound semiotics in recurrent ovarian cancer after optimal cytoreductive surgery. Problems in Oncology. 2014;60(3):323–326.
4. Belyaeva O.A., Bespalov V.G. Experimental technology of chemoperfusion treatment for peritoneal carcinomatosis in ovarian cancer. Problems in Oncology. 2014;60(1):71–79.
5. Vyshenskaya E.A., Bespalov V.G. New regimens of intraperitoneal polychemotherapy for treatment of peritoneal carcinomatosis in ovarian cancer. Journal of Obstetrics and Women's Diseases. 2014;63(5):38–45.
6. Guseynov K.D., Belyaev A.M. Hyperthermic intraperitoneal chemotherapy (HIPEC) in the treatment of recurrent ovarian cancer. Problems in Oncology. 2014;60(3):343–347.
7. Sulaymanov D.A., Khasanov A.I., Abdikhakimov A.N., Abdukhakimov A.N. Immediate results of combined treatment in ascitic forms of ovarian cancer. Clinical and Experimental Oncology. Uzbekistan (Fergana). 2018;3(5):122.
8. Sulaymanov D.A., Abdikhakimov A.N., Khasanov A.I., Abdukhakimov A.N. Closed technique of hyperthermic intraperitoneal chemotherapy in intraperitoneally disseminated cancer. Clinical and Experimental Oncology. Uzbekistan (Fergana). 2018;3(5):144.
9. Arimoto T., Nakagawa S., Oda K., Kawana K., Yasugi T., Taketani Y. Second-line chemotherapy with docetaxel and carboplatin in paclitaxel- and platinum-pretreated ovarian, fallopian tube, and peritoneal cancer. Medical Oncology. 2015.

10. Carrabin N., Mithieux F., Meeus P., Trédan O., Guastalla J.P., Bachelot T., et al. Hyperthermic intraperitoneal chemotherapy with oxaliplatin without adjuvant chemotherapy in stage IIIC ovarian cancer. Bulletin du Cancer. 2015;98(6): E32.

11. Chi D.S., Cui Y., Cui M.H., Jia Y. Choice of chemotherapy methods for epithelial ovarian carcinoma: a meta-analysis. Zhonghua Yi Xue Za Zhi. 2016;97(20):177.

12. Muenyi C.S., States V.A., Masters J.H., Fan T.W., Helm C.W., States J.C. Sodium arsenite and hyperthermia modulate cisplatin–DNA damage responses and enhance platinum accumulation in murine metastatic ovarian cancer xenografts after HIPEC. Journal of Ovarian Research. 2016;9:28.

HEALTHCARE REFORMS AS A STRATEGIC FACTOR IN POVERTY REDUCTION

Daripov Shahobiddin Israiljonovich
Central Asian Medical University, Senior Lecturer, Department of Social Sciences and Physical Culture. Uzbekistan, Fergana
ORCID.ID. https://orcid.org/0009-0003-2813-7623

ABSTRACT
This article analyzes the role and significance of ongoing healthcare reforms in improving population welfare and reducing poverty levels. The author justifies that the efficiency of the healthcare system directly affects not only medical indicators but also the socio-economic stability of the country. The article focuses on the strategic changes implemented in the sector under the leadership of the President of the Republic of Uzbekistan, Shavkat Mirziyoyev, including strengthening primary healthcare, expanding emergency medical services, developing the pharmaceutical industry, and supporting the private sector. It also highlights the negative impact of unexpected medical expenses on family budgets and the role of state-provided preferential medical services as a shield for social protection. The main conclusion of the article is that investment in healthcare is the most effective means of breaking the intergenerational chain of poverty through the development of human capital.
Keywords: healthcare reforms, poverty reduction, social protection, human capital, primary healthcare, health economics, prevention, pharmaceuticals, private medicine, digital medicine.

INTRODUCTION

At the center of social state policy aimed at honoring human dignity and ensuring their interests lies the issue of building a healthy society. It is a globally recognized fact that a country's economic power is determined not only by its natural resources or industrial potential, but primarily by healthy and productive human capital. From this perspective, healthcare reforms represent not only the modernization of the healthcare system but also the most crucial strategic factor in reducing poverty and achieving social welfare.

The intrinsic connection between economic hardship and health can be articulated through the concept of the "vicious cycle of poverty". Loss of working capacity due to illness leads to a decrease in income, while unexpected expenditures on medical services cause family budget crises. Consequently, the vulnerable segment of the population in need of social protection falls into a deeper vortex of economic problems. This makes expanding access to quality and affordable medical services through healthcare reform an urgent task.

In recent years, a process of fundamental transformation in the medical field has begun as a priority direction of state policy in Uzbekistan. As President Shavkat Mirziyoyev noted, "If the medical system is properly established and people's health is strengthened, a solid foundation will be created for reducing poverty in society". Today, organizing medical services on a neighborhood-based (mahallabay) system, implementing digital medicine, and alleviating the financial burden on the population by developing medical insurance mechanisms have become the main criteria of these reforms.

This article analyzes the impact of qualitative changes in the medical system on poverty reduction, the economic efficiency of prevention and patronage services, and the prospects for achieving sustainable economic growth through human capital development. Global experience shows that economic distress and health problems form an interlocking chain. Only when a person is healthy can they work, earn an income, and contribute to the development of society. Therefore, today, reforming the medical system has become one of the most effective tools for

combating poverty. In many cases, unexpected severe illnesses and the large expenses required for their treatment cause families to fall into poverty. The primary goal of the reforms is to reduce the burden on people's pockets by making medical services affordable and high-quality.

CONCEPTUAL FRAMEWORK

To this end, on October 18, 2017, President of the Republic of Uzbekistan Shavkat Mirziyoyev held a meeting dedicated to analyzing reforms aimed at further improving the healthcare system, enhancing the quality and efficiency of medical services, strengthening public health, and ensuring the provision of quality medicines. State-issued warrants and free drug supply protect low-income families from economic crises.

Prevention is economic thrift. Preventing a disease is several times cheaper for both the state and the citizen than treating it. Tremendous work is being done to provide the population with quality and affordable medicines, protect motherhood and childhood, improve emergency medical care, further enhance the primary healthcare system, and develop the private sector and specialized medical care. Detecting diseases at an early stage ensures that the patient does not lose their ability to work. This improves the quality of labor resources and reduces the need for social benefits.

Within the framework of implementing the resolution on improving primary healthcare, 793 rural medical centers were reorganized into rural family polyclinics, and 441 emergency medical care stations were established. The buildings of 658 discontinued rural medical centers were reallocated to doctors to be used as official housing. In addition, the activities of 5 types of specialized professionals were launched in rural family polyclinics and 10 types in Tashkent city family polyclinics.

To improve staffing in the primary system, 79 percent of the 3,197 young specialists who graduated from the bachelor's degree programs of medical higher educational institutions this year were directed to work in the primary healthcare sector. Investment in medicine is an investment in human capital. A healthy generation receives a better education and is capable of working in high-tech jobs.

The insufficient quality of medical services provided to the population in the regions is one of the pressing problems facing us. Admittedly, people having to travel from the regions to Tashkent to receive high-tech medical care causes numerous objections from the population. One of the pressing issues causing rightful public concern relates to emergency medical services. We still have a lot of work to do in this regard; transportation vehicles are still insufficient, and there are interruptions in the supply of medicines. While 806 emergency medical care stations and substations operated in 2016, this indicator exceeded 2,100 in 2017. Naturally, this is of great importance in ensuring the timely arrival of every emergency call. Along with the services provided in this direction, it is necessary to shape the culture of the population regarding the proper utilization of this service.

METHODOLOGY

One of the urgent issues facing global healthcare is social diseases. Branches of republican specialized centers are being established in the provinces, and over 50 of our compatriots working in foreign clinics have returned to our country.

The reduction of undergraduate medical education from 7 to 6 years, and from 6 to 5 years for the "medical prevention" direction, serves to adapt our country's medical education system to world standards, improve localized staffing, and ensure the targeted spending of state budget funds. Under the decree of the President regarding the improvement of the state support system for disabled persons, 8,400 disabled individuals are provided with prosthetic and

orthopedic products, and over 9,300 needy citizens are provided with technical means for rehabilitation free of charge. Additionally, more than 23,000 veterans of the 1941-1945 war, lonely elderly individuals, pensioners, and disabled people receive free recuperation at sanatoriums.

Following the resolution on improving the development of the private healthcare sector, the types of medical specialties that private medical institutions can engage in were increased from 50 to 126. As a result, joint clinics are being established in Tashkent with countries like South Korea, India, and Turkey; in Jizzakh with South Korea; and in Fergana, Karshi, and the Kumkurgan district with India. Over the past 9 months, the number of private medical institutions has increased by more than 400, reaching over 3,860, which reflects the effectiveness of broad privileges and preferences granted to private medical institutions.

In 2017, the production of 76 new products was mastered by the "Uzfarmsanoat" joint-stock concern. The production of medicines and medical devices increased fourfold compared to 2011. To assimilate the production of new, high-tech products in the pharmaceutical industry, credit resources amounting to 250 million dollars were allocated from the Fund for Reconstruction and Development of Uzbekistan. The implementation of more than 70 investment projects with a total value of almost 400 million dollars has begun, 33 of which will be launched this year.

The consistent work carried out in this direction has begun to yield results. For instance, a 40% increase in pharmaceutical exports has been achieved, and production efficiency per worker has grown by 35% due to the introduction of high technologies. The types of medicines produced by local pharmaceutical enterprises have exceeded 2,300. Special attention was paid to further improving the emergency medical service, strengthening the material and technical base of screening centers, and enhancing the quality of manufactured prosthetic and orthopedic products and technical rehabilitation aids.

The healthy growth of children lays the foundation for them to become active and productive members of society in the future. This, in turn, helps to break the chain of intergenerational poverty. To fully cover the population with patronage and systematic clinical examination, further improve maternal and child screening and medical-genetic services, and bring these services closer to the population, the State Program for 2018-2022 was developed.

The reduction of corruption and bureaucracy in healthcare ensures the success of social reforms; the distribution of medicines and provision of medical services through electronic systems guarantee that the most vulnerable segments receive genuine assistance.

CONCLUSION AND DISCUSSION

An analysis of the socio-economic aspects of healthcare reforms shows that the healthcare system is not only a place for treating diseases but also a strategic mechanism for reducing poverty and ensuring sustainable development in the country. Based on the research and analyses conducted, the following final conclusions can be drawn:

- Medical and Economic Protection: The digitalization of medical services and the expansion of state-guaranteed free medical care in our country have significantly reduced unexpected (catastrophic) medical expenses for the population, especially for low-income segments. This serves as the main factor preventing families from falling into poverty due to medical reasons.
- Quality of Human Capital: Investments directed toward healthcare increase the productivity of human capital. A healthy workforce becomes competitive in the labor market, which in turn leads to consistent growth in population income and increased economic activity in society.
- Efficiency of Prevention: Strengthening primary healthcare and the local introduction of the patronage system have enabled the detection of diseases at an early stage. This not only extends

human life expectancy but also saves the state budget and citizens' funds from high treatment costs.

- Pharmaceuticals and Innovation: Developing the local pharmaceutical industry and paving the way for private medicine have ensured the affordability of medical products for the population. This has been an important step toward establishing the principles of social equality.

As a concluding thought, it is worth emphasizing that the success of the medical system directly defines the social image of a society. A healthy nation means a society free from poverty, economically independent, and looking to the future with confidence. Therefore, consistently continuing reforms in healthcare remains the most reliable foundation for achieving the strategic goals of New Uzbekistan.

REFERENCES

I. Normative-legal documents and official sources

1. Constitution of the Republic of Uzbekistan. (New edition). — Tashkent: "Uzbekistan", 2023. (Especially articles concerning the right to medical care).
2. Decree of the President of the Republic of Uzbekistan. On the "Uzbekistan — 2030" Strategy. No. UP-158. September 11, 2023.
3. Decree of the President of the Republic of Uzbekistan. "On comprehensive measures to radically improve the healthcare system of the Republic of Uzbekistan." No. UP-5590. December 7, 2018.
4. Resolution of the President of the Republic of Uzbekistan. "On introducing new mechanisms for organizing the activities of primary healthcare institutions." No. PP-4890. November 12, 2020.
5. Decree of the President of the Republic of Uzbekistan. "On the development strategy of New Uzbekistan for 2022–2026." No. UP-60. January 28, 2022.

II. Works and speeches of the President

6. Mirziyoyev Sh.M. Strategy of New Uzbekistan. — Tashkent: "Uzbekistan" publishing house, 2021. — 464 p.
7. Mirziyoyev Sh.M. The consent of our people is the highest assessment given to our activities. Volume 2. — Tashkent: "Uzbekistan", 2018.
8. Mirziyoyev Sh.M. We will resolutely continue our path of national development and raise it to a new level. Volume 1. — Tashkent: "Uzbekistan", 2017.

III. Scientific literature, textbooks, and articles

9. Shodmonov A.K. Health Economics. Textbook. — Tashkent: "Iqtisod-Moliya", 2019.
10. Alimov A.V. et al. Stages of reforming the system of protection of motherhood and childhood in Uzbekistan. Scientific-practical manual. — Tashkent, 2020.
11. Ibragimov A.Yu. Management in medicine and the effectiveness of social reforms. — Tashkent: "Abu Ali ibn Sino", 2021.
12. Mahmudova G.A. Social policy and public health: an analysis of interconnection // Economics and Finance (Uzbekistan). — 2022. — No. 4.
13. Rasulov B.M. Digitalization of the healthcare system as a factor in public welfare // Journal of Society and Management. — 2023. — No. 2.
14. State Committee of the Republic of Uzbekistan on Statistics. Statistical collection "Social development and living standards of Uzbekistan". — Tashkent, 2024.
15. Report of the international expert group. The impact of poverty reduction and health reforms in Uzbekistan: UNICEF and World Bank analysis. — Tashkent, 2022.

AGE-SPECIFIC PREVENTIVE STRATEGIES FOR ATHEROSCLEROSIS AND THEIR EFFECTIVENESS

Shakhnoza Kholboboyeva Asadullayeva
Central Asian Medical University, Fergana, Uzbekistan
ORCID ID: https://orsid.org/0009-0008-2123-6991

Abstract
Atherosclerosis is a major underlying cause of cardiovascular diseases, with aging being a critical determinant in its development. This study provides an in-depth analysis of age-related vascular changes, including endothelial dysfunction, lipid metabolism disorders, oxidative stress, and chronic inflammation. The effectiveness of age-specific preventive strategies was evaluated across different age groups. The findings indicate that tailored preventive approaches significantly reduce the progression of atherosclerosis and associated cardiovascular risks.
Keywords: atherosclerosis, aging, prevention, endothelial dysfunction, lipid metabolism, inflammation

RELEVANCE OF THE STUDY

Atherosclerosis is one of the most pressing problems in modern medicine, serving as the primary pathogenetic substrate of cardiovascular diseases and imposing a significant socio-economic burden on the global healthcare system. According to the World Health Organization, cardiovascular diseases account for approximately 30–32% of all deaths worldwide, with atherosclerotic processes identified as the principal etiopathogenetic factor [1]. The significance of this condition is determined not only by its high mortality rates but also by early disability, reduced work capacity, and increased healthcare expenditures.

The development of atherosclerosis is a complex and multistage process involving lipid infiltration, accumulation of oxidized lipoproteins, activation of the monocyte-macrophage system, formation of foam cells, and the development of fibrous plaques. During this process, structural changes occur in all layers of the vascular wall—intima, media, and adventitia. In particular, the oxidation of low-density lipoproteins (LDL) and their subsequent phagocytosis by macrophages play a decisive role in the early stages of atherogenesis [2]. In addition, oxidative stress and the action of free radicals intensify lipid peroxidation, leading to damage to cell membranes and impairment of endothelial function.

Age is considered a central determinant in the pathogenesis of atherosclerosis. During the aging process, the elastic components of the vascular wall—particularly elastin fibers—undergo degeneration and are replaced by more rigid collagen fibers, resulting in reduced vascular elasticity and increased stiffness. The increase in arterial stiffness leads to elevated hemodynamic load, higher systolic blood pressure, and increased cardiac workload. Furthermore, age-related decline in the proliferative and regenerative capacity of endothelial cells results in impairment of the endothelial barrier function.

Endothelial dysfunction is one of the earliest and most critical stages of atherosclerosis, characterized by reduced bioavailability of nitric oxide (NO), impaired vasodilation mechanisms, and increased thrombogenicity. A decrease in NO synthesis leads to increased vascular tone, enhanced platelet aggregation, and activation of inflammatory mediators, thereby promoting plaque formation and progression [2].

According to modern scientific concepts, atherosclerosis is not merely a consequence of lipid metabolism disorders but is also regarded as a chronic low-grade inflammatory disease.

The aging-associated inflammatory process, known as "inflammaging," is characterized by increased levels of cytokines such as IL-6, TNF-α, and C-reactive protein (CRP), which accelerate the formation of atherosclerotic plaques [3]. Age-related alterations in the immune system (immunosenescence) further contribute to uncontrolled inflammatory responses, leading to the progression of pathological changes in the vascular wall[11,12,13].

In addition, epigenetic changes, mitochondrial dysfunction, and cellular senescence play a significant role in the development of atherosclerosis. Senescent cells release pro-inflammatory mediators, known as the senescence-associated secretory phenotype (SASP), which promote inflammation and fibrosis within the vascular wall. This leads to decreased plaque stability and an increased risk of plaque rupture.

Recent epidemiological and morphological studies have demonstrated that atherosclerotic processes begin as early as childhood and adolescence. For example, studies conducted by McGill and colleagues revealed the presence of early lipid streaks in the aorta and coronary arteries of young individuals [4]. This finding highlights the necessity of viewing atherosclerosis not merely as a disease of old age but as a lifelong progressive pathological process[5,6,7].

From this perspective, the development and implementation of age-specific preventive strategies for atherosclerosis are of paramount importance. Preventive approaches should be differentiated according to age: in young individuals, the primary focus should be on establishing healthy lifestyle habits; in middle-aged individuals, metabolic control and reduction of risk factors become crucial[8,9,10]; and in older adults, pharmacological prevention and the management of complications are of primary importance.

Thus, the in-depth study of age-related features of atherosclerosis and the development of appropriate preventive strategies represent one of the key directions of modern medicine. Such research significantly expands the possibilities for effective disease management in clinical practice.

Aim of the study. The aim of this study was to comprehensively investigate the mechanisms of atherosclerosis development across different age groups and to scientifically evaluate the effectiveness of age-specific preventive strategies.

MATERIALS AND METHODS

This study was conducted in the city of Fergana between 2023 and 2025. A total of 150 participants were included and divided into three age groups: young adults aged 18–30 years, middle-aged individuals aged 31–50 years, and elderly individuals aged 51–75 years. Each group consisted of 50 participants, selected in a balanced manner with respect to gender and socio-demographic characteristics.

The study methodology was based on a comprehensive approach. Biochemical analyses included the determination of total cholesterol, low-density lipoproteins (LDL), high-density lipoproteins (HDL), and triglyceride levels. Instrumental assessments involved Doppler ultrasound examination to measure carotid artery intima-media thickness. In addition, arterial stiffness was evaluated by measuring pulse wave velocity (PWV). Endothelial function was assessed using the flow-mediated dilation (FMD) method.

Preventive strategies included a balanced diet based on the Mediterranean dietary model, regular physical activity (at least 150 minutes per week), limitation of harmful habits, and pharmacological prevention (statins and, when indicated, antiplatelet agents). Statistical analysis was performed using SPSS version 26, and results were considered statistically significant at $p<0.05$.

RESULTS AND DISCUSSION

The results of the study once again confirmed that age is a significant determinant in the development of atherosclerosis. According to the obtained data, progressive pathological changes occur in lipid metabolism, vascular wall elasticity, and endothelial function with increasing age. In particular, a significant increase in low-density lipoprotein (LDL) levels was observed in the 51–75 age group, indicating the formation of an atherogenic lipid profile. This condition can be explained by age-related decline in hepatic lipid metabolism, decreased activity of lipoprotein lipase, and functional insufficiency of LDL receptors. As a result, the accumulation of atherogenic fractions in the blood increases, and lipid infiltration into the vascular intima accelerates.

In addition, arterial stiffness indicators, particularly pulse wave velocity (PWV), were found to increase progressively with age. This phenomenon is associated with degeneration of elastin fibers, increased collagen content, and enhanced vascular calcification. Increased arterial stiffness leads to elevated hemodynamic load, increased systolic blood pressure, and left ventricular hypertrophy. From this perspective, arterial stiffness can be considered not only a marker of atherosclerosis but also a factor that accelerates its progression.

Endothelial dysfunction indicators also showed significant differences among age groups. In elderly individuals, a decrease in flow-mediated dilation (FMD) was observed, indicating reduced bioavailability of nitric oxide (NO). Reduced NO synthesis impairs the vasoprotective function of the endothelium, resulting in vasoconstriction, enhanced platelet aggregation, and increased activity of inflammatory mediators. These processes contribute to the formation and destabilization of atherosclerotic plaques. Furthermore, increased oxidative stress accelerates the inactivation of NO molecules and leads to the formation of toxic metabolites such as peroxynitrite.

The effectiveness of the preventive strategies applied during the study was demonstrated by positive outcomes across all age groups; however, their intensity and impact varied depending on age. The Mediterranean diet, used as a rational nutritional model, played a crucial role in normalizing the lipid profile. In particular, a diet rich in unsaturated fatty acids reduced LDL levels and increased HDL levels, leading to a decrease in the atherogenic index. On average, LDL levels decreased by 18–25%, which is considered clinically significant.

Increased physical activity also had a positive effect on metabolic processes. Regular aerobic exercise improved insulin sensitivity, enhanced lipid metabolism, and stimulated nitric oxide synthesis in the vascular wall. In addition, physical activity exerted anti-inflammatory effects, reducing levels of C-reactive protein (CRP) and other pro-inflammatory markers. This contributes to the attenuation of the “inflammaging” process.

Pharmacological prevention, particularly statin therapy, proved to be one of the most effective approaches for controlling lipid levels. Statins not only reduce LDL levels but also exert pleiotropic effects, including improvement of endothelial function, reduction of oxidative stress, and stabilization of atherosclerotic plaques. During the study, participants receiving statin therapy demonstrated a significantly lower risk of cardiovascular complications.

The results showed that the implementation of a comprehensive preventive approach reduced the risk of cardiovascular complications by 25–30%. These findings are consistent with international clinical studies. For example, the PREDIMED study conducted by Estruch et al. demonstrated that the Mediterranean diet significantly reduces the risk of cardiovascular diseases [5]. In addition, numerous randomized controlled trials on statins have confirmed their role in stabilizing atherosclerotic plaques and reducing thrombotic complications [6].

During the discussion, particular attention was given to age-related inflammatory processes. The phenomenon of “inflammaging” is characterized by increased levels of cytokines, particularly IL-6, TNF-α, and C-reactive protein. These mediators enhance

inflammatory infiltration in the vascular wall, accelerating plaque formation and progression. Furthermore, age-related remodeling of the immune system (immunosenescence) leads to a weakening of protective mechanisms and contributes to the progression of pathological processes.

Moreover, mitochondrial dysfunction and cellular senescence play an important role in the pathogenesis of atherosclerosis. Senescent cells produce components of the senescence-associated secretory phenotype (SASP), which sustain chronic inflammation and promote fibrotic processes within the vascular wall. This reduces plaque stability and increases the risk of plaque rupture, potentially leading to acute coronary syndromes.

Thus, the obtained results and their discussion indicate that the application of age-specific preventive strategies is essential for the prevention of atherosclerosis. This approach demonstrates high effectiveness in normalizing lipid metabolism, restoring endothelial function, reducing inflammatory processes, and lowering overall cardiovascular risk.

CONCLUSION

Age is a fundamental determinant in the development of atherosclerosis and is closely associated with the general biological mechanisms of aging. With advancing age, complex morphological and functional changes occur in the vascular wall, including degradation of elastic fibers, increased collagen deposition, proliferation and migration of smooth muscle cells, and activation of calcification processes. These alterations lead to reduced vascular elasticity and increased stiffness, resulting in elevated hemodynamic load and the development of arterial hypertension. These conditions, in turn, create a significant pathophysiological basis for the formation and progression of atherosclerotic plaques.

In addition, age-related decline in endothelial cell function leads to reduced nitric oxide (NO) synthesis and weakening of anti-atherogenic defense mechanisms. As a result of endothelial dysfunction, vasodilation is impaired, platelet aggregation is enhanced, inflammatory mediator release increases, and favorable conditions for lipid infiltration into the vascular wall are created. At the same time, increased oxidative stress intensifies lipid peroxidation and promotes the formation of oxidized LDL, which is phagocytosed by macrophages, leading to foam cell formation and activation of early atherogenesis.

Another important factor associated with aging is chronic low-grade inflammation, known as “inflammaging.” This process is characterized by increased levels of pro-inflammatory cytokines (IL-6, TNF-α, CRP), which enhance inflammatory infiltration in the vascular wall and accelerate plaque formation and destabilization. Furthermore, age-related remodeling of the immune system (immunosenescence) reduces the effectiveness of protective mechanisms and contributes to the uncontrolled progression of pathological processes.

The findings of this study demonstrate that the use of age-specific preventive strategies in atherosclerosis prevention is highly clinically effective. Preventive measures developed based on a differentiated approach across age groups contribute to normalization of lipid metabolism, restoration of endothelial function, improvement of arterial elasticity, and reduction of inflammatory processes. In particular, the Mediterranean diet exerts anti-atherogenic effects by improving the lipid profile, reducing oxidative stress, and enhancing antioxidant defense mechanisms. Physical activity, in turn, promotes metabolic homeostasis, increases insulin sensitivity, and stimulates nitric oxide synthesis in the vascular wall, thereby slowing the progression of atherosclerosis.

Pharmacological prevention, especially the use of statins and antiplatelet agents, plays an important role in preventing disease progression in high-risk patients. The pleiotropic effects of statins—including lipid-lowering, improvement of endothelial function, reduction of inflammation, and stabilization of plaques—make them a key component of atherosclerosis

prevention. At the same time, modern approaches emphasize the need to individualize preventive strategies based on risk assessment.

The results and their analysis indicate that effective prevention of atherosclerosis requires early, continuous, and stepwise implementation of preventive measures throughout life. The main principle of prevention is the lifelong minimization of risk factors and the promotion of a healthy lifestyle. In this regard, the widespread implementation of age-specific preventive strategies not only at the individual level but also at the population level can significantly reduce the burden of cardiovascular diseases.

Future research should focus on in-depth investigation of the molecular and genetic mechanisms of atherosclerosis, the development of early diagnostic systems based on biomarkers, and the improvement of personalized preventive and therapeutic approaches. This will expand the possibilities for early detection and effective management of the disease, ultimately reducing its complications.

REFERENCES

1. World Health Organization. Cardiovascular diseases (CVDs). Geneva; 2023.
2. Libby P. The changing landscape of atherosclerosis. *Nature*. 2021;592:524–533.
3. Franceschi C., Campisi J. Chronic inflammation (inflammaging). *J Gerontol A Biol Sci Med Sci*. 2014;69:S4–S9.
4. McGill H.C. et al. Atherosclerosis in youth. *Circulation*. 2000;102:374–379.
5. Estruch R. et al. Mediterranean diet and cardiovascular prevention. *N Engl J Med*. 2018;378:25–34.
6. Stone N.J. et al. 2019 ACC/AHA cholesterol guideline. *Circulation*. 2019;139:e1082–e1143.
7. Ross R. Atherosclerosis — an inflammatory disease. *N Engl J Med*. 1999;340:115–126.
8. Ridker P.M. et al. Anti-inflammatory therapy with canakinumab for atherosclerotic disease. *N Engl J Med*. 2017;377:1119–1131.
9. Benjamin E.J. et al. Heart disease and stroke statistics—2020 update. *Circulation*. 2020;141:e139–e596.
10. North B.J., Sinclair D.A. The intersection between aging and cardiovascular disease. *Circ Res*. 2012;110:1097–1108.
11. Lakatta E.G., Levy D. Arterial and cardiac aging: major shareholders in cardiovascular disease. *Circulation*. 2003;107:139–146.
12. Hansson G.K. Inflammation, atherosclerosis, and coronary artery disease. *N Engl J Med*. 2005;352:1685–1695.
13. O'Donnell C.J., Elosua R. Cardiovascular risk factors. *Circ Res*. 2008;102:20–28.

ТУПРОҚДАГИ ЎТА ХАВФЛИ ЮҚУМЛИ КАСАЛЛИКЛАР ТАРҚАТУВЧИ МИКРООРГАНИЗМЛАР ВА УЛАРНИНГ ЭКОЛОГИЯСИ

MICROORGANISMS CAUSING HIGHLY DANGEROUS INFECTIOUS DISEASES IN SOIL AND THEIR ECOLOGY

Shodmonov Usmonbek Bakhodir ugli
Central Asian Medical University, Fergana, Uzbekistan
https://orcid.org/0000-0003-2247-4672

Аннотация
Тезисда тупроқ муҳитида узоқ муддат сақланиб қолиши мумкин бўлган ва одам ҳамда ҳайвонлар учун ўта хавфли юқумли касалликларни келтириб чиқарувчи асосий микроорганизмлар, хусусан Bacillus anthracis, Clostridium tetani, Clostridium botulinum ва Clostridium perfringensнинг биологик хусусиятлари ҳамда экологик омиллар билан боғлиқлиги ёритилган. Шунингдек, уларнинг тупроқда сақланишига таъсир этувчи намлик, ҳарорат, pH, органик моддалар миқдори, аэрация ва инсон хўжалик фаолияти каби омиллар таҳлил қилинган. Тупроқ орқали тарқалувчи инфекцияларнинг эпидемиологик аҳамияти, профилактика чоралари ва санитария-экологик назоратнинг муҳим жиҳатлари баён этилган.
Калит сўзлар: тупроқ, патоген микроорганизмлар, куйдирги, қоқшол, ботулизм, газли гангрена, экология, профилактика, санитария, эпидемиология
Abstract
This thesis examines the main microorganisms that can persist in the soil environment for long periods and cause highly dangerous infectious diseases in humans and animals, particularly *Bacillus anthracis, Clostridium tetani, Clostridium botulinum,* and *Clostridium perfringens*, as well as their biological characteristics and relationship with ecological factors. It also analyzes the factors affecting their persistence in soil, including moisture, temperature, pH, the amount of organic matter, aeration, and human economic activity. The epidemiological significance of soil-borne infections, preventive measures, and the important aspects of sanitary-ecological control are also described.
Keywords: soil, pathogenic microorganisms, anthrax, tetanus, botulism, gas gangrene, ecology, prevention, sanitation, epidemiology.

INTRODUCTION

Тупроқ табиий экотизимнинг энг мураккаб ва биологик жиҳатдан фаол муҳитларидан бири бўлиб, унда миллиардлаб микроорганизмлар яшайди. Уларнинг аксарияти моддалар айланиши, органик қолдиқларни парчалаш ва тупроқ унумдорлигини таъминлашда ижобий аҳамиятга эга. Бироқ айрим патоген микроорганизмлар тупроқда узоқ муддат сақланиб, одам ва ҳайвонлар учун ўта хавфли юқумли касалликлар манбаи бўлиб хизмат қилади. Айниқса спора ҳосил қилувчи бактериялар экологик жиҳатдан чидамли бўлгани сабабли улар тупроқда йиллар, ҳатто айрим ҳолатларда ўнлаб йиллар давомида сақланиши мумкин. Шу боис тупроқ фақат агрономик ресурс эмас, балки санитария-эпидемиология нуқтаи назаридан ҳам муҳим объект ҳисобланади.

Долзарблиги

Иқлим ўзгариши, сув тошқинлари, қурғоқчилик, чорвачилик айланмаси, тупроқ қазиш ишлари, санитария қоидаларининг етарли бажарилмаслиги ва чиқиндилар билан ифлосланиш каби омиллар тупроқдаги патоген микроорганизмларнинг фаоллашуви ҳамда инсонга ўтиш хавфини кучайтириши мумкин. Тиббиёт, ветеринария, экология ва жамоат саломатлиги соҳалари кесишган нуқтада мазкур муаммони ўрганиш профилактика чораларини илмий асосда ташкил этиш учун зарурдир.
Мақсад. Тупроқда учрайдиган ва ўта хавфли юқумли касалликларни қўзғатиши мумкин бўлган асосий микроорганизмлар, уларнинг биологик хусусиятлари, тарқалиш йўллари, сақланиб қолишини белгиловчи экологик омиллар ҳамда профилактика тамойилларини ёритиш.

Асосий қисм

1. Тупроқ муҳити ва патоген микроорганизмлар учун экологик ниша. Тупроқ намлик, ҳарорат, pH, органик моддалар миқдори, кальций ва бошқа минерал тузлар, аэрация даражаси, микро- ва макрофауна фаоллиги каби кўрсаткичлар билан тавсифланади. Ушбу омиллар патогенларнинг сақланиши, споралaниши, вегетатив шаклга ўтиши ёки аксинча инактивациясини белгилайди. Тупроқдаги микроёриқлар, органик қолдиқлар ва анаэроб микрозоналар айрим бактериялар учун қулай яшаш жойини ҳосил қилади. Шу билан бирга, тупроқдаги микроблар орасида рақобат, антагонизм ва фаглар таъсири ҳам патогенлар тақдирига таъсир кўрсатади.

2. Куйдирги қўзғатувчиси – *Bacillus anthracis*. Куйдирги тупроқ билан боғлиқ энг хавфли зооноз касалликлардан бири ҳисобланади. Bacillus anthracis граммусбат, аэроб, спора ҳосил қилувчи таёқчасимон бактерия бўлиб, унинг споралари ташқи муҳитга жуда чидамли. Инфекция манбаи асосан касал ёки нобуд бўлган чорва ва ёввойи ҳайвонлар ҳисобланади. Улардан чиққан биологик материал тупроқни ифлослантиради, натижада ўчоқ ҳудудлар узоқ муддат давомида хавфли бўлиб қолади. Тупроқнинг ишқорийга яқин ёки ишқорий pH кўрсаткичи, кальций ва органик моддаларнинг нисбатан юқори бўлиши, намлик ва сув тошқинлари споралaрнинг сақланиб қолиши ва юзага чиқишида муҳим омил сифатида қайд этилади. Чорва молларининг инфекцияланган ерларда ўтлаши куйдирги циклини қайта тиклайди. Одамларга эса касал ҳайвон билан мулоқот, ифлосланган тери ва жун маҳсулотлари, кам ҳолларда аэрозол ва озиқ-овқат орқали юқиши мумкин.

3. Қоқшол қўзғатувчиси – *Clostridium tetani*. Қоқшол тупроқ, чанг ва ҳайвон ахлатида кенг тарқалган спора ҳосил қилувчи анаэроб бактерия билан боғлиқ касалликдир. У одатда инсондан инсонга юқмайди; касаллик чуқур жароҳат, тешилган яра, куйиш, совқотиш ёки некрозланган тўқимага споралар тушганда ривожланади. Анаэроб шароитда споралар вегетатив шаклга ўтиб, кучли нейротоксин – тетаноспазмин ишлаб чиқаради. Тупроқнинг ўзида бактерия кўпинча спора шаклида сақланади; унинг патогенлик механизми эса хўжайин организми ичида намоён бўлади. Қишлоқ хўжалиги ишлари, ифлос асбоблардан фойдаланиш, санитария талабларига риоя этмаслик ва вакциналаниш даражасининг пастлиги қоқшол хавфини оширади.

4. Ботулизм қўзғатувчиси – *Clostridium botulinum*. Ушбу бактерия споралари табиатда, жумладан тупроқда кенг тарқалган. Clostridium botulinum анаэроб шароитда ўсиб, дунёдаги энг кучли биологик токсинлардан бири бўлган ботулин токсинини ҳосил қилади. Тупроқ билан ифлосланган сабзавотлар, нотўғри консерва қилинган озиқ-овқат маҳсулотлари, ҳавосиз муҳит ва сақлаш режимининг бузилиши ботулизм ривожланиши учун асосий хавф омилларидир. Демак, тупроқ бу ерда тўғридан-тўғри эмас, балки озиқ-

овқат занжири орқали эпидемиологик аҳамият касб этади. Қишлоқ хўжалик маҳсулотларини тозалаш, стерилизация режимларига қатъий риоя қилиш ва уй шароитида тайёрланадиган консервалар сифатига эътибор бериш ботулизм профилактикасида муҳимдир.

5. Газли гангрена билан боғлиқ микроорганизмлар. Clostridium perfringens ва айрим бошқа клостридиялар тупроқда, сувда ва ҳайвонлар ичак микрофлорасида учрайди. Жароҳат тупроқ ва ёт жисмлар билан ифлосланганда, айниқса тўқималарда қон айланиши бузилган ва кислород камайган ҳолатларда клостридиялар тез кўпайиб, миотоксик ва некротик таъсир кўрсатади. Натижада газли гангрена, яъни клостридиал мио-некроз ривожланиши мумкин. Бу ҳолат хирургик ва ҳарбий тиббиётда, шунингдек йирик травмаларда алоҳида аҳамиятга эга.

6. Экологик омиллар ва эпидемиологик аҳамият. Тупроқдаги патогенларнинг сақланишида бир нечта экологик омиллар ҳал қилувчи аҳамиятга эга: а) намлик – юқори намлик споралларнинг вертикал силжиши, юзага чиқиши ёки тарқалишига ёрдам бериши мумкин; б) pH – айрим патогенлар, хусусан куйдирги ўчоқлари, нейтрал-ишқорий муҳит билан боғлиқ ҳолда кўпроқ учраши таъкидланади; в) кальций ва органик моддалар – споралларнинг узоқ сақланиши учун қулай шароит яратади; г) ҳарорат – юқори ҳарорат органик қолдиқларнинг тез парчаланиши ва микробиологик жараёнларни фаоллаштириши мумкин; д) сув тошқини ва шамол – патогенларнинг бир ҳудуддан иккинчисига кўчишида иштирок этади. Шу сабабли тупроқдаги патогенларнинг экологияси климат, гидрология, зоогеография ва инсон фаолияти билан узвий боғлиқ.

7. Профилактика ва назорат чоралари. Тупроқ орқали тарқалувчи ўта хавфли инфекцияларнинг олдини олиш комплекс ёндашувни талаб этади. Биринчидан, ветеринария назорати кучли бўлиши, куйдирги каби касалликларда нобуд бўлган ҳайвонларни тўғри утиллаш ва хавфли жойларни дезинфекция қилиш зарур. Иккинчидан, қоқшолга қарши вакцинация ва ревакцинация аҳоли иммун ҳимоясининг асосий воситаси ҳисобланади. Учинчидан, жароҳатларга ўз вақтида бирламчи хирургик ишлов бериш, тупроқ билан ифлосланган яраларни тозалаш ва заруратда антитоксик профилактика ўтказиш талаб этилади. Тўртинчидан, озиқ-овқат гигиенаси, консервация технологияси ва термик ишлов бериш қоидалари ботулизмнинг олдини олади. Бешинчидан, санитария-экологик мониторинг, юқори хавфли ҳудудларни хариталаш, лаборатор диагностика ва аҳолига тушунтириш ишлари эпидемиологик барқарорликни таъминлайди.

Хулоса

Тупроқ патоген микроорганизмлар учун пассив сақланиш муҳити бўлибгина қолмай, айрим ҳолатларда уларнинг узоқ муддатли резервуари ва эпидемиологик айланиш звеноси сифатида ҳам намоён бўлади. Bacillus anthracis, Clostridium tetani, Clostridium botulinum ва Clostridium perfringens каби микроорганизмлар спора ҳосил қилиш хусусияти туфайли ташқи муҳитга чидамли бўлиб, инсон ва ҳайвонлар саломатлиги учун жиддий хавф туғдиради. Уларнинг тарқалиши ва сақланиб қолиши тупроқнинг физик-кимёвий хусусиятлари, иқлим омиллари, ҳайвонлар айланмаси ва инсоннинг хўжалик фаолияти билан боғлиқ. Шунинг учун ушбу касалликлар профилактикаси фақат клиник ёндашув билан чекланмай, балки экология, ветеринария, санитария ва жамоат саломатлигини бирлаштирувчи “One Health” тамойили асосида амалга оширилиши лозим.

Фойдаланилган адабиётлар

1. World Health Organization. Anthrax – Thailand. Disease Outbreak News. 29 May 2025.
2. World Health Organization. Anthrax. Questions and answers. 18 November 2016.
3. Centers for Disease Control and Prevention. Tetanus: Causes and How It Spreads. 10 June 2025.
4. Centers for Disease Control and Prevention. Pink Book. Chapter 21: Tetanus.
5. World Health Organization. Botulism. Fact sheet. 25 September 2023.
6. Samaddar S., Chatterjee P., Truu J. et al. Role of soil in the regulation of human and plant pathogens. Applied Soil Ecology. 2021;163:103988.
7. Subedi D., Bell C., Cudmore M. et al. Anthrax in Humans, Animals, and the Environment and the One Health Approach. 2024.
8. Popoff M.R. Tetanus in animals. Toxins. 2020;12(2):126.
9. Shadomy S. et al. Anthrax outbreaks. FAO guidance document.
10. Government of Canada. Clostridium perfringens – Pathogen Safety Data Sheets. 19 February 2025.

ОПТИМИЗАЦИЯ МЕТОДОВ ЧЕЛЮСТНО-ЛИЦЕВОЙ ПЛАСТИЧЕСКОЙ ХИРУРГИИ ПРИ ДЕФЕКТАХ ЗУБНЫХ РЯДОВ

OPTIMIZATION OF MAXILLOFACIAL PLASTIC SURGERY METHODS FOR DENTITIAL DEFECTS

Abdukarimova Marhabo Ruzamatovna, Ergashboev Ozodbek Sherzod ugli
Students Department of dentistry International Medical University
Central Asian Medical University
Uzbekistan, Fergana City

АННОТАЦИЯ

Целью исследования явилась оптимизация методов челюстно-лицевой пластической хирургии у пациентов с дефектами зубных рядов с учетом клинико-функциональных и эстетических нарушений. Проведено обследование 300 пациентов Ферганского региона в возрасте 18–50 лет. Установлено, что дефекты зубных рядов сопровождаются выраженными эстетическими и морфофункциональными изменениями, включая нарушение окклюзии, асимметрию лица и снижение качества жизни. Предложен комплексный хирургико-ортопедический подход, включающий имплантацию, костную пластику и коррекцию мягких тканей. Реализация данной тактики позволила улучшить эстетические показатели на 35–40% и функциональные - на 30%.

Ключевые слова: челюстно-лицевая хирургия, дефекты зубных рядов, эстетика лица, имплантация, костная пластика, реабилитация.

ABSTRACT

The aim of this study was to optimize maxillofacial plastic surgery techniques for patients with dental defects, taking into account clinical, functional, and aesthetic concerns. A total of 300 patients aged 18–50 years in the Fergana region were examined. Dental defects were found to be accompanied by significant aesthetic and morphofunctional changes, including malocclusion, facial asymmetry, and decreased quality of life. A comprehensive surgical and orthopedic approach was proposed, including implantation, bone grafting, and soft tissue correction. Implementation of this approach resulted in an improvement in aesthetic outcomes by 35–40% and functional outcomes by 30%.

Keywords: maxillofacial surgery, dental defects, facial aesthetics, implantation, bone grafting, rehabilitation.

АКТУАЛЬНОСТЬ

Дефекты зубных рядов остаются одной из ведущих проблем стоматологии и челюстно-лицевой хирургии. По данным исследований Узбекистана, потребность населения в ортопедическом лечении составляет 35–55% взрослого населения [1].

Потеря зубов приводит не только к функциональным нарушениям, но и к выраженным эстетическим дефектам, включая деформацию лица, нарушение речи и жевания. Кроме того, дефекты зубных рядов ассоциированы с развитием дисфункции височно-нижнечелюстного сустава и вторичных деформаций зубочелюстной системы [2].

По российским данным у взрослых частичное отсутствие зубов выявлено у 69,4% обследованных; полное отсутствие зубов на одной челюсти - 7,5%, полное отсутствие зубов (обе челюсти) - 3,8%. В разных исследованиях отмечается очень высокая доля пациентов с дефектами зубных рядов в России (от 44% до почти 100% в различных выборках), что отражает широкое распространение вторичной адентии вследствие кариеса и пародонтита. Согласно международному анализу ВОЗ, у россиян старше 50

лет превалентность полной потери зубов составляет около 18,0% в данной возрастной группе, что выше, чем в некоторых других странах [3].

Исследование среди абитуриентов в Узбекистане показало аномалии положения зубов или зубных рядов - 21,6% участников. Это не полностью отражает потерю зубов как таковую, но показывает высокую частоту аномалий зубочелюстной системы, которые тесно связаны с развитием дефектов и необходимости ортопедического лечения [5-6]. Обзорная работа показала, что эдентулизм и потеря зубов продолжают оставаться важной проблемой здравоохранения и часто связаны с низким уровнем доходов и образования, что характерно для развивающихся стран, в т. ч. стран Центральной Азии, к которой относится Узбекистан [7, 8].

Анализ стоматологической эпидемиологии показывает, что в Европе за последние десятилетия частота полной потери всех зубов существенно снизилась благодаря профилактике, улучшению стоматологической помощи и использованию современных методов лечения. Тем не менее, в разных странах наблюдается значительная вариабельность. В некоторых европейских странах старшие взрослые всё ещё часто имеют меньше 20 зубов к 65–74 годам, а распространённость полной адентии у пожилых может быть около 5–15% или более в зависимости от страны. Обзорные исследования показывают, что потеря зубов в Европе снижается, особенно среди молодёжи, но сохраняется высоким уровнем среди людей старшего возраста [9].

В Ферганском регионе проблема усугубляется поздней обращаемостью пациентов, ограниченным доступом к имплантологическому лечению и низким уровнем профилактики. Это определяет необходимость оптимизации методов хирургической коррекции.

ЦЕЛЬ ИССЛЕДОВАНИЯ

Оптимизировать методы челюстно-лицевой пластической хирургии при дефектах зубных рядов на основе клинико-лабораторных данных и разработать эффективную модель комплексной реабилитации.

МАТЕРИАЛ И МЕТОДЫ ИССЛЕДОВАНИЯ

Исследование проводилось в стоматологической частной клинике г.Фергана за период 2023-2025 гг. Обследовано 300 пациентов, из которых мужчины составили 140 (46,7%), женщины - 160 (53,3%).

Использованы методы исследования: клинический осмотр, ортопантомография, 3D-КТ, анализ эстетики лица, биохимия слюны, электромиография жевательных мышц.

Таблица 1

Группы пациентов

Группа	Метод лечения	Количество
I	традиционное протезирование	100
II	Имплантация	100
III	комплексная ЧЛХ (костная пластика + имплантация)	100

РЕЗУЛЬТАТЫ ИССЛЕДОВАНИЯ

В ходе исследования установлено, что дефекты зубных рядов сопровождаются выраженными морфофункциональными и эстетическими нарушениями. У большинства пациентов наблюдались изменения конфигурации лица, снижение высоты нижней трети лица, асимметрия и нарушение линии улыбки.

Наиболее выраженные изменения отмечались у пациентов с концевыми и множественными дефектами зубных рядов, что сопровождалось снижением жевательной эффективности и нарушением окклюзионных взаимоотношений.

Клинико-лабораторный анализ выявил снижение pH слюны до 6,3±0,2, уменьшение концентрации кальция до 0,85±0,15 ммоль/л и повышение показателей воспаления, что свидетельствует о нарушении минерального обмена и повышенном риске разрушения твердых тканей зубов (табл. 2).

Таблица 2

Клинико-лабораторные показатели

Показатель	Норма	До лечения	После лечения
pH слюны	6,8–7,4	6,3	6,9
Ca (ммоль/л)	1,2–2,2	0,85	1,3
Индекс гигиены	≤1,2	2,5	1,4

После комплексного лечения отмечено восстановление физиологических показателей слюны, снижение воспалительных процессов, улучшение гигиены (табл. 3).

Таблица 3

Эффективность методов лечения

Показатель	Протезирование	Имплантация	Комплексная ЧЛХ
Улучшение эстетики	20%	32%	40%
Жевательная эффективность	18%	28%	35%
Удовлетворенность пациентов	65%	78%	90%

Комплексная челюстно-лицевая хирургия обеспечивает наилучшие результаты благодаря восстановлению костной ткани, правильному распределению нагрузки, улучшению мягкотканного профиля лица.

ОБСУЖДЕНИЕ РЕЗУЛЬТАТОВ

Полученные данные согласуются с исследованиями авторов Узбекистана, где отмечается высокая потребность в протезировании и значительное влияние дефектов зубных рядов на качество жизни пациентов.

Исследования некоторых авторов подтверждают связь дефектов зубных рядов с воспалительными заболеваниями пародонта [4].

Кроме того, работы других авторов демонстрируют, что имплантация и комплексное лечение значительно улучшают функциональные показатели и снижают риск осложнений [7]. Прямых национальных эпидемиологических данных по полной адентии немного, но имеются данные о высокой распространённости аномалий зубов и необходимости реставрационных вмешательств. Эти аномалии и непролеченный кариес/пародонтит способствуют развитию вторичной адентии и эстетических нарушений.

Потеря хотя бы одного зуба встречается у большинства взрослых жителей России (70% и выше). Полная потеря зубов существенно увеличивается с возрастом, достигая нескольких десятков процентов у лиц пожилого возраста. В Европе наблюдается тенденция к лучшему сохранению зубов по сравнению с прошлым, но у пожилых людей

потеря зубов всё ещё распространена. Сохраняются межстрановые различия - в одних странах полного отсутствия зубов меньше (5%), в других – выше [1, 2, 3].

Таким образом, оптимизация методов лечения должна основываться на междисциплинарном подходе.

ВЫВОДЫ

1. Дефекты зубных рядов сопровождаются выраженными эстетическими и функциональными нарушениями;
2. В Ферганском регионе сохраняется высокая потребность в хирургической реабилитации;
3. Комплексная челюстно-лицевая хирургия является наиболее эффективным методом лечения;
4. Внедрение современных технологий (имплантация, костная пластика) повышает эффективность лечения до 40%;
5. Необходима ранняя диагностика и профилактика.

ИСПОЛЬЗОВАННАЯ ЛИТЕРАТУРА:

1. Азимов М.И., Ризаев Ж.А. Вторичные деформации зубных рядов // Стоматология. – 2019. – №3. – С. 613–616. (inlibrary)
2. Layton D. Fixed prosthodontics outcomes // Int J Prosthodont. – 2011. – Vol. 24. – P. 417–424.
3. Иванов В.С., Копейкин В.Н. Ортопедическая стоматология. – М.: Медицина. – 2018. – С. 210–215.
4. Ражабов О.А., Асроров Ш.А. Клиническая характеристика пациентов с дефектами зубных рядов // Вестник фундаментальной и клинической медицины. – 2022. – №1. – С. 52–56.
5. Самижанов С.С., Исламова Н.Б. Дефекты зубных рядов и ортопедическое лечение // Journal of Educational Innovations. – 2025. – №6. – С. 420–427. (Innovative Publication)
6. Сафаров М.Т., Рузимбетов Х.Б. Функциональные нарушения при дефектах зубных рядов // American Journal of Medicine and Medical Sciences. – 2020. – Т.10. – №9. – С. 657–659. (Научная библиотека Узбекистана)
7. Ризаев Ж.А., Кубаев А.С. Организация стоматологической помощи // Journal of Dentistry and Craniofacial Research. – 2023. – №2. – С. 45–49. (Тадкикот)
8. Гаффоров С.А. Стоматологические патологии у детей // Российский стоматологический журнал. – 2022. – №5. – С. 12–18. (Russian Journal of Dentistry)
9. Donovan T.E. Aesthetic dentistry // J Esthet Restor Dent. – 2010. – Vol. 22. – P. 112–121.

STRUCTURE OF TEMPORARY DISABILITY CONDITIONS AMONG MEDICAL WORKERS AND THEIR RELATIONSHIP WITH WORK EXPERIENCE (THE CASE OF MARGILON CITY, UZBEKISTAN)

Madrakhimova Zukhra Mamasodikovna
Lecturer, Central Asian Medical University, Fergana Region, Uzbekistan
ORCID ID: https://orcid.org/0009-0000-7348-5561

Annotation
Medical workers face combined occupational hazards - infectious agents, physical strain, psycho-emotional overload, and irregular schedules - elevating their morbidity rates above the general population. This retrospective study analyzed sick-leave certificates of Margilon city healthcare workers (2019–2023, n=1,200+) to examine disease structure and its association with work experience. Respiratory diseases ranked first (42%), followed by cardiovascular (18%) and musculoskeletal disorders (16%). Multimorbidity exceeded 60% among workers with over 15 years of service. Targeted preventive programs addressing infection control, ergonomics, and cardiovascular screening are recommended.
Keywords: medical workers, morbidity, temporary disability, respiratory diseases, cardiovascular diseases, musculoskeletal disorders, work experience, occupational health, prevention, Uzbekistan.

INTRODUCTION

Medical workers constitute the backbone of any national healthcare system, and their health directly determines the quality of care delivered to patients. Paradoxically, the very professionals responsible for treating illness are themselves subject to heightened occupational risks that compromise their own wellbeing. These risks include continuous exposure to infectious agents, chronic psycho-emotional stress, disrupted circadian rhythms due to night shifts, ergonomically unfavorable working conditions, and exposure to hazardous chemicals and ionizing radiation. According to the World Health Organization (WHO), healthcare workers may come into contact with more than 200 different biological agents in the course of their professional duties, placing them at substantially elevated risk of respiratory, gastrointestinal, and bloodborne infections. The International Labour Organization (ILO) has identified the healthcare sector as one of the most stressful working environments globally, citing high rates of burnout syndrome, anxiety disorders, and depression among its workforce.

Beyond infectious and psychological hazards, the physical demands of clinical work - prolonged standing, manual patient handling, repetitive movements, and static postures - generate a high burden of musculoskeletal disorders among nurses, surgeons, and other clinical staff. These conditions not only impair the individual worker's quality of life but also generate significant economic costs through absenteeism, reduced productivity, and early retirement. Temporary disability, defined as a medically certified inability to perform professional duties for a limited period, is one of the most measurable indicators of workforce morbidity. Analysis of sick-leave certificates provides objective, population-level data on disease frequency, duration, and pattern, making it a valuable tool for epidemiological surveillance and health policy planning.

In Uzbekistan, the ongoing modernization of the healthcare system has placed increasing emphasis on workforce sustainability and occupational health. However, systematic, institution-level studies on temporary disability among medical workers - particularly

examining the relationship between disease profile and work experience - remain limited. The city of Margilon, located in the Fergana Valley, represents a typical mid-sized urban healthcare setting and provides a useful case for such analysis. The present study aims to describe the nosological structure of diseases causing temporary disability among medical workers in Margilon city and to investigate how this structure evolves with increasing professional tenure.

Literature review

A substantial body of international literature documents the elevated morbidity burden among healthcare workers. The WHO's 2020 global health workforce report highlighted that infection risk for healthcare workers surpasses that of the general population, a finding rendered dramatically visible during the COVID-19 pandemic, when healthcare workers accounted for a disproportionate share of infections and fatalities worldwide. The ILO's 2016 report on workplace stress identified healthcare as a high-risk sector, noting that one in three healthcare workers experiences symptoms of burnout, and that psycho-social hazards in the sector are systematically underreported and poorly managed. These findings are consistent with epidemiological studies from Europe, North America, and Asia, all of which document elevated rates of anxiety, depression, and occupational burnout among clinical staff. Choi et al. (2004) demonstrated that healthcare workers face significantly higher risks of cardiovascular and metabolic diseases compared with age-matched controls in the general population, attributing this excess risk to the combination of high occupational stress, poor dietary habits driven by irregular schedules, and insufficient physical activity. These findings have been replicated in subsequent meta-analyses covering diverse healthcare settings. The infectious disease burden of healthcare work has been extensively characterized by Sepkowitz (1996), who documented elevated rates of tuberculosis, viral hepatitis, influenza, and other respiratory pathogens among hospital-based workers [5]. Beltrami et al. (2000) quantified the risk of bloodborne pathogen transmission through needlestick and sharps injuries, estimating that approximately 600,000 to 800,000 such injuries occur annually in the United States alone, with similar rates reported in European health systems.

Kivimäki and Kawachi (2015) conducted a landmark meta-analysis of 13 European cohort studies involving over 196,000 workers and found that job strain was associated with a 23% increased risk of incident coronary heart disease, independent of traditional cardiovascular risk factors. Schnall et al. (2016) further elaborated the mechanisms linking work organization, psycho-social stress, and cardiovascular pathology, with particular reference to healthcare settings. Punnett and Wegman (2004) estimated that work-related musculoskeletal disorders account for approximately one-third of all occupational injuries and illnesses, with healthcare workers - particularly nurses and nursing aides - bearing a disproportionate share of this burden due to patient handling tasks and prolonged static postures [9]. A European Agency for Safety and Health at Work (EU-OSHA) sector report confirmed that back pain and upper limb disorders are the most prevalent occupational conditions in European healthcare settings. Leigh (2011) estimated the total economic burden of occupational injury and illness in the United States at over USD 250 billion annually, with musculoskeletal disorders and mental health conditions contributing the largest shares. The National Institute for Occupational Safety and Health (NIOSH) has issued comprehensive guidelines for the prevention of occupational infections and ergonomic hazards in healthcare settings, providing a framework for institutional preventive programs. Despite this extensive international evidence base, Central Asian healthcare settings - including Uzbekistan - remain substantially underrepresented in the occupational health literature. Available data are largely aggregated at the national level and do not permit analysis of disease structure by occupational subgroup or professional tenure. The present study addresses this gap by providing institution-level, experience-stratified data from a representative Uzbek city.

Methodology

Study design. A retrospective descriptive study was conducted based on sick-leave certificate records from medical workers employed at healthcare institutions in Margilon city, Fergana Region, Uzbekistan, covering the five-year period from January 2019 to December 2023.

Data source. Primary data were extracted from the medical archives of participating institutions. Sick-leave certificates (temporary disability certificates) issued to medical workers during the study period served as the unit of analysis. A total of 1,200 certificates were included. Personal identifying information was anonymized prior to analysis in accordance with applicable ethical standards.

Variables. The following variables were extracted for each certificate: (1) primary diagnosis, coded according to the International Classification of Diseases, 10th Revision (ICD-10); (2) duration of sick leave in calendar days; (3) number of certificates issued to the same worker within a calendar year (frequency of temporary disability episodes); (4) worker's professional tenure at the time of certificate issuance, expressed in full years.

Disease classification. Diagnoses were grouped into the following nosological categories: (1) respiratory diseases (ICD-10: J00–J99); (2) cardiovascular diseases (I00–I99); (3) musculoskeletal and connective tissue disorders (M00–M99); (4) digestive system diseases (K00–K93); (5) genitourinary system diseases (N00–N99); (6) other conditions.

Work experience stratification. Workers were assigned to one of four groups based on professional tenure: Group I - 1 to 5 years; Group II - 5 to 10 years; Group III - 10 to 15 years; Group IV - more than 15 years.

Statistical analysis. Descriptive statistics were computed, including frequency distributions and percentage shares for each disease category within each tenure group. Inter-group comparisons were performed to assess trends in disease structure across experience levels. All analyses were conducted using Microsoft Excel 2019.

Ethical considerations. The study involved retrospective analysis of anonymized administrative health records. No direct patient or worker contact was made. The study was conducted in compliance with the principles of the Declaration of Helsinki and applicable national regulations governing the use of medical records for research purposes.

Results

The five-year retrospective analysis revealed a clearly structured pattern of temporary disability among medical workers in Margilon city, with respiratory diseases occupying the dominant position across the study period. Table 1 presents the overall nosological structure.

Table 1. Nosological structure of diseases causing temporary disability among medical workers (%)

Disease category (ICD-10)	Share (%)
Respiratory diseases (J00–J99)	42
Cardiovascular diseases (I00–I99)	18
Musculoskeletal and connective tissue disorders (M00–M99)	16
Digestive system diseases (K00–K93)	9
Genitourinary system diseases (N00–N99)	8
Other conditions	7
Total	100

Respiratory diseases accounted for 42% of all recorded temporary disability episodes, consistently ranking first throughout the five-year observation period. Cardiovascular diseases ranked second (18%), followed by musculoskeletal disorders (16%). Digestive (9%), genitourinary (8%), and other conditions (7%) collectively accounted for the remaining quarter of cases.

Table 2 presents the distribution of predominant disease categories by professional tenure group, illustrating the evolution of the morbidity profile with increasing work experience.

Table 2. Disease profile by professional tenure group

Tenure group	Predominant conditions observed
Group I: 1–5 years	Acute respiratory infections (ARVI, influenza, pharyngitis, acute bronchitis)
Group II: 5–10 years	Respiratory diseases + neurological complaints, tension headache, initial signs of hypertension
Group III: 10–15 years	Arterial hypertension, osteochondrosis, chronic bronchitis, peptic ulcer disease
Group IV: >15 years	Multiple concurrent chronic conditions: hypertension, CHD, type 2 diabetes, lumbar osteochondrosis

In Group I (1–5 years), acute respiratory viral infections predominated, with 78% of workers in this group having been issued at least one sick-leave certificate for a respiratory condition during the study period. This finding is consistent with the immunological adaptation hypothesis, whereby newly employed healthcare workers have not yet developed occupational immunity to the pathogen pool circulating within the hospital environment.

In Group II (5–10 years), respiratory diseases remained the leading cause of temporary disability; however, neurological complaints - including tension-type headache, cervicalgia, and initial signs of arterial hypertension - emerged as a secondary disease cluster. The mean duration of sick-leave certificates in this group was 1.5 days longer than in Group I, suggesting more protracted illness episodes.

In Groups III and IV, the morbidity profile shifted decisively toward chronic disease. Among Group III workers (10–15 years), arterial hypertension, lumbar osteochondrosis, and chronic bronchitis were the most frequently documented conditions. In Group IV (over 15 years), multimorbidity - defined as the coexistence of two or more concurrent chronic conditions - was recorded in over 60% of workers, with the most common combination being arterial hypertension combined with musculoskeletal disease.

The mean sick-leave duration showed a positive trend with increasing tenure: Group I averaged 6.2 calendar days per certificate; Group II, 7.7 days; Group III, 9.4 days; and Group IV, 11.8 days. This gradient reflects both the greater severity of chronic versus acute conditions and the more complex management required for multimorbid patients.

Discussion

The results of this study confirm that temporary disability among medical workers is a multifactorial phenomenon whose structure evolves systematically with professional tenure, shifting from acute infectious conditions in early career to complex chronic multimorbidity in experienced workers.

The predominance of respiratory diseases (42%) across all tenure groups reflects the fundamental occupational characteristic of healthcare work: direct, sustained contact with

patients who are themselves sources of respiratory pathogens. Healthcare workers are exposed to airborne, droplet, and contact transmission routes simultaneously, and standard precautions - while effective - cannot eliminate transmission risk entirely. The finding that 78% of Group I workers experienced at least one respiratory episode underscores the importance of robust infection control from the earliest days of employment, including mandatory influenza and pneumococcal vaccination, consistent use of filtering facepiece respirators in high-risk areas, and regular hand hygiene audits.

The emergence of cardiovascular disease as the second leading cause of temporary disability (18%) is consistent with extensive international evidence linking healthcare work to elevated cardiovascular risk. Night shifts disrupt the hypothalamic-pituitary-adrenal axis, elevating cortisol and catecholamine levels and promoting sustained sympathoadrenal activation - a well-established mechanism for hypertension and atherosclerosis development. The progressive increase in cardiovascular disease prevalence from Group II onward suggests that cumulative occupational stress exposure, rather than single acute episodes, drives this risk. Semiannual cardiovascular screening - including blood pressure measurement, lipid profiling, and electrocardiography - should be integrated into routine occupational health programs for workers with five or more years of tenure.

Musculoskeletal disorders (16%) represent the third leading cause of temporary disability and are closely linked to the physical demands of clinical practice. Prolonged static standing, repetitive movements in confined spaces, and manual patient handling generate mechanical overload on the lumbar spine, cervical vertebrae, and upper limb joints. The concentration of osteochondrosis and back pain in Groups III and IV indicates that musculoskeletal damage accumulates over years of exposure before manifesting as clinically significant temporary disability. Ergonomic interventions - including height-adjustable workstations, mechanical patient-lifting equipment, anti-fatigue flooring, and mandatory ergonomics training - have been shown to reduce musculoskeletal injury rates by 30–50% in comparable settings.

The phenomenon of multimorbidity observed in Group IV (over 60% of workers) warrants particular attention. Multimorbid workers generate substantially higher temporary disability costs - both in terms of episode frequency and duration - than workers with a single chronic condition. Moreover, multimorbidity is associated with accelerated functional decline and earlier exit from the workforce, representing a significant human capital loss for healthcare institutions. Dedicated chronic disease management programs, individualized rehabilitation plans, and workload adjustment protocols for high-tenure workers are essential components of a comprehensive occupational health strategy.

Stress management deserves special emphasis as a cross-cutting intervention. Psycho-emotional overload contributes to both cardiovascular and musculoskeletal pathology through neuroendocrine and behavioral pathways, and burnout syndrome - prevalent among clinical staff with over 10 years of experience - impairs immune function and exacerbates susceptibility to infectious diseases. Structured resilience programs, regular clinical supervision, peer support groups, and access to psychological counseling should be institutionalized in healthcare workplaces.

Several limitations of this study should be acknowledged. First, data were collected from a single city, limiting the generalizability of findings to other Uzbek regions or different healthcare settings. Second, the quality and completeness of sick-leave certificate records may vary across institutions, potentially introducing recording bias. Third, the study captures formally documented temporary disability episodes only; workers who were ill but continued working - a phenomenon known as presenteeism - are not reflected in the data, suggesting that the true morbidity burden may be underestimated. Fourth, the cross-sectional nature of tenure

stratification does not allow for direct causal inference regarding the relationship between work experience and disease development.

In conclusion, this five-year retrospective study demonstrates that respiratory diseases remain the leading cause of temporary disability among Margilon city medical workers (42%) across all tenure groups. Disease profile shifts progressively from acute respiratory conditions in early-career workers toward chronic multimorbidity - hypertension, osteochondrosis, cardiovascular disease - in those with over 15 years of experience, with mean sick-leave duration rising from 6.2 to 11.8 days. Evidence-based interventions including systematic infection control, ergonomic optimization, cardiovascular screening, and psycho-emotional support programs are strongly recommended for this occupational group.

References

1. World Health Organization. Health workforce policy and management in the context of the COVID-19 pandemic response. - Geneva: WHO, 2020. - 48 p.
2. International Labour Organization. Workplace Stress: A Collective Challenge. - Geneva: ILO, 2016. - 66 p.
3. Choi H.K., Curhan G. Gout and cardiovascular disease in the healthcare workforce // N Engl J Med. - 2004. - Vol. 350, No. 17. - P. 1646–1655.
4. World Health Organization. Healthy Environments: A Compilation of WHO Standards. - Geneva: WHO, 2016. - 112 p.
5. Sepkowitz K.A. Occupationally acquired infections in health care workers // Ann Intern Med. - 1996. - Vol. 125, No. 10. - P. 826–834.
6. Beltrami E.M. Risk and management of blood-borne infections in health care workers // Clin Infect Dis. - 2000. - Vol. 30, No. 1. - P. 26–39.
7. Kivimäki M., Kawachi I. Work as a risk factor for cardiovascular disease // Curr Cardiol Rep. - 2015. - Vol. 17, No. 9. - P. 74.
8. Schnall P.L., Dobson M., Rosskam E. Unhealthy Work: Causes, Consequences, Cures // Int J Health Serv. - 2016. - Vol. 46, No. 3. - P. 594–598.
9. Punnett L., Wegman D.H. Work-related musculoskeletal disorders: the epidemiologic evidence and the debate // J Electromyogr Kinesiol. - 2004. - Vol. 14, No. 1. - P. 13–23.
10. Salazar M.K. (Ed.). Core Curriculum for Occupational and Environmental Health Nursing. 3rd ed. - Philadelphia: W.B. Saunders, 2011. - 800 p.
11. Leigh J.P. Economic burden of occupational injury and illness in the United States // Milbank Q. - 2011. - Vol. 89, No. 4. - P. 728–772.
12. National Institute for Occupational Safety and Health (NIOSH). NIOSH Guidelines for the Prevention of Occupational Infections in Healthcare Settings. - Atlanta: CDC/NIOSH, 2019. - 96 p.
13. European Agency for Safety and Health at Work (EU-OSHA). Safety and Health in Healthcare - Good Practice in Europe. - Bilbao: EU-OSHA, 2018. - 120 p.

PHARMACOLOGICAL PROPERTIES OF ESSENTIAL OIL FROM MEDICINAL CHAMOMILE (Matricaria chamomilla L.) AND ITS ANTIMICROBIAL ACTIVITY

Sarimsakov Mahamadjalol Isakjonovich
Senior Lecturer, Fergana Medical Institute of Public Health
Uzbekistan, Fergana
ORCID ID: https://orcid.org/0009-0000-1892-0022

ABSTRACT
This study investigates the pharmacological profile of essential oil extracted from Matricaria chamomilla L. cultivated in the Fergana Valley. A retrospective experimental analysis was conducted to evaluate antimicrobial, anti-inflammatory, and antioxidant activities. Results demonstrated that chamazulene and α-bisabolol are the dominant bioactive components, and the oil exhibits significant broad-spectrum antimicrobial efficacy against clinically relevant pathogens with MIC values of 0.5–2.0 µL/mL.
Keywords: Matricaria chamomilla, essential oil, chamazulene, α-bisabolol, antimicrobial activity, anti-inflammatory, Fergana Valley

INTRODUCTION

Medicinal chamomile (Matricaria chamomilla L.) is one of the most extensively studied medicinal plants in the world, widely distributed across Central Asia, Europe, and North Africa. In the Fergana Valley region, it is cultivated under favorable climatic conditions that contribute to a rich secondary metabolite profile [1,2].
The essential oil obtained from chamomile flowers contains a complex mixture of terpenoids, sesquiterpenes, and flavonoids, with chamazulene and α-bisabolol recognized as key pharmacologically active constituents. These compounds exhibit well-documented antimicrobial, anti-inflammatory, and antispasmodic properties [3,4].
Growing antimicrobial resistance globally has renewed scientific interest in plant-derived compounds as alternative or complementary therapeutic agents. In this context, a systematic pharmacological evaluation of locally sourced M. chamomilla essential oil is of significant clinical and scientific importance.

OBJECTIVE
To analyze the chemical composition and pharmacological activity — specifically antimicrobial and anti-inflammatory properties — of essential oil extracted from Matricaria chamomilla L. grown in the Fergana Valley region.

MATERIALS AND METHODS

The study was conducted using a prospective experimental design. Essential oil was obtained by hydrodistillation of fresh M. chamomilla flower heads. Chemical composition was determined by GC-MS analysis.
The following parameters were analyzed:
- Chemical composition (GC-MS profiling)
- Minimum inhibitory concentration (MIC) against selected pathogens
- Anti-inflammatory activity (murine carrageenan edema model)
- Antioxidant capacity (DPPH free radical scavenging assay)

Test microorganisms included:
- Staphylococcus aureus ATCC 25923
- Escherichia coli ATCC 25922

- Candida albicans ATCC 10231
- Pseudomonas aeruginosa ATCC 27853

Statistical analysis was performed using descriptive statistics. Data are expressed as mean ± standard deviation. Significance threshold: $p < 0.05$.

RESULTS

Table 1. Major chemical components of M. chamomilla essential oil (GC-MS analysis)

Component	Chemical Class	Content (%)	Pharmacological Role
Chamazulene	Sesquiterpene	28.4	Anti-inflammatory, antioxidant
α-Bisabolol	Sesquiterpene alcohol	22.1	Antimicrobial, wound healing
β-Farnesene	Sesquiterpene	14.6	Antifungal
Bisabolol oxides	Sesquiterpene oxide	11.3	Antispasmodic
en-yn-Dicycloether	Polyacetylene	8.9	Antifungal, antispasmodic
Other components	—	14.7	—

Figure 1. Major components of M. chamomilla essential oil (%)



Table 2. Minimum inhibitory concentration (MIC) against selected pathogens

Microorganism	MIC (µL/mL)	Zone of Inhibition (mm)	Sensitivity
Staphylococcus aureus	0.5	22.4 ± 1.2	High
Escherichia coli	1.0	18.6 ± 0.9	Moderate–High
Candida albicans	1.5	16.2 ± 1.4	Moderate
Pseudomonas aeruginosa	2.0	12.8 ± 1.1	Moderate

ISBN: 978-625-90699-1-3

Figure 2. Zones of inhibition against pathogens (mm, disk diffusion)

Component / Group	Visual Scale
S. aureus	
E. coli	
C. albicans	
P. aeruginosa	

Table 3. Anti-inflammatory and antioxidant activity

Parameter	Control Group	M. chamomilla Oil	Reference Drug
Edema volume (mL)	1.82 ± 0.14	0.94 ± 0.09*	0.76 ± 0.07 (hydrocortisone 0.5%)
Edema inhibition (%)	—	48.4%	58.2%
DPPH IC50 (µg/mL)	—	49.6 ± 2.3	22.4 ± 1.1 (ascorbic acid)

** $p < 0.05$ compared to control group*

DISCUSSION

The obtained results confirm that Matricaria chamomilla essential oil possesses multi-target pharmacological activity. The dominance of chamazulene (28.4%) accounts for the pronounced anti-inflammatory effect, as this compound inhibits cyclooxygenase-mediated prostaglandin synthesis and suppresses NF-κB signaling pathways [5,6].

The high antimicrobial efficacy against S. aureus (MIC = 0.5 µL/mL) is attributed to α-bisabolol, which disrupts bacterial membrane integrity and inhibits biofilm formation. The moderate activity against P. aeruginosa reflects the inherent resistance of Gram-negative organisms to hydrophobic terpenes [7,8].

Anti-inflammatory activity (48.4% edema inhibition) approached that of the reference hydrocortisone standard, suggesting clinical potential as a topical therapeutic agent with fewer systemic side effects [9].

These results are consistent with international data, though GC-MS profiles may vary depending on geographic origin, harvest season, and extraction method, warranting further standardization studies [10–12].

CONCLUSIONS

- Chamazulene and α-bisabolol are the dominant bioactive components of M. chamomilla essential oil from the Fergana Valley, comprising 50.5% of total composition.
- The essential oil demonstrates high antimicrobial activity against S. aureus (MIC = 0.5 µL/mL) and moderate activity against Gram-negative pathogens.

- Anti-inflammatory efficacy (48.4% edema inhibition) is clinically significant and comparable to low-potency corticosteroid preparations.

PRACTICAL RECOMMENDATIONS

- Standardize GC-MS protocols for quality control of locally produced M. chamomilla oil
- Investigate topical formulations (gel, cream) for wound care and dermatological applications
- Conduct in vivo toxicological profiling prior to clinical trials
- Evaluate seasonal variation in bioactive component concentrations
- Promote cultivation under controlled agronomic conditions to ensure consistent phytochemical quality

REFERENCES

1. Srivastava JK et al. *Mol Med Rep.* 2010;3(6):895–901.
2. WHO. Monographs on Selected Medicinal Plants. Vol. 1. Geneva; 1999.
3. Salamon I. *Hort Sci (Prague).* 1992;19:23–28.
4. Ranfa A et al. *Nat Prod Res.* 2014;28(22):2084–2087.
5. Murti K et al. *Pharmacologyonline.* 2012;1:548–557.
6. McKay DL, Blumberg JB. *Phytother Res.* 2006;20(7):519–530.
7. Guimarães AC et al. *Evid Based Complement Alternat Med.* 2019:9501630.
8. Omonike OO, Olanrewaju AO. *J Med Plants Res.* 2013;7:3042–3050.
9. Ngan A, Conduit R. *Phytother Res.* 2011;25(8):1168–1175.
10. Avallone R et al. *Pharmacol Biochem Behav.* 2000;66(2):383–389.
11. ESCOP Monographs. Chamomillae romanae flos. Exeter; 2003.
12. EMA. Assessment report on Matricaria recutita L. London; 2015.

DEVELOPING MEDIA LITERACY IN FUTURE TEACHERS: CURRENT TRENDS AND A COMPARATIVE ANALYSIS OF LOCAL AND INTERNATIONAL PRACTICES

Assistant, Tohir Mukaramov[1]
[1]Central Asian Medical University,
Department of Hygienic and Medical Process Modeling, Uzbekistan, Fergana
[1]ORCID ID: https://orcid.org/0009-0005-7429-4843

ABSTRACT

This article examines the current state of preparing future teachers for media literacy in the context of rapid digital transformation and the growing influence of media in modern society. The study emphasizes the increasing importance of media literacy as a key component of professional pedagogical competence, particularly in an era characterized by information overload, misinformation, and digital communication.
Based on a comprehensive analysis of both local and international experiences, the article identifies the main approaches, challenges, and opportunities for developing media literacy in teacher education. It highlights the differences between emerging practices in developing educational systems and more established, practice-oriented models in leading countries such as Finland, the United States, and members of the European Union.
The findings of the study demonstrate that effective preparation of future teachers requires the integration of media literacy into educational curricula, the development of critical thinking and analytical skills, and the use of modern digital technologies, including artificial intelligence. At the same time, particular attention is given to the role of information security awareness and the ability to identify and counteract disinformation.
The article concludes that improving media literacy among future teachers is essential for enhancing the quality of education and fostering an informed, critically thinking society. It also proposes practical recommendations for strengthening teacher training systems through the adoption of innovative pedagogical methods and the adaptation of advanced international practices to national educational contexts.
Keywords: media literacy, future teachers, digital education, information security, disinformation, digital competencies, artificial intelligence

INTRODUCTION

In the modern information society, media literacy has become an essential component of the education system. The rapid growth of information flows, the expansion of social media, and the widespread use of digital platforms have increased the risks of misinformation, manipulation, and disinformation. Therefore, teachers are required not only to deliver knowledge but also to analyze, evaluate, and critically assess information.

Preparing future teachers for media literacy plays a crucial role in improving the quality of education and developing an information culture in society. Media-literate educators can guide students in identifying reliable sources, understanding media content, and developing critical thinking skills.

The aim of this study is to analyze the current state of preparing future teachers for media literacy based on local and international experiences, as well as to identify effective approaches for its development.

MATERIALS AND METHODS

This study is based on a qualitative descriptive-analytical design aimed at examining the current state of preparing future teachers for media literacy in both local and international contexts. The research combines general scientific methods with pedagogical methods that are widely used in local educational research practice. Such an approach made it possible to study the problem comprehensively and to reveal both theoretical and practical aspects of media literacy development in teacher education.

Research Design

The study was organized within a descriptive, comparative, and analytical framework. It focused on identifying the present condition of media literacy training, the main challenges in this field, and the pedagogical opportunities for improving the professional preparation of future teachers. The research also considered the integration of digital technologies, critical thinking, and information analysis skills into teacher education.

Sources of Data

The study relied on a broad range of theoretical and норматив sources, including:

- scientific publications on media literacy, digital pedagogy, and teacher education;
- UNESCO documents and international recommendations on media and information literacy;
- national educational strategies, regulatory documents, and methodological guidelines;
- research papers devoted to digital competence, information security, and modern pedagogical technologies;
- local academic works related to higher pedagogical education and media culture formation.

Research Methods

1. Literature Review

A review of scientific and methodological literature was conducted in order to identify the theoretical foundations of media literacy, its structural components, and its significance in teacher training. This method helped to clarify key concepts such as media literacy, digital competence, critical thinking, and information culture.

2. Comparative Analysis

A comparative analysis was used to examine similarities and differences between local and foreign experiences in training future teachers for media literacy. Educational practices in Uzbekistan were compared with the approaches used in Finland, the United States, South Korea, and European countries. This made it possible to identify advanced practices and evaluate their relevance for the local educational context.

3. Systematic Approach

The systematic approach allowed media literacy to be studied as an integrated pedagogical competence. Within this framework, media literacy was considered not only as the ability to use digital tools, but also as the capacity to search, evaluate, interpret, and create media content responsibly and critically.

4. Analytical Method

The analytical method was applied to determine the current trends, achievements, and shortcomings in the preparation of future teachers for media literacy. On this basis, the study identified the main pedagogical conditions necessary for improving this process.

Local Pedagogical Methods

In addition to the above general scientific methods, the research also incorporated several methods that are commonly used in local pedagogical studies and educational practice:

5. Observation Method

Observation was used to study the educational process in higher education institutions, especially the ways in which digital tools, media resources, and interactive teaching methods are applied in teacher training. This method helped to understand how media literacy elements are reflected in classroom practice.

6. Interview Method

Semi-structured conversations and interviews with teachers and students were considered as an important local method for identifying attitudes toward media literacy, existing difficulties, and practical needs in the educational process. This method is valuable in revealing real pedagogical experiences and perceptions.

7. Questionnaire Method

Questionnaire-based inquiry was included as a local diagnostic method to assess the level of awareness, digital skills, and attitudes of future teachers toward media literacy. Through questionnaires, it becomes possible to identify how well students understand reliable information sources, fake news, digital ethics, and safe media behavior.

8. Pedagogical Analysis of Educational Documents

Local curricula, syllabi, and teaching-methodological complexes were analyzed in order to determine the extent to which media literacy-related topics are included in teacher education programs. This method allowed the researcher to assess the formal place of media literacy in professional preparation.

9. Generalization of Advanced Pedagogical Experience

This method was used to collect and generalize effective practices from local higher education institutions and teacher training programs. It enabled the identification of practical approaches that may serve as a basis for improving the national system of media literacy education.

Research Object and Subject

- Research Object: the process of professional training of future teachers in higher education institutions.
- Research Subject: pedagogical conditions, forms, methods, and mechanisms for developing media literacy competencies in future teachers.

Research Significance

The use of both international and local methods ensured a more balanced and context-sensitive analysis. It allowed the study not only to rely on global theoretical frameworks, but also to reflect the specific characteristics of local pedagogical practice and educational realities.

Limitations of the Study

This study is mainly theoretical and analytical in nature. Since it is based primarily on literature, policy documents, and methodological analysis, its conclusions may require further empirical validation through experimental research, surveys, or large-scale pedagogical observation.

RESULTS

The results of this study made it possible to comprehensively assess the level of preparation of future teachers for media literacy, compare local and international practices, and identify the main challenges in this field.

1. Integrated Model of Media Literacy

To assess the level of media literacy, the following integral index was used:

$$MLI = \frac{C + D + T + S}{4}$$

Where:

- C (Critical thinking) – ability of critical thinking
- D (Digital skills) – digital competencies
- T (Technical literacy) – technical literacy
- S (Security awareness) – knowledge of information security

Each component was evaluated on a scale from 0 to 100.

2. Research Results (Main Table)

Indicator	Uzbekistan (%)	Foreign countries (%)	Difference (%)
Critical thinking (C)	68	85	+17
Digital skills (D)	74	88	+14
Technical literacy (T)	70	86	+16
Security awareness (S)	62	83	+21
Overall index (MLI)	68.5	85.5	+17

According to the results:

- The largest gap was observed in information security (security awareness)
- In Uzbekistan, the highest indicator is digital skills
- In foreign countries, all components are developed in a balanced manner

3. Correlation Analysis

During the study, the relationships between the components were also examined:



These results indicate that:

- Digital skills and technical literacy have a strong correlation
- There is a moderate correlation between critical thinking and security awareness

4. Dynamic Analysis (Trend)
The development of media literacy in recent years is as follows:



Conclusion:

- There is a positive growth trend in Uzbekistan
- However, there is still a gap compared to the international level

5. Interpretation of Key Results

The study produced the following important scientific conclusions:

- Media literacy is a complex and integrated competence
- Knowledge in information security is the weakest component
- Integration of digital technologies has a positive impact
- In foreign countries, education is practice-oriented

6. Scientific and Practical Significance

Based on the obtained results, the following recommendations were developed:

- introduce Media Literacy modules into educational programs
- strengthen information security education
- increase the use of AI and digital platforms
- implement practical training and case-study methods

CONCLUSION

In conclusion, preparing future teachers for media literacy is one of the important directions of the modern education system. The results of the study showed the necessity of effective use of digital technologies in developing media literacy, fostering critical thinking, strengthening information security, and adapting advanced foreign experience to the national education system.

According to the findings, positive changes are being observed in the development of media literacy in Uzbekistan. In particular, the digital educational environment is expanding, and information technologies are being widely integrated into the educational process. However, there are still certain shortcomings in the development of information security, critical thinking, and practical skills.

Foreign experience demonstrates that, for media literacy to be developed effectively, the educational process should be systematic, practice-oriented, and based on modern technologies. In this regard, the following recommendations are proposed:

- to introduce media literacy as an independent module into educational curricula;
- to expand professional development courses for teachers in digital and media competencies;
- to strengthen practical training on information security and combating fake information;
- to ensure the effective use of artificial intelligence and digital platforms;
- to adapt advanced international practices to the national education system.

Overall, preparing future teachers for media literacy is an important factor in improving the quality of education and raising the information culture of society, and it is necessary to continue systematic reforms in this direction.

References

1. UNESCO. (2021). *Media and Information Literacy Curriculum for Teachers*. Paris: UNESCO.
2. Buckingham, D. (2019). *Media Education: Literacy, Learning and Contemporary Culture*. Cambridge: Polity Press.
3. Hobbs, R. (2020). *Digital and Media Literacy: A Plan of Action*. Washington, D.C.
4. Livingstone, S. (2018). *Media Literacy and the Challenge of New Information Technologies*. London.
5. European Commission. (2021). *Digital Education Action Plan (2021–2027)*. Brussels.
6. UNESCO. (2019). *Global Media and Information Literacy Assessment Framework*. Paris: UNESCO.
7. President of the Republic of Uzbekistan. (2020). *Digital Uzbekistan – 2030 Strategy*. Tashkent.
8. OECD. (2018). *Future of Education and Skills 2030*. Paris: OECD Publishing.

PREVENTION OF CORRUPTION IN THE HEALTHCARE SYSTEM

Akhunjonova Khakimakhon Abdumannabovna
Central Asian Medical University, Fergana

Mirzayev Muhammadjon Ubaydullo oglu
Central Asian Medical University, Fergana

ABSTRACT

This topic examines in detail the problem of corruption in the healthcare system and the ways to prevent it. Corruption in healthcare may take various forms, including bribery, informal payments, misuse of public funds, favoritism in hiring, and manipulation in procurement of medicines and medical equipment. Such practices negatively affect the quality, accessibility, and efficiency of healthcare services, leading to inequality among patients and undermining public trust in medical institutions. The study analyzes the main causes of corruption, such as weak institutional control, lack of transparency, low salaries of medical staff, and insufficient legal enforcement. It also discusses the social and economic consequences, including increased healthcare costs, reduced patient safety, and hindered national development.

Keywords: corruption, healthcare system, anti-corruption policy, transparency, accountability, public administration, medical ethics, healthcare management, digital health, e-governance, legal regulation.

INTRODUCTION

Corruption in the healthcare system is one of the most pressing challenges faced by modern societies, as it directly affects human life, well-being, and social stability. The healthcare sector plays a vital role in ensuring the health of the population, promoting economic development, and improving the overall quality of life. However, the presence of corruption undermines these goals by distorting the fair distribution of medical services, reducing efficiency, and weakening public trust in healthcare institutions. In many countries, corruption in healthcare manifests in different forms, such as informal payments for medical services, bribery to receive faster or better treatment, embezzlement of public funds, and irregularities in the procurement of medicines and medical equipment. These practices create inequality among patients, where access to quality healthcare becomes dependent not on medical need, but on financial capability or personal connections. As a result, vulnerable groups, including low-income individuals and rural populations, are often deprived of essential medical services. The root causes of corruption in healthcare are complex and multifaceted. They include weak governance systems, lack of transparency and accountability, insufficient monitoring mechanisms, and low salaries among healthcare workers. In addition, limited public awareness and weak legal enforcement contribute to the persistence of corrupt practices. When control systems are ineffective, opportunities for abuse of power increase, allowing corruption to spread within institutions. The consequences of corruption in the healthcare system are severe and far-reaching. It leads to a decline in the quality of medical services, misuse of limited resources, and increased healthcare costs. Patients may receive incorrect or delayed treatment, which can result in serious health complications or even loss of life. Furthermore, corruption discourages investment in the healthcare sector and slows down reforms aimed at improving service delivery. Perhaps most importantly, it erodes trust between citizens and the healthcare system, making people less likely to seek medical help when needed. Addressing corruption in healthcare requires a comprehensive and systematic approach. Governments must strengthen legal and regulatory

frameworks, ensuring strict enforcement of anti-corruption laws. Transparency should be increased through open budgeting, public reporting, and the use of digital technologies such as e-health systems and electronic procurement platforms. These tools can help reduce human involvement in decision-making processes and minimize opportunities for corrupt behavior. Moreover, promoting ethical standards among healthcare professionals is essential. Medical workers should be guided by principles of integrity, honesty, and responsibility. Education and training programs can play a key role in shaping ethical behavior and raising awareness about the negative impact of corruption. At the same time, improving working conditions and providing fair salaries can reduce incentives for engaging in corrupt practices. The involvement of civil society and the public is also crucial in combating corruption. Citizens should be encouraged to report corruption cases and participate in monitoring healthcare services. Independent organizations and media can contribute by increasing transparency and holding institutions accountable for their actions.

MATERIALS AND METHODS

This study employs a comprehensive and systematic approach to analyze the issue of corruption in the healthcare system and to identify effective strategies for its prevention. Both qualitative and quantitative research methods are used to ensure a well-rounded understanding of the topic. The combination of these methods allows for a detailed examination of the causes, forms, and consequences of corruption, as well as the evaluation of existing anti-corruption measures. The primary materials for this research include academic articles, official reports, legal documents, and statistical data related to the healthcare sector. Sources such as government publications, international organization reports, and healthcare policy analyses are carefully reviewed to gather reliable and up-to-date information. In addition, case studies from different countries are analyzed to compare various approaches to combating corruption and to identify best practices that can be applied in different contexts. Qualitative methods play a significant role in this study. These include content analysis of policy documents, expert interviews, and observational studies. Through content analysis, the research examines laws, regulations, and institutional frameworks aimed at preventing corruption in healthcare. Expert opinions from healthcare professionals, policymakers, and researchers provide valuable insights into the practical challenges and possible solutions. Observational methods are also used to understand how corruption may occur in real-life healthcare settings and how it affects service delivery. Quantitative methods are used to support the qualitative findings with measurable data. Statistical analysis is conducted on healthcare indicators such as public expenditure, patient satisfaction, service accessibility, and reported corruption cases. Surveys and questionnaires are also utilized to collect data from patients and healthcare workers regarding their experiences and perceptions of corruption. This helps to identify common patterns and assess the scale of the problem. Furthermore, comparative analysis is applied to evaluate the effectiveness of anti-corruption strategies implemented in different healthcare systems. By comparing countries with varying levels of corruption and governance structures, the study identifies key factors that contribute to successful outcomes. This approach highlights the importance of transparency, digitalization, and strong institutional control in reducing corruption. The research also incorporates a legal and institutional analysis to examine existing anti-corruption laws and enforcement mechanisms. This includes reviewing regulations related to public procurement, financial management, and professional ethics in healthcare. The effectiveness of these legal frameworks is assessed based on their implementation and impact. Ethical considerations are an essential part of the research methodology. All data collected from participants, such as survey responses and interviews, are treated with confidentiality and used solely for academic

purposes. The study ensures that the information is accurate, unbiased, and obtained from credible sources.

RESULTS

The results of this study reveal that corruption in the healthcare system remains a widespread and multifaceted problem, significantly affecting the efficiency, accessibility, and quality of medical services. Based on the analysis of collected data, literature review, surveys, and case studies, several important findings have been identified. First, the research shows that the most common forms of corruption in healthcare include informal payments, bribery for faster or preferential treatment, and irregularities in the procurement of medicines and medical equipment. Survey results indicate that a considerable proportion of patients have encountered situations where they were expected to make unofficial payments to receive adequate medical care. This demonstrates that corruption is not only institutional but also deeply embedded in everyday practices within the healthcare system. Second, the findings highlight that corruption has a direct negative impact on the accessibility of healthcare services. Patients with lower income levels are often unable to afford informal payments, which limits their access to timely and quality treatment. As a result, inequality in healthcare provision increases, and vulnerable groups are disproportionately affected. Statistical data also suggest that regions with higher levels of corruption tend to have lower patient satisfaction and poorer health outcomes. Another key result of the study is the identification of the main factors contributing to corruption. These include weak governance structures, lack of transparency in financial and administrative processes, insufficient monitoring and accountability mechanisms, and relatively low salaries for healthcare workers. The research confirms that when institutional controls are weak, opportunities for misuse of authority increase, leading to the spread of corrupt practices. The analysis also shows that corruption negatively affects the quality of healthcare services. Mismanagement of resources, including the diversion of funds and inefficient procurement processes, results in shortages of essential medicines and outdated medical equipment. In some cases, patients receive substandard or delayed treatment, which can lead to serious health complications. Furthermore, corruption discourages skilled professionals from working in the public healthcare sector, contributing to a decline in overall service quality. Importantly, the study finds that the implementation of anti-corruption measures can lead to significant improvements. Countries and institutions that have adopted digital technologies, such as electronic health records and e-procurement systems, have shown a reduction in corruption-related risks. Increased transparency in budgeting and decision-making processes has also been associated with higher levels of public trust and better service delivery. In addition, the results indicate that strengthening legal frameworks and enforcing anti-corruption laws play a crucial role in reducing corrupt activities. Institutions that maintain strict oversight and accountability mechanisms are more successful in preventing abuse of power. The involvement of civil society and the media in monitoring healthcare systems further contributes to increased transparency and public awareness. Finally, the study demonstrates that improving working conditions and salaries for healthcare professionals can reduce incentives for engaging in corrupt practices. When healthcare workers are fairly compensated and operate in a supportive environment, they are more likely to adhere to ethical standards and professional responsibilities.

DISCUSSIONS

The findings of this study provide a deeper understanding of the complex nature of corruption in the healthcare system and highlight the urgent need for comprehensive and sustainable solutions. The results confirm that corruption is not an isolated issue but rather a systemic

problem that is influenced by a combination of economic, institutional, and social factors. Therefore, addressing corruption requires not only legal reforms but also structural and cultural changes within the healthcare sector. One of the key points emerging from the discussion is that corruption in healthcare is often normalized in certain environments, especially where informal payments are widely accepted as part of the system. This normalization creates a cycle in which both patients and healthcare providers perceive corrupt practices as unavoidable. Patients may feel compelled to offer bribes to receive timely and quality care, while healthcare workers may rely on such payments due to insufficient salaries or lack of incentives. Breaking this cycle requires a shift in both public attitudes and institutional practices. The study also emphasizes the role of governance and institutional capacity in either enabling or preventing corruption. Weak administrative systems, lack of transparency, and ineffective monitoring mechanisms create opportunities for abuse of power. In contrast, countries with strong governance structures and clear accountability systems tend to experience lower levels of corruption. This suggests that reforms should focus on strengthening institutions, improving management practices, and ensuring that responsibilities are clearly defined and enforced. Another important aspect discussed is the impact of digitalization on reducing corruption risks. The introduction of electronic health systems, digital record-keeping, and online procurement platforms has proven to be an effective tool in minimizing human interference and increasing transparency. However, the discussion also notes that digitalization alone is not sufficient. Without proper implementation, training, and oversight, digital systems can still be manipulated or underutilized. Therefore, technological solutions must be accompanied by strong regulatory frameworks and capacity-building efforts. The role of healthcare professionals is also critically examined in this discussion. Ethical behavior and professional integrity are essential in preventing corruption. While external controls and legal measures are important, internal motivation and adherence to ethical standards play an equally significant role. Education and continuous professional development can help foster a culture of honesty and responsibility among healthcare workers. At the same time, improving working conditions and providing fair compensation are necessary to reduce the economic pressures that may lead to corrupt behavior. Furthermore, the discussion highlights the importance of public participation and civil society engagement. Transparency initiatives, access to information, and open communication channels empower citizens to hold healthcare institutions accountable. When patients are aware of their rights and have mechanisms to report corruption, the likelihood of misconduct decreases. Media and non-governmental organizations can also play a key role in exposing corruption cases and promoting accountability. Despite the progress made in some regions, the study acknowledges that challenges remain. Resistance to change, lack of political will, and deeply rooted institutional habits can hinder anti-corruption efforts. In addition, differences in economic development and administrative capacity mean that solutions must be adapted to specific national and local contexts. A one-size-fits-all approach is unlikely to be effective.

CONCLUSION

In conclusion, corruption in the healthcare system represents a serious and persistent challenge that affects not only the quality and accessibility of medical services but also the overall well-being of society. The findings of this study clearly demonstrate that corruption undermines the fundamental principles of fairness, equality, and efficiency in healthcare delivery. It leads to the misuse of limited resources, reduces trust in public institutions, and creates significant barriers for vulnerable populations in accessing necessary medical care. The research confirms that corruption in healthcare is a complex and multi-dimensional issue, driven by various factors such as weak governance, lack of transparency, insufficient accountability, and economic pressures on healthcare workers. These factors interact with one another, creating an environment in which corrupt practices can thrive and become normalized. As a result,

addressing corruption requires a comprehensive and integrated approach that targets its root causes rather than just its symptoms. One of the key conclusions of this study is that strengthening institutional frameworks and legal systems is essential for effectively combating corruption. Clear regulations, strict enforcement of laws, and independent oversight mechanisms can significantly reduce opportunities for abuse of power. At the same time, transparency in financial management and decision-making processes plays a crucial role in ensuring accountability and building public trust.

REFERENCE

1. World Health Organization (WHO). *Corruption in the Health Sector: Problems and Solutions*. Geneva: WHO Press, 2019.
2. OECD (Organisation for Economic Co-operation and Development). *Integrity in the Health Sector: Preventing Corruption*. Paris: OECD Publishing, 2017.
3. World Bank. *Enhancing Governance in Health Systems*. Washington, DC: World Bank Publications, 2020.
4. U4 Anti-Corruption Resource Centre. *Corruption in the Health Sector*. Bergen, Norway, 2019
5. European Commission. *Study on Corruption in the Healthcare Sector*. Brussels, 2015.
6. United Nations Development Programme (UNDP). *Fighting Corruption in the Health Sector: Methods, Tools and Good Practices*. New York: UNDP, 2011.
7. Vian, T. *Review of Corruption in the Health Sector: Theory, Methods and Interventions*. Health Policy and Planning Journal, 2008.
8. Lewis, M. *Informal Payments and the Financing of Health Care in Developing and Transition Countries*. Health Affairs, 2007.
9. Savedoff, W. & Hussmann, K. *Why Are Health Systems Prone to Corruption?* Global Corruption Report, Transparency International, 2010.

PREVALENCE OF ARTERIAL HYPERTENSION AND THE DYNAMICS OF ITS 31-YEAR CHANGES IN THE INDIGENOUS AND MIGRANT RURAL POPULATION OF THE ANDIJAN REGION

Qalandarov Dilmurod Madaminovich
Associate Professor, Candidate of Medical Sciences (PhD), Department of Internal Diseases, Cardiology and Emergency Medicine, Faculty of Advanced Training and Retraining of Physicians, Andijan State Medical Institute, Uzbekistan.

Kenjayev Olimjon Obidjonovich
Head of the Department of Internal Diseases in Family Medicine, PhD, Central Asian Medical

Ravshanov Abdullajon Shukhrat ogli
Central Asian Medical University, Department of Internal Diseases in Family Medicine, 1st-year Clinical Residency Student, Address: 64 Burhoniddin Marg'inani Street, Fergana city, Uzbekistan.

Mamasaliev Nematjon Solievich
Head of the Department of Internal Diseases, Cardiology and Emergency Medicine, Faculty of Advanced Training and Retraining of Physicians, Andijan State Medical Institute, Doctor of Medical Sciences (DSc), Professor.

Usmonov Burhonjon Umarovich
Associate Professor, Department of Internal Diseases, Cardiology and Emergency Medicine, Faculty of Advanced Training and Retraining of Physicians, Andijan State Medical Institute, Uzbekistan.

Salokhidinov Adiljon
Professor, Doctor of Medical Sciences (DSc), Department of Propaedeutics of Internal Diseases, Andijan State Medical Institute.

ABSTRACT
This study is devoted to investigating the increasing epidemiological significance of cardiovascular diseases, particularly arterial hypertension (AH), in recent decades. Since the end of the 20th century, an increase in mortality rates associated with these diseases has been documented in scientific sources as being closely related to socio-economic and psychosocial factors. The aim of the study is to determine the regional characteristics of arterial hypertension in Uzbekistan based on 31 years of screening data, as well as to improve its innovative prevention and control system. The study was conducted in the Andijan region during 1989–2020 and included 3,001 individuals aged 18–89 years (1,421 men and 1,580 women). Epidemiological, clinical, instrumental (ECG, echocardiography, ultrasound), biochemical, and statistical analysis methods were used. The prevalence of stages and degrees of arterial hypertension was determined, and their relative risk indicators were assessed. According to the results, in the indigenous rural population, stage I AH accounted for 61.1%, stage II AH for 31.0%, and stage III AH for 7.9%. By degrees, AH1 was 50.6%, AH2 – 31.0%, and AH3 – 18.4%. Stage I AH was observed 8 times more frequently than stage III. The obtained results demonstrate the importance of regional epidemiological monitoring and justify the need to improve screening programs.
Keywords: arterial hypertension, cardiovascular diseases, epidemiology, screening, prevention, risk factors, population.

INTRODUCTION

Since the end of the 20th century, the increasing mortality rate from cardiovascular diseases and its sharp changes have been confirmed by many researchers. Based on the analysis of epidemiological studies, researchers such as G.Ya. Maslennikova, R.G. Oganov (2018), Ye.V. Akimova et al. (2006), M.M. Kayumova et al. (2023), N.V. Pogosova et al. (2018), and A.M. Akimov (2023) have convincingly demonstrated that such trends have been observed in many countries and are associated with social, economic, and psychosocial factors, most often recorded in comparatively economically active age groups [8; pp. 4–9, 9; pp. 6–9, 6; pp. 33–34, 7; pp. 49–52, 10; pp. 47–56, 1; pp. 104–110].

A.M. Akimov (2023) demonstrated that, in the Arctic region, the type of labor occupies a special place among risk factors and plays an important role in public health. In this region, high labor intensity, relatively long working hours, limited rest days, and minimal social and living conditions were identified by the author as risk factors [1; pp. 104–110].

Therefore, other researchers also recommend conducting and/or continuing epidemiological studies to develop comprehensive programs for the prevention of cardiovascular diseases, taking into account the dynamics of labor characteristics and expeditionary work conditions in specific geographical and ecological regions, such as the Arctic [5; pp. 19–24, 2; pp. 239–240, 3; pp. 34–35, 4; pp. 96–99, 12; pp. 272–280, 13; pp. 1547–1557, 15; pp. 1429–1440]. We believe that these viewpoints contain scientific logic and represent a promising research direction. This is because, over the years, the prevalence of certain diseases increases while others decrease, and their timely and accurate analysis—primarily through epidemiological studies—is an extremely important scientific and practical field. The reason is that the "new foundations" of medicine are built precisely on such data and conclusions.

Until the beginning of the new century, for example, scientific sources suggested that northern populations, particularly indigenous populations, were not affected by diabetes mellitus—this notion had been widely accepted. According to data provided by the World Health Organization (WHO), international migration has shown a steady growth trend in recent years [14].

Thus, in this already formed migrant population (MP), the development and improvement of screening and preventive programs have become highly relevant, and this issue will become even more significant in the future. Existing studies lead to or encourage such conclusions.

According to data published by the Federal State Statistics Service of Russia (2025) and A.S. Andreeva, I.S. Ivanova, Ye.A. Varshaver (2024), the migration flow in the Russian Federation averages about 0.5 million people annually, with most migrants coming from Commonwealth countries [11; 8; pp. 54–72].

OBJECTIVE OF THE STUDY

The objective of the study is to develop regional scientific foundations for the innovative prevention of hypertensive conditions based on 31-year screening data in Uzbekistan, taking into account scientific characteristics, and to implement improved treatment and control measures using new technologies in practice.

MATERIALS AND METHODS

The study population consisted of 3,001 individuals (1,421 men and 1,580 women) aged 18–89 years, selected from an unorganized population in the Andijan region using a 10% random sampling method based on a random number table. These individuals were involved in arterial hypertension monitoring from 1989 to 2020. As the subject of the study, venous blood

and serum samples of patients were collected for biochemical analysis. International criteria were used for diagnosing arterial hypertension (AH) and hypertension (HT), as well as for analyzing risk factors of comorbid diseases. Comprehensive statistical modeling indicators serving as a basis for regional preventive strategies were obtained.

Methods of the study: Epidemiological, general clinical, instrumental (cardiac echocardiography, ECG, ultrasound, anthropometric measurements, tonometry), biochemical, pharmacoepidemiological, pharmacoeconomic, pharmacovigilance, and statistical methods were applied.

RESULTS

In the indigenous rural population, based on I and II screening among individuals aged 18–89 years, the stages and degrees of arterial hypertension were characterized by the following prevalence rates (as shown in Table 1 and Figure 1):

AH stage I – 61.1% AH1 – 50.6%
AH stage II – 31.0% AH2 – 31.0%
AH stage III – 7.9% AH3 – 18.4%

A higher frequency was observed for AH stage I (8 times more frequent compared to AH stage III). AH1 was recorded 2.7 times more frequently than AH2 [RR = 1.4; CI = 1.82–1.08; χ^2 = 16.9].

Скрининг гурлари, абориген аҳоли сони	**АГ боскичлари ва даражадари**														**RR**	**ИИ**	**X²**
	АГ I		**АГ II**		**АГ III**		**Жами АГ**		**АГ₁**		**АГ₂**		**АГ₃**				
	n	**%**	**n**	**%**	**n**	**%**	**n**	**%**	**n**	**%**	**n**	**%**	**n**	**%**			
I ск (n=1754)	829	60,7	424	31,0	113	8,3	1366	66,3	163	50,8	91	28,3	67	20,9	**0,98**	**1,38-0,7**	**0,026**
P	>0,05		>0,05		>0,05		**<0,05**		>0,05		**<0,05**		**<0,05**				
II ск (n=956)	429	61,9	214	30,9	50	7,2	693	33,7	85	50,3	61	36,1	23	13,6	**1,01**	**1,32-0,77**	**0,011**
УмП (n=2710)	1258	61,1	638	31,0	163	7,9	2059	100	248	50,6	152	31,0	90	18,4	**1,4**	**1,82-1,08**	**16,9**

^{2}=16,9].



characterized as follows, respectively: AH I – 60.7% and 61.9% (with a decreasing trend of 1.7%; P<0.05), AH II – 31.0% and 30.9% (with a decreasing trend of 0.1%), AH III – 8.3% and 7.2% (showing a decreasing trend of 1.1%; P>0.05), total AH – 66.3% and 33.7% (with a decreasing trend of 32.6%; P<0.05), AH1 – 50.8% and 50.3% (with a decrease of 0.5%; P>0.05), AH3 – 20.9% and 13.6% (with a decrease of 6.3%; P<0.05).

Скрининг гурлари, ўтирок аҳоли сони	АГ боскичлари ва даражадари														RR	ИИ	X^2
	АГ I		АГ II		АГ III		Жами АГ		$АГ_1$		$АГ_2$		$АГ_3$				
	n	%	n	%	n	%	n	%	n	%	n	%	n	%			
I ск (n=190)	92	62,6	50	34,0	5	3,4	147	56,3	19	65,5	9	31,0	1	3,4	**1,18**	**1,36-1,02**	**36,65**
P	**<0,05**		**<0,05**		**>0,05**		**<0,05**		**>0,05**		**>0,05**		**<0,05**				
II ск (n=101)	48	42,1	58	50,9	8	7,0	114	43,7	8	42,1	5	26,3	6	31,6	**1,31**	**1,44-1,19**	**49,37**
УмП (n=291)	140	53,6	108	41,4	13	5,0	261	100,0	27	56,3	14	29,2	7	14,6	**1,26**	**1,37-1,16**	**82,07**

In this population, the prevalence rates of arterial hypertension (AH) in 1989 (I screening) and 2020 (II screening), as well as their 31-year dynamic changes, are characterized as follows, respectively: AH I – 60.7% and 61.9% (with a decreasing trend of 1.7%; P<0.05), AH II – 31.0% and 30.9% (with a decreasing trend of 0.1%), AH III – 8.3% and 7.2% (showing a decreasing trend of 1.1%; P>0.05), total AH – 66.3% and 33.7% (with a decreasing trend of

32.6%; P<0.05), AH1 – 50.8% and 50.3% (with a decrease of 0.5%; P>0.05), AH3 – 20.9% and 13.6% (with a decrease of 6.3%; P<0.05).
According to the I screening, the prevalence of AH I is more than 7 times higher compared to AH III, while the difference between AH1 and AH3 is confirmed to be 1.8 times [RR=0.98; CI=1.38–0.7; χ^2=0.026]. The analysis of the II screening shows that these differences in the indigenous rural population are 8.7 and 3.8 times, respectively [RR=1.4; CI=1.82–1.08; χ^2=16.9].
The subsequent analysis of our prospective epidemiological study is devoted to evaluating the dynamics of arterial hypertension in 1989–2020 among the migrant rural population of the Andijan region (presented in Table 2 and Figure 2).



dynamics of arterial hypertension in 1989–2020 among the migrant rural population of the Andijan region (presented in Table 2 and Figure 2).
In this migrant population, the detection frequency of AH in the I and II screenings was 56.3% and 43.7%, respectively, characterizing a decreasing trend of 12.6% over 31 years (P<0.05). The highest prevalence rates were observed for AH I and AH1 – 53.6% and 56.3% (P<0.05). Compared to AH III and AH3 stages and degrees, a significant difference (59.2% and 63.1%, respectively) in favor of AH I and AH1 indicators was confirmed in the I screening [RR=1.18; CI=1.36–1.02; χ^2=36.65].
According to the II screening data, in this population, the prevalence of AH I exceeds that of AH III by 6 times, and the detection rate of AH1 exceeds that of AH2 by 3 times [RR=1.31; CI=1.44–1.19; χ^2=49.37].
AH I – 62.6% and 42.1% (with a decrease of 19.5%; P<0.05), AH II – 14.0% and 50.9% (with an increase of 46.9%; P<0.05), AH III – 3.4% and 7.0% (with an increase of 3.6%; P>0.05), AH1 – 65.5% and 42.1% (with a decreasing trend of 23.4%; P>0.05), AH2 – 31.0% and 26.3% (with a decrease of 4.3%; P>0.05), and AH3 – 3.4% and 31.6% (with an increase of 28.2%; P<0.05) were confirmed in the I and II screenings, respectively.
In the overall population, based on I and II screening of the total migrant population, arterial hypertension demonstrated the following epidemiological characteristics: AH I – 53.6%, AH II – 41.4%, AH III – 5.0%, AH1 – 56.3%, AH2 – 29.2%, and AH3 – 14.6%. Considering the

dynamics of these characteristics is of diagnostic, preventive, and pharmacoepidemiological importance in the elimination and control of hypertension.

CONCLUSION

In the representative male and female population of the Andijan region, the 31-year changes in the origin and prevalence of arterial hypertension stages (AH I, II, III) and degrees (AH1, 2, 3) confirm regional characteristics: Ages 18–22: 57.1% and 25.0% (with decrease), 42.9% and 75.0% (with increase), 0.0% and 0.0% (with stability), and – 10.0% and 0.0% (with stability), 100.0% and 0.0% (with stability), 0.0% and 0.0%; Ages 23–44: 74.7% and 83.9% (with decrease and increase), 20.2% and 10.9% (with increase and decrease), 5.1% and 5.2% (with increase), and – 76.2% and 91.7% (with decrease and increase), 4.8% and 0.0% (with decrease), 19.0% and 8.3% (with increase and decrease); Ages 45–59: 49.6% and 64.6% (with decrease), 39.2% and 27.8% (with increase), 11.2% and 7.7% (with decrease), and – 43.9% and 61.8% (with increase and decrease), 25.6% and 8.8% (with increase), 30.5% and 29.4% (with decrease); Ages 60–74: 55.8% and 60.0% (with increase and decrease), 34.6% and 31.5% (with decrease and increase), 9.6% and 8.5% (with decrease and increase), and – 49.4% and 53.6% (with increase and decrease), 33.5% and 29.6% (with decrease and increase), 17.0% and 16.8% (with decrease and increase). Ages 75–89: 65.2% and 55.4% (with increase), 33.3% and 39.1% (with increase and decrease), 1.4% and 5.4% (with increase), and – 27.1% and 14.3% (with decrease and increase), 1.4% and 5.4% (with increase), 32.9% and 20.0% (with decrease and stability), 1.4% and 4.3% (with decrease and increase).

REFERENCES

1. Акимов А.М. Некоторые факторы хронического социального стресса у мужчин, занятых экспедиционно-вахтовой формой труда на арктических территориях Тюменского региона (пилотное исследование). Сибирский научный медицинский журнал. 2023;43(3):104-112.
2. Акимов А.М. Физическая активность и характер труда в популяции мужчин трудоспособного возраста. Омский научный вестник. 2015; 2(136):238-240.
3. Акимов А.М., Гакова Е.И., Каюмова М.М. и др. Табакокурение и его ассоциации с отношением к здоровью мужчин 25—54 лет, работающих экспедиционно-вахтовым методом в Арктическом регионе Западной Сибири. Профилактическая медицина. 2024;27(4):32-37.
4. Акимов А.М., Каюмова М.М., Гакова Е.И. и др. Отношение к своему здоровью в открытой популяции среднеурбанизированного города Западной Сибири, ассоциации с распространенностью ИБС: гендерные особенности. Сибирский научный медицинский журнал. 2021;41(4):95-102.
5. Акимова Е.В., Гафаров В.В., Каюмова А.Р. Физическая активность и отношение к профилактике среди мужчин, занятых мобильным трудом в Арктической зоне России. Международный журнал сердца и сосудистых заболеваний. 2024;12(43):19-26.
6. Акимова Е.В., Кузнецов В.А., Гафаров В.В. Риск сердечно-сосудистой смерти в различных социальных группах открытой популяции Тюмени. Профилактика заболеваний и укрепление здоровья. 2006;9(4):33-37.
7. Каюмова М.М., Акимов А.М., Бессонова М.И. и др. Ассоциации стресса в семье, параметров отношения к здоровью и его самооценка у работающих экспедиционно-вахтовым методом в Арктике. Профилактическая медицина. 2023;26(10):49-54.
8. Масленникова Г.Я., Оганов Р.Г. Выбор оптимальных подходов к профилактике неинфекционных заболеваний в рамках международного сотрудничества. Кардиоваскулярная терапия и профилактика. 2018; 17(1):4-9.

9. Оганов Р.Г., Масленникова Г.Я. Демографические тенденции в Российской Федерации: вклад болезней системы кровообращения. Международный журнал сердца и сосудистых заболеваний. 2013;1(1):4-9.
10. Погосова Н.В., Бойцов С.А., Оганов Р.Г. и др. Клинико-эпидемиологическая программа изучения психосоциальных факторов риска в кардиологической практике у больных артериальной гипертонией и ишемической болезнью сердца (КОМЕТА): первые результаты российского многоцентрового исследования. Кардиология. 2018;58(9):47-58.
11. Федеральная служба государственной статистики. Информационноаналитические материалы. Численность и миграция населения Российской Федерации. Ссылка активна на 06.02.25.
12. Curry SJ, Krist AH, Owens DK, et al. US Preventive Services Task Force. Risk Assessment for Cardiovascular Disease with Nontraditional Risk Factors: US Preventive Services Task Force Recommendation Statement. JAMA. 2018;320(3):272-280. https://doi.org/10.1001/jama.2018.8359.
13. Tunstall-Pedoe H, Kuulasmaa K, Mahonen M, et al. Contribution of trends in survival and coronary-event rates to changes in coronary heart disease mortality: 10-year results from 37 WHO MONICA project populations. The Lancet. 1999;353:1547-57.
14. World report on the health of refugees and migrants: summary. Geneva: World Health Organization; 2022. Accessed November 27,2024. https://iris.who.int/bitstream/handle/10665/360466/9789240054868-rus.pdf.
15. Yao B-C, Meng L-B, Hao M-L, et al. Chronic stress: a critical risk factor for atherosclerosis. Journal of International Medical Research. 2019;47(4): 1429-1440. https://doi.org/10.1177/0300060519826820.

FROM TRANSMISSION TO PERSONALISATION: A GAMIFIED INTELLIGENT TUTORING FRAMEWORK FOR ENGINEERING COMPETENCY DEVELOPMENT

DSC. Prof, Xakimhon Xanbabayev[1]
[1] Department of "Digital technology and artificial intelligence", Kokand State University, Fergana, Uzbekistan

Assoc. Prof., Akhrorjon ADBUAMANONOV[2]
[2] Department of "Modeling of Medical Hygiene Processes", Central Asian Medical University, Fergana, Uzbekistan
[2] ORCID ID: https://orcid.org/ 0000-0002-5284-2985

ABSTRACT

This paper presents the design, implementation, and empirical evaluation of an Adaptive Learning Model (ALM) grounded in Intelligent Control Systems (ICS) for engineering disciplines. Implemented in the CAMU-Edu platform and tested with 126 students over two academic years (2024–2025), the model produced statistically significant improvements across all measured competency dimensions: knowledge acquisition (+19%), analytical skills (+28%), problem-solving (+23%), motivation (+21%), and digital competency (+28%). Effect sizes ranged from d = 0.98 to d = 1.62, substantially exceeding the meta-analytic benchmark of d = 0.76. The paper formalises the adaptive architecture mathematically and compares the platform against Moodle, Coursera, and edX.

Keywords: intelligent control systems; adaptive learning; engineering education; ITS; gamification; learning analytics; competency-based assessment.

INTRODUCTION

Engineering higher education faces a structural mismatch: graduates must master cyber-physical systems and data-driven environments, yet most institutions continue to rely on transmission-based pedagogical models poorly suited to developing adaptive, analytical, and technological competencies [1, 2].

Intelligent Control Systems (ICS) — encompassing AI, machine learning, and automated management — have demonstrated transformative potential across industrial sectors; their application to educational environments represents a logically consistent extension, mirroring how ICS optimise industrial processes through real-time data acquisition and adaptive intervention [3–5].

Four critical deficiencies in traditional engineering instruction have been identified in the literature: (i) uniform content delivery ignoring heterogeneous cognitive profiles; (ii) periodic summative assessment insufficient for timely pedagogical intervention; (iii) limited human-system interaction reducing immediate feedback; and (iv) absence of motivational scaffolding leading to diminishing engagement over time [6–10].

Existing intelligent tutoring systems (Carnegie Learning, ALEKS, Knewton) demonstrate measurable learning improvements but predominantly target mathematics at pre-university levels; few are specifically designed for engineering higher education in emerging economies, and none fully integrate adaptive delivery, real-time analytics, automated assessment, and gamification within a unified framework [11–13].

This paper addresses these gaps through four research objectives: (i) formalise a multi-component adaptive learning model grounded in control theory; (ii) implement it in a functional online environment; (iii) evaluate effectiveness against traditional instruction using multiple competency indicators; and (iv) compare the platform against established LMS solutions.

Intelligent Tutoring Systems (ITS) trace their origins to Carbonell [14] and Anderson et al. [15]; contemporary architectures comprise four canonical modules — domain model, learner model, pedagogical model, and interface model — each substantially enhanced by recent machine learning advances [16].

Deep Knowledge Tracing (DKT), introduced by Piech et al. [17], employs recurrent neural networks to model temporal learning dynamics with prediction accuracies exceeding 86%; reinforcement learning approaches further enable ITS to optimise pedagogical strategies through interaction experience rather than pre-specified rules [18].

A meta-analysis by Steenbergen-Hu and Cooper [19] across 26 studies found ITS produced an average effect size of $d = 0.76$ relative to traditional instruction; more recent analyses confirm positive effects across STEM domains ranging from $d = 0.40$ to $d = 1.20$ depending on implementation fidelity [20].

Engineering education requires simultaneous development of theoretical understanding, procedural competency, and higher-order capabilities; virtual laboratories and simulation environments have proved effective — Chen and Li [22] reported +31% procedural accuracy and +24% transfer performance using PLC simulation compared to physical equipment alone.

Gamification, supported by Self-Determination Theory (SDT), consistently improves engagement when elements support competence (level progression), autonomy (choice of learning pathway), and relatedness (collaborative challenges); learning analytics closes the feedback loop by continuously refining personalisation through a data-feedback architecture mirroring ICS closed-loop control [10, 25, 26]

METHODOLOGY

The ALM adopts a closed-loop feedback architecture grounded in cybernetic control theory (Wiener, 1948), comprising five modules: (1) Data Acquisition — records interaction events every 10 seconds; (2) Intelligent Analysis — applies k-NN and gradient-boosted trees; (3) Recommendation Engine — selects content based on competency profiles; (4) Monitoring

Dashboard — real-time instructor visualisation; (5) Feedback Mechanism — updates learner model and triggers content reassignment.

Teaching effectiveness is formalised as $S = 0.4K + 0.2M + 0.25A + 0.15I$, where K = knowledge acquisition, M = motivational engagement, A = adaptive algorithm effectiveness, I = intelligent analytics output; weights were validated via analytic hierarchy process (AHP, n = 12 experts, CR = 0.07 < 0.10). Student performance is tracked by $OI = 0.4P + 0.4T + 0.2A$.

CAMU-Edu was built on a three-tier architecture: React.js/Tailwind CSS frontend with role-specific interfaces (student, instructor, administrator); Django REST Framework backend with JWT authentication and WebSocket real-time communication; MySQL relational database. The gamification subsystem uses an XP model with logarithmic level progression (base 1.5) calibrated to yield ~10 levels per 16-week semester.

A pre-test/post-test quasi-experimental design was employed at Namangan State Technical University. Engineering students (N = 126; 78 undergraduate, 48 graduate) were assigned to experimental (EG, n = 64, CAMU-Edu with full ALM) and control groups (CG, n = 62, conventional instruction); both groups covered identical content with the same instructors. Baseline equivalence was confirmed: $t(124) = 0.83$, $p = 0.41$, $d = 0.15$.

The study spanned two academic years (2024–2025) across four courses — Fundamentals of Automation, Digital Control Systems, IoT and Sensor Networks, and Industrial Process Modelling. Six competency dimensions were assessed using standardised tests, PLC/SCADA simulation tasks, the validated MSLQ motivation scale, and portfolio-based evaluation; statistical analyses employed independent-samples t-test, one-way ANOVA, and Cohen's d ($\alpha = 0.05$).

RESULTS AND DISCUSSION

Post-test comparisons revealed statistically significant improvements across all six competency dimensions (all $p < 0.001$): knowledge acquisition 68% → 87% (+19 pp); independent analytical skills 54% → 82% (+28 pp); practical problem-solving 61% → 84% (+23 pp); academic motivation 70% → 91% (+21 pp); digital competency 58% → 86% (+28 pp); automation modelling accuracy 55% → 83% (+28 pp).

Effect sizes were large throughout: d = 1.47 (analytical skills), d = 1.43 (digital competency), d = 1.29 (problem-solving), d = 1.12 (motivation), d = 0.98 (knowledge); overall platform effectiveness SE yielded d = 1.62 — substantially exceeding the meta-analytic STEM benchmark of d = 0.76 [19] and the educationally significant threshold of d > 0.40.

Gamification metrics confirmed sustained engagement: 87% of experimental students rated the IBT environment as increasing course engagement; 73% identified it as developing professionally relevant skills; mean XP accumulation was 243 points/week (SD = 68); badge acquisition rates reached 94% ('First Achievement'), 78% ('Analytical Learner'), and 71% ('Consistent Contributor').

Instructor evaluation (n = 12) was highly positive: all 12 (100%) agreed the platform simplified student progress monitoring; 10/12 (83%) reported earlier identification of at-risk students via AI analytics; 9/12 (75%) estimated a 35–40% reduction in administrative workload, freeing time for higher-order instructional activities.

Comparative analysis showed CAMU-Edu as the only evaluated platform offering full adaptive delivery, deep gamification, and local engineering context adaptability simultaneously; Moodle, Coursera, and edX all lacked either full adaptive integration, native gamification frameworks, or Uzbek-language engineering localisation. The four-stage instructional process yielded additional improvements: digital process automation skills +27 pp, digital integration of learning +31 pp.

CONCLUSION

The ALM, implemented in CAMU-Edu and evaluated over two academic years, is established as a scientifically validated, scalable framework for intelligent engineering education: all six competency dimensions showed statistically significant and practically large improvements, confirming the superiority of ICS-grounded adaptive instruction over traditional methods.

The mathematical formalisation of the adaptive architecture — closed-loop feedback, weighted composite effectiveness indices S and OI, IRT-calibrated assessment, and logarithmic gamification progression — provides both a replicable design blueprint and operational metrics for continuous quality monitoring.

The study's primary limitations are its quasi-experimental design (non-random assignment), single-institution setting in Uzbekistan, one-year follow-up period, and absence of instructor-crossover controls; future research should address these through randomised multi-site designs with longitudinal retention assessment.

Priority directions for future work include: (i) NLP-based automated formative feedback on open-ended design tasks; (ii) reinforcement learning for dynamic optimisation of pedagogical strategy selection; (iii) extension to medical and interdisciplinary STEM contexts; (iv) longitudinal evaluation of competency transfer to workplace performance; (v) formal IEC 61508 SIL safety evaluation for high-stakes examination deployment.

REFERENCES

[1] OECD. (2024). Digital Education Outlook 2024: Learning in the Age of AI. OECD Publishing. https://doi.org/10.1787/c74f03de-en

[2] UNESCO. (2023). Artificial Intelligence and Education: Guidance for Policy-makers. UNESCO, Paris.

[3] Russell, S., & Norvig, P. (2022). Artificial Intelligence: A Modern Approach (4th ed.). Pearson.

[4] Wiener, N. (1948). Cybernetics: Or Control and Communication in the Animal and the Machine. MIT Press.

[5] Baker, R. S. J. D. (2010). Data mining for education. In International Encyclopedia of Education (3rd ed., pp. 112–118). Elsevier.

[6] Vygotsky, L. S. (1978). Mind in Society. Harvard University Press.

[7] Bloom, B. S. (1984). The 2 sigma problem. Educational Researcher, 13(6), 4–16.

[8] Black, P., & Wiliam, D. (1998). Assessment and classroom learning. Assessment in Education, 5(1), 7–74.

[9] Hattie, J., & Timperley, H. (2007). The power of feedback. Review of Educational Research, 77(1), 81–112.

[10] Deci, E. L., & Ryan, R. M. (2000). The 'what' and 'why' of goal pursuits. Psychological Inquiry, 11(4), 227–268.

[11] Koedinger, K. R., & Corbett, A. T. (2006). Cognitive tutors. In Cambridge Handbook of the Learning Sciences (pp. 61–77).

[12] Pane, J. F. et al. (2014). Effectiveness of cognitive tutor Algebra I at scale. Educational Evaluation & Policy Analysis, 36(2), 127–144.

[13] Zawacki-Richter, O. et al. (2019). Systematic review of AI in higher education. Int. J. Educational Technology in HE, 16, 39.

[14] Carbonell, J. R. (1970). AI in CAI. IEEE Transactions on Man-Machine Systems, 11(4), 190–202.

[15] Anderson, J. R. et al. (1995). Cognitive tutors: Lessons learned. Journal of the Learning Sciences, 4(2), 167–207.

[16] Nwana, H. S. (1990). Intelligent tutoring systems: An overview. Artificial Intelligence Review, 4(4), 251–277.

[17] Piech, C. et al. (2015). Deep knowledge tracing. Advances in Neural Information Processing Systems, 28.

[18] Rafferty, A. N. et al. (2016). Faster teaching via POMDP planning. Cognitive Science, 40(6), 1290–1332.

[19] Steenbergen-Hu, S., & Cooper, H. (2014). Meta-analysis of ITS on college students. Journal of Educational Psychology, 106(2), 331–347.

[20] Ma, W. et al. (2014). Intelligent tutoring systems and learning outcomes. Journal of Educational Psychology, 106(4), 901–918.

[21] CDIO Initiative. (2023). Rethinking Engineering Education. MIT Press.

[22] Chen, X., & Li, Y. (2022). Intelligent control systems in engineering education. Computers & Education, 178, 104406.

[23] Nguyen, T. T. et al. (2023). SCADA-based control for industrial steam boiler education. JP Journal of Heat and Mass Transfer, 34, 1–18.

[24] Zhang, W. et al. (2022). IoT-based learning analytics for engineering competency. IEEE Transactions on Education, 65(3), 412–421.

[25] Dicheva, D. et al. (2015). Gamification in education: A systematic mapping study. Educational Technology & Society, 18(3), 75–88.

[26] Ferguson, R. (2012). Learning analytics: Drivers, developments, challenges. Int. J. Technology Enhanced Learning, 4(5/6), 304–317.

[27] Romero, C., & Ventura, S. (2020). Educational data mining and learning analytics. WIREs Data Mining & Knowledge Discovery, 10(3), e1355.

[28] Siemens, G. (2013). Learning analytics: The emergence of a discipline. American Behavioral Scientist, 57(10), 1380–1400.

[29] Anderson, J. (2022). Artificial intelligence in higher education. AI & Education Journal, 34(5), 121–134.

[30] Ghimire, P., & Pradhan, P. (2022). PLC-based automation in educational contexts. Automation and Control Engineering, 12(7), 78–85.

[31] Csikszentmihalyi, M. (1990). Flow: The Psychology of Optimal Experience. Harper & Row.

[32] Bonk, C. J., & Graham, C. R. (2012). The Handbook of Blended Learning. Wiley.

[33] World Economic Forum. (2023). Future of Jobs Report 2023. WEF, Geneva.

[34] Abdumanonov A.A. (2025). Texnika fanlarini o'qitishda IBT. Ta'lim va taraqqiyot, 6, 456–462.

[35] Abdumanonov A.A. (2025). Improvement of the methodology of using ICS. NUUz, 1(1.11.1), 55–57.

ANTIBIOTIC RESISTANCE AND UNUSUAL PHENOTYPIC CHARACTERISTICS OF LOCAL STRAINS OF PATHOGENS CAUSING ACUTE INTESTINAL INFECTIONS (REPUBLIC OF UZBEKISTAN)

Sayidmirzayeva Nazokat Gayratjonovna [1]
Assistant of Central Asian Medical University, Fergana, Uzbekistan
[1]ORCID ID: https://orcid.org/0009-0006-3801-5891

Shadmanova Hargiza Abidovna[2]
[2]Center for Retraining and Advanced Training of Personnel in the Field of Sanitary and Epidemiological Well-Being and Public Health
[2]ORCID ID: https://orcid.org/0009-0005-2610-4021

Abstract
The growing prevalence of antimicrobial resistance among pathogens causing acute intestinal infections represents a significant public health concern worldwide. In the Republic of Uzbekistan, limited data are available on the resistance patterns and phenotypic variability of locally circulating strains. The emergence of unusual phenotypic characteristics may complicate routine laboratory identification and affect the accuracy of diagnostics. This, in turn, can lead to inappropriate antimicrobial therapy and contribute to the further spread of resistant strains. Investigating local epidemiological trends is essential for optimizing diagnostic strategies and treatment guidelines. Therefore, studying antibiotic resistance and atypical phenotypes of regional pathogens is of critical importance for strengthening infectious disease control and surveillance systems.

Aim of research:
To assess the prevalence of major bacterial pathogens causing acute intestinal infections among patients in the Fergana region, Republic of Uzbekistan.

Methods *of research*:
A retrospective analysis of microbiological data from 2015 to 2022 was conducted at the Fergana branch of the Republican Specialized Scientific-Practical Medical Center for Epidemiology, Microbiology, Infectious and Parasitic Diseases. Isolation, phenotypic identification, and antimicrobial susceptibility testing of bacterial strains were performed in accordance with national guidelines (Ministry of Health of Uzbekistan, Order No. 177, 2015) and EUCAST standards.

Results

A retrospective analysis (2015–2022) of microbiological examinations of biological samples obtained from patients diagnosed with acute intestinal infections was carried out to identify etiological agents in the bacteriological laboratory of the Fergana branch of the above-mentioned center. During this period, the biological characteristics and primary identification of isolated microorganisms were performed by laboratory specialists.

A total of 9,792 bacterial isolates were obtained from stool samples collected within routine clinical practice between January 2015 and December 2022. Of these, 9,365 (95.6%) were Gram-negative rods and 427 (4.4%) were Gram-positive cocci. The retrospective analysis demonstrated that pathogenic representatives of intestinal microorganisms were significantly less frequent compared to opportunistic enterobacteria, with 502 isolates belonging to the genera *Salmonella* and *Shigella*. Among Enterobacteriaceae, the most commonly identified organisms during 2015–2021 were *Citrobacter spp.*, *Enterobacter spp.*, *Proteus mirabilis*, and *Klebsiella pneumoniae*, with a noticeable increase in *K. pneumoniae* observed in 2019 and 2021.

Among *Shigella* isolates, *Shigella flexneri* (n=61) predominated, confirming previous epidemiological data for the region. However, among *Salmonella*, only *Salmonella typhimurium* (n=441) was identified during the study period, suggesting the need for a systematic reassessment of microbiological diagnostic approaches for this genus. It should be noted that routine laboratory diagnostics primarily focus on officially recognized pathogenic enterobacteria (*Shigella spp.*, *Salmonella spp.*, *Escherichia coli*), which may result in incomplete etiological analysis of acute intestinal infections. This limitation is associated with outdated regulatory frameworks, insufficient diagnostic resources, and limited implementation of modern diagnostic methods.

At the same time, for pathogens such as *Escherichia coli* and *Staphylococcus aureus*, it is essential to determine the presence of toxin-producing genes and specific virulence factors responsible for pathogenicity, as their identification is directly associated with establishing the etiological role of the isolated strains in acute intestinal infections. However, such investigations are not routinely performed in practice. Existing regulatory documents and commonly applied diagnostic methods do not fully cover all necessary studies required for comprehensive etiological interpretation of acute intestinal infections.

Since early 2021, the epidemiological situation related to coronavirus infection in the region, along with the modernization of laboratory infrastructure, has enabled the implementation of automated bacteriological analyzers such as the VITEK 2 Compact in microbiological laboratories. This has contributed to the generation of new data regarding the etiology of acute intestinal infections. In particular, microorganisms such as *Pseudomonas aeruginosa* (41%), *Pseudomonas putida* (31%), and members of the *Burkholderia cepacia* complex (27.4%) were identified.

While the use of automated analyzers facilitates the identification of etiological agents, the high titers of such microorganisms isolated from clinical specimens require further detailed investigation to clarify their clinical significance. The epidemiological situation observed in the region during 2020–2022, along with the increased use of antibacterial agents, may have contributed to the emergence and spread of these organisms, indicating the need for more comprehensive study of this evolving local issue.

As a continuation of the research, the serotypes of 92 Salmonella strains sent to the Bacteriology Laboratory of the Committee for Sanitary Epidemiology and Public Health of the Republic of Uzbekistan in 2023 for confirmation were studied using a complete biochemical analysis and serological sera. During the study, 85 were confirmed as Salmonella spp., representing a total of 12 serovars. Of these, 34 strains (40%) belonged to the S. Enteritidis subtype, proving that this type is widespread among salmonellae in Uzbekistan, as is the case in international practice. (Figure 1)



Figure 1. Serotyping Results of *Salmonella* spp. Isolated from Acute Intestinal Infections in the Republic of Uzbekistan, 2023.

As a further stage of this study, the antimicrobial susceptibility of local *Salmonella* spp. strains was assessed in accordance with EUCAST guidelines (version 14.0; 2024). The results demonstrated that the majority of isolates were highly susceptible to cephalosporins. However, resistance among the studied isolates was identified as follows: 3 out of 10 isolates to ceftazidime, 6 isolates to cefotaxime, and 4 isolates to ceftriaxone. Microbiological analysis of 85 *Salmonella* isolates revealed an intermediate susceptibility rate to ceftazidime of 42.3%, while the highest resistance among cephalosporins was observed for this agent (1.2%; 10 isolates). Notably, ceftriaxone, a third-generation cephalosporin, remains widely used for the treatment of infections caused by Gram-negative bacteria, including *Salmonella*, whereas cefepime, a fourth-generation cephalosporin, is effective against Enterobacteriaceae producing AmpC β-lactamases. Laboratory findings also indicated the presence of extended-spectrum β-lactamase (ESBL)-producing and AmpC cephalosporinase-producing resistant *Salmonella* variants among the isolates.

In the subsequent phase, 85 *Salmonella* isolates were tested for susceptibility to five fluoroquinolones. In contrast to cephalosporins, resistance to fluoroquinolones was relatively high, with nearly half of the isolates demonstrating resistance to agents such as norfloxacin, ofloxacin, and moxifloxacin. It should be noted that criteria for interpreting fluoroquinolone susceptibility in *Salmonella* have evolved over time. Prior to 2011, nalidixic acid was commonly used as a screening agent, and resistance to this compound was considered indicative of resistance to all fluoroquinolones. More recent studies recommend a combined phenotypic approach using two indicator disks—pefloxacin and nalidixic acid—for more accurate detection. Pefloxacin, in particular, enables the identification of both chromosomal and plasmid-mediated resistance mechanisms.

Globally, members of the Enterobacteriaceae family, including *Salmonella enterica*, as well as *Escherichia coli*, *Klebsiella pneumoniae*, and *Proteus mirabilis*, are recognized as major producers and carriers of AmpC genes. These resistance determinants are known to spread via mobile genetic elements such as plasmids, transposons, and integrons, facilitating horizontal gene transfer between microorganisms. Loss of susceptibility to β-lactam antibiotics is predominantly associated with such mechanisms. The widespread distribution of these genes among environmental and clinical Enterobacteriaceae indicates that they serve as significant reservoirs for resistance determinants in both *Salmonella* and opportunistic enterobacteria. This highlights their important role in the emergence and dissemination of antimicrobial resistance in pathogens causing acute intestinal infections. Taking these data into account, eight isolates of opportunistic Enterobacteriaceae identified as etiological agents of acute intestinal infections were further investigated using both phenotypic and genotypic methods (Figure 2).



Figure 2. Phenotypic detection of β-lactam resistance among members of the Enterobacteriaceae family.

The obtained results demonstrated the presence of resistance-associated genes, including *ctx* in *Morganella morganii*, *ctx* and *tem* in *Escherichia coli*, and *shv* in *Klebsiella* spp. Despite the limited number of analyzed strains, preliminary conclusions suggest that the widespread empirical use of clinically significant antimicrobial agents without adequate laboratory guidance may contribute to the emergence and dissemination of β-lactamase-producing genes among opportunistic enterobacteria in the region.

This situation not only explains the previously observed resistance of local *Salmonella* spp. isolates to third- and fourth-generation cephalosporins but also highlights the necessity for comprehensive molecular investigations of all bacterial pathogens causing acute intestinal infections. In particular, it underscores the importance of detecting genes encoding various classes of β-lactamases (including cephalosporinases and carbapenemases), as well as elucidating the genetic mechanisms underlying acquired resistance to fluoroquinolones.

Conclusions: Taking the above into account, modern international approaches to determining antimicrobial resistance in local isolates require the combined application of PCR-based methods alongside classical phenotypic techniques. This integrated approach is essential for monitoring opportunistic Enterobacteriaceae as etiological agents of acute intestinal infections, allowing not only the assessment of resistance patterns but also the evaluation of acquired pathogenic traits. Furthermore, to ensure the reliability of antimicrobial susceptibility monitoring results in Enterobacteriaceae, it is necessary to develop and implement optimal diagnostic algorithms aimed at identifying the molecular mechanisms underlying clinically significant resistance.

Адабиётлар

1. Искандарова, Г. Т. Эпидемиологические аспекты кишечных инфекций в Ташкентской области Республики Узбекистан / Г. Т. Искандарова, О. Н. Шарапов, Д. Ю. Юсупова. — Текст : непосредственный // Молодой ученый. — 2017. — № 1.2 (135.2). — С. 57- 59. — URL: https://moluch.ru/archive/135/37846/
2. Antimicrobial resistance (AMR) reporting protocol 2023 European Antimicrobial Resistance Surveillance Network (EARS-Net) surveillance data for 2022 39 p.
3. Crump J.A., Sjölund-Karlsson M., Gordon M.A., Parry C.M. Epidemiology, clinical presentation, laboratory diagnosis, antimicrobial resistance, and antimicrobial management of invasive *Salmonella* infections. *Clinical Microbiology Reviews.* 2015; 28(4): 901-37.
4. CDC. National Antimicrobial Resistance Monitoring System (NARMS) Now: Human Data. Atlanta, Georgia: U.S. Department of Health and Human Services, CDC. Available at: https:// www.cdc. gov/narmsnow.
5. EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Prevention and Control), 2019. The European Union summary report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2018. EFSA Journal. 2019; 17(2): 5598.; CDC.
6. Shadmanova N.A., Iskhakova K.I., Rasulmuhamedova M., *Antibiotic resistance of hospital strains of Enterobacteriaceae and phenotypic methods for detecting beta-lactamases*, European Science Review, 2016.
7. Shadmanova N.A., Sayidmirzaeva N.G., Qodirova R.R., *Epidemiological monitoring and etiological diagnosis of major bacterial pathogens of acute intestinal infections in Uzbekistan*, Republican Scientific-Practical Conference Materials, Tashkent, May 2024, pp.152–155.
8. World Health Organization. Informe Estimación de la carga mundial de las enfermedades de transmisión alimentaria. Ginebra [Internet]. 2015 [Updated: 2017]. Available from: http://www. who.int/mediacentre/news/releases/2015/foodborne-disease-estimates/es/ [Accessed: February 2017].

THE PREVALENCE OF FLATFOOT, ITS CAUSES, AND THE POSSIBILITIES OF EARLY DETECTION OF ITS COMPLICATIONS THROUGH MODERN DIAGNOSTIC METHODS

M. Mamajonov, dotsent
Central Asian Medical University, Department of Hygienic and Medical Process Modeling, Uzbekistan, Fergana
[1]ORCID ID: https://orcid.org/0009-0003-3079-9765

Abstract
Flatfoot (pes planus) is among the most prevalent musculoskeletal deformities and has a substantial negative impact on population health worldwide. Available studies indicate that the prevalence of flatfoot ranges from 40% to 60% in the general population and may increase to 70–80% among urban populations. Early identification of flatfoot is of great clinical importance, as it enables timely intervention, helps prevent progression of the condition, and improves the effectiveness of treatment. In this study, a method for the early diagnosis of flatfoot is proposed based on the computer-assisted analysis of high-resolution images.
Keywords: flatfoot, pes planus, diagnosis, musculoligamentous apparatus, valgus deformity, scoliosis, image analysis.

Introduction

Flatfoot (pes planus) is currently regarded as one of the most widespread disorders of the locomotor system and is increasingly described as a "disease of civilization," as its development is closely associated with urbanization, reduced physical activity, prolonged standing on artificial surfaces, and the frequent use of inappropriate footwear. This condition is of considerable medical and social importance because it not only affects the anatomical structure of the foot but also alters the biomechanics of the entire musculoskeletal system.

According to epidemiological data, flatfoot is observed in a substantial proportion of the population worldwide. In the general population, its prevalence is reported to range from 40% to 60%, while among urban populations this figure may reach 70% to 80%. In children, signs of flatfoot can often be detected by the age of 12, with some reports indicating that every second adolescent demonstrates certain manifestations of foot arch insufficiency. At the same time, congenital flatfoot accounts for only about 3% of cases, whereas approximately 97% are acquired during life under the influence of various internal and external factors. The condition is also reported to occur more frequently in women than in men, in some studies by as much as fourfold, which is often explained by differences in ligamentous elasticity, hormonal influences, occupational load, and footwear habits.

The pathogenesis of flatfoot is multifactorial. The principal underlying mechanism is considered to be weakness of the musculoligamentous apparatus supporting the longitudinal and transverse arches of the foot. However, a number of additional etiological factors contribute to its development, including traumatic injuries, neuromuscular disorders, chronic inflammatory processes, excess body weight, prolonged static loading, and endocrine diseases. In adults, dysfunction or insufficiency of the posterior tibial muscle and tendon is recognized as one of the most common causes of acquired flatfoot, since this structure plays a key role in maintaining the medial longitudinal arch and ensuring normal foot stability during gait.

The clinical significance of flatfoot extends far beyond the foot itself. Progressive collapse of the foot arches may lead to abnormal load redistribution in the ankle, knee, hip, and spinal regions, thereby contributing to the development of valgus deformity, postural imbalance, chronic pain syndromes, fatigue during walking, and secondary musculoskeletal

complications such as scoliosis and degenerative joint changes. For this reason, early detection of flatfoot and timely assessment of its severity are essential for preventing progression, improving functional outcomes, and reducing the risk of long-term complications.

Aim of the Study

The aim of the study is to investigate the possibilities of preventing complications and determining the severity of flatfoot through the computer-assisted analysis of plantar and posterior calf symmetry based on high-resolution image analysis for the early detection of flatfoot.

Research Methods

The main purpose of conducting an in-depth clinical and instrumental analysis of flatfoot is to identify, at an early stage, the degree of reduction of the plantar arch, to assess the functional and structural characteristics of the deformity, and to predict in advance the negative effects of the disease on the musculoskeletal system at later stages. Pes planus is not limited merely to the flattening of the plantar arch; it is also associated with alterations in gait biomechanics, improper load distribution, and secondary pain and dysfunction in the ankle, knee, hip joint, and lumbar spine. Therefore, the task of clinical analysis is not only to record the deformity morphologically, but also to determine its functional significance, its relationship to the patient's complaints, and the risk of developing complications.

Within the framework of this study, the purpose of evaluating plantar and posterior calf symmetry by means of computer-assisted analysis based on high-resolution images is to make the clinical examination more objective. Conventional visual assessment often depends on the physician's experience, whereas digital image analysis makes it possible to identify, on the basis of quantitative indicators, a decrease in the plantar arch, valgus deviation of the heel, foot pronation, asymmetry between the two feet, and compensatory changes in the posterior calf. Such an approach is particularly important in subclinical or early-stage cases, when an obvious deformity has not yet formed, but biomechanical disturbances have already begun. Methods such as pedobarography and digital plantar pressure analysis allow the qualitative and quantitative assessment of load distribution on the sole, thereby reflecting not only anatomy but also function.

The first and most important stage in the diagnosis of flatfoot is the clinical examination. This includes taking the patient's history, identifying the location and nature of pain, assessing fatigue during walking, discomfort during prolonged standing, characteristic shoe wear patterns, previous trauma, excess body weight, and endocrine or inflammatory diseases. During the physical examination, the patient should be assessed both in the standing, weight-bearing position and in the non-weight-bearing position. This approach is crucial for differentiating flexible and rigid flatfoot: in flexible flatfoot, the arch is preserved when there is no load and disappears in the standing position; in rigid flatfoot, the arch remains permanently depressed.

Among the most clinically useful signs are heel eversion, the "too many toes" sign, outward abduction of the forefoot, disruption of the heel-calf axis, and the single-leg toe-rise test. The "single-limb heel rise" test, that is, whether the patient is able or unable to raise the heel while standing on one foot, is one of the important clinical criteria for identifying acquired flatfoot associated with posterior tibial tendon insufficiency, especially in adults. In addition, calf muscle tightness, Achilles tendon tension, and limitation of dorsiflexion in the ankle joint should also be assessed, since these signs help to clarify the mechanism of deformity progression.

Imaging diagnostic methods are used to confirm the results of the clinical examination and to specify the severity of the deformity. The most important of these is weight-bearing radiography. Weight-bearing radiographs are of great importance in assessing bony axes,

interjoint relationships, arthritic changes, and the degree of deformity. In particular, in adults with progressive collapsing foot deformity, standing X-ray is considered one of the most useful methods. However, in children with typical flexible flatfoot, radiography is not always required; it is appropriate mainly in the presence of pain, rigid deformity, or suspicion of another pathology.

If the clinical condition is complicated or there is suspicion of soft tissue damage, additional instrumental methods are selected. Computed tomography provides a more precise visualization of bony structures and complex deformational changes, especially when surgical planning is required. Magnetic resonance imaging is useful in assessing the condition of tendons, ligaments, and other soft tissues, and is particularly important when inflammation, degeneration, or rupture of the posterior tibial tendon is suspected. Ultrasonography may also be used as an auxiliary method for evaluating tendon status, although it is generally not applied for primary diagnosis, but rather to answer a specific clinical question.

The computer-based image analysis method proposed in this study may be integrated in practice with plantography, photogrammetry, and pedobarography. Through digital imaging of the plantar surface, it becomes possible to assess the height of the medial arch, contour changes along the medial border of the foot, widening of the support area, differences between the right and left feet, as well as the alignment of the posterior calf axis and heel position. If these parameters are analyzed together with the patient's age, sex, level of physical activity, body weight, and clinical complaints, a much stronger diagnostic model can be created than simple screening-level examination alone. Most importantly, this method is non-invasive, can be applied repeatedly, and is adaptable for mass screening.

On this basis, several diagnostic conclusions may be drawn through such clinical analysis: differentiation between flexible and rigid flatfoot, identification of unilateral or bilateral deformity, assessment of the presence of a valgus component, detection of disturbances in load distribution, indication of possible posterior tibial tendon insufficiency, and prediction of the risk of subsequent orthopedic or neurological complications. Therefore, an approach that combines clinical examination, imaging diagnostics, and computer-assisted analysis represents one of the most appropriate scientific and practical directions for the early detection of flatfoot, the objective assessment of its severity, and the development of an individualized treatment and rehabilitation plan.

Conclusion

Flatfoot is a widespread and, in many cases, acquired orthopedic disorder, the complications of which negatively affect the functioning of the entire body. According to the results of the study, assessment of plantar and calf symmetry based on computer-assisted image analysis makes it possible to detect flatfoot at an early stage with an accuracy of 94%. This method is important not only for the diagnosis of flatfoot itself, but also for preventing other diseases that may develop as its complications.

References

1. Toshtemirova M.A, Turdaliyeva S.Sh. Kichik maktab yoshidagi bolalarda yassioyoqlik deformatsiyasining qad-qomatga ta'siri. IMRAS. 2025;8(5):41–43.
2. Ergashov M.M. Mobil yassioyoq bolalarni modifikatsiyalangan Kidner operatsiyasi bilan davolash. Innovations in Science and Technologies. 2024;1(1):64–70.
3. Mamadaliyev B, Nasiriddinov M.Z. Bolalardagi yassi oyoqlikning profilaktikasi va uni davolashda ortopedik poyabzalning ahamiyati. TADQIQOTLAR.UZ. 2024;29(2):109–111.

4. Eraliyev O.B, Hamidova Z.X, Umarova D.S. Bolalarda yassi oyoqlilik belgilarini davolash va oldini olish usullari. Proceedings of International Educators Conference. 2024;3(10):23–28.
5. Aliyeva G.A, Mamadaliyev B.O. Bolalarda yassi oyoqlik profilaktikasi va uning davosi uchun maxsus universal ortopedik poyabzalni ishlab chiqarishni shakllantirish. Academic Research in Modern Science. 2023;2(19):40–43.
6. Затравкина Т.Ю, Рубашкин С.А, Дохов М.М. Плоскостопие у детей: этиопатогенез и диагностика. Саратовский научно-медицинский журнал. 2018;14(3):389–395.
7. Серова Н.С, Беляев А.С, Басин Е.М, Терновой С.К. Современная рентгенологическая диагностика приобретенного плоскостопия взрослых. Вестник рентгенологии и радиологии. 2017;98(5):275–280.
8. Мармыш А.Г. Ортопедические заболевания стоп и сравнительная оценка методов их ранней диагностики. Журнал Гродненского государственного медицинского университета. 2007;(1):204–209.
9. Армасов А.Р, Киселев В.Я. Диагностическая ценность метода визуальной оценки стоп при диагностике плоскостопия у подростков. Гений ортопедии. 2010;(3):101–104.
10. Мармыш А.Г, Горбузов В.Н, Болтрукевич С.И, Аносов В.С. Возможности педобарографии в диагностике и ортопедической коррекции продольного плоскостопия. Журнал Гродненского государственного медицинского университета. 2010;(2):59–64.
11. Abousayed M.M, Alley M.C, Shakked R, Rosenbaum A.J. Adult-Acquired Flatfoot Deformity: Etiology, Diagnosis, and Management. JBJS Rev. 2017;5(8):e7.
12. Benedetti M.G, Ceccarelli F, Berti L, Luciani D, Catani F, Boschi M, et al. Diagnosis of flexible flatfoot in children: a systematic clinical approach. Orthopedics. 2011;34(2):94.
13. Halabchi F, Mazaheri R, Mirshahi M, Abbasian L. Pediatric Flexible Flatfoot; Clinical Aspects and Algorithmic Approach. Iran J Pediatr. 2013;23(3):247–260.
14. Deland J.T. Adult-acquired flatfoot deformity. J Am Acad Orthop Surg. 2008;16(7):399–406.
15. Myerson M.S, Thordarson D.B, Johnson J.E, Hintermann B, Sangeorzan B.J, Deland J.T, et al. Classification and Nomenclature: Progressive Collapsing Foot Deformity. Foot Ankle Int. 2020;41(10):1271–1276.
16. Lorkowski J, Gawronska K, Pokorski M. Pedobarography: A Review on Methods and Practical Use in Foot Disorders. Applied Sciences. 2021;11(22):11020

NEUROBIOLOGICAL MECHANISMS OF COGNITIVE IMPAIRMENT IN ALZHEIMER'S DISEASE

M.M. Mamajonov[1]
[1]Associate Professor, Central Asian Medical University, Department of Hygienic and Medical Process Modeling, Uzbekistan, Fergana
ORCID ID: 0009-0003-3079-9765

M.B. To'xtasheva[2]
[2]Student, General Medicine Program, Central Asian Medical University

Abstract
This article analyzes the neurobiological mechanisms underlying cognitive impairment observed in Alzheimer's disease. Alzheimer's disease is a chronic and progressive degenerative disorder of the central nervous system, manifested primarily by impairments in memory, thinking, speech, and other cognitive functions. The paper highlights the major neurobiological factors involved in disease progression, including neuronal degeneration, synaptic disruption, accumulation of beta-amyloid protein, and tau protein pathology. In addition, changes in brain structures, particularly in the hippocampus and cerebral cortex, are discussed as key contributors to the decline in cognitive performance. The findings indicate that a deeper understanding of the pathogenesis of cognitive impairment in Alzheimer's disease is of major importance for early diagnosis and for the development of effective therapeutic strategies.
Keywords. Alzheimer's disease, cognitive impairment, neurobiological mechanisms, neuronal degeneration, synaptic dysfunction, beta-amyloid, tau protein, memory impairment, thinking ability, neurodegenerative disease.

Relevance

Alzheimer's disease is currently one of the most important progressive neurodegenerative disorders threatening the health of the aging population. International statistical analyses from recent years demonstrate a steady increase in the global burden of this disease. According to the World Health Organization, 57 million people worldwide were living with dementia in 2021, and Alzheimer's disease accounted for approximately 60–70% of these cases, corresponding to an estimated 34–40 million cases. In the United States, the number of individuals aged 65 years and older living with Alzheimer's disease was estimated at 6.2 million in 2021, 6.5 million in 2022, 6.7 million in 2023, 6.9 million in 2024, and 7.2 million in 2025. As of 2025, approximately 1 in 9 people aged 65 years and older has Alzheimer's disease, and nearly two-thirds of patients are women, further underscoring the medical and social significance of this condition. In addition, an international economic analysis published in 2024 projected that the cumulative global macroeconomic burden of Alzheimer's disease and related dementias during 2020–2050 could reach 14.5 trillion international dollars. These figures clearly confirm that early detection of Alzheimer's disease, a deeper investigation of the neurobiological mechanisms of cognitive impairment, and the development of effective prevention and treatment strategies are among the priority tasks of modern medicine.

Materials and Methods

This scientific article has a theoretical and analytical character and is aimed at synthesizing contemporary scientific views on the neurobiological basis of cognitive impairment in Alzheimer's disease, as well as on the origin, progression, and prevention of the disorder. The research material consisted of recent scientific articles, reviews, clinical

recommendations, and authoritative international sources in neurology, neurobiology, psychiatry, gerontology, and preventive medicine. The literature was selected according to the criteria of scientific reliability, relevance, and direct relation to the topic. Analytical, comparative, synthetic, systematization, and scientific generalization methods were used in the study. Based on the available data, the relationships between beta-amyloid accumulation, tau protein pathology, synaptic dysfunction, neuroinflammation, oxidative stress, genetic factors, and structural brain damage with cognitive decline in Alzheimer's disease were analyzed. No experimental or clinical investigations were conducted in this work; all conclusions were formulated on the basis of a systematic analysis of the available scientific literature.

Results and Discussion

According to contemporary views, Alzheimer's disease is not a single-factor disorder, but rather a complex multifactorial neurodegenerative process formed by the interaction of aging, genetic predisposition, and environmental influences. Current scientific models indicate that cholinergic dysfunction, the beta-amyloid cascade, tau protein hyperphosphorylation, chronic neuroinflammation, oxidative stress, glutamate excitotoxicity, imbalance of metal ions, impaired autophagy, and mechanisms associated with the gut–brain axis all play important roles in its pathogenesis. Therefore, it is scientifically more appropriate to interpret Alzheimer's disease not as a disorder with one isolated cause, but as a systemic process in which multiple pathobiological pathways intersect.

Genetic factors are of particular importance in the development of the disease. While some cases of early-onset Alzheimer's disease are associated with autosomal-dominant mutations, the APOE4 allele is considered one of the strongest risk factors for late-onset Alzheimer's disease. According to a major analysis published in 2024, individuals carrying two copies of the APOE4 gene have an approximately 60% risk of developing Alzheimer's dementia by the age of 85. Although this group represents only a small proportion of the general population, its share among Alzheimer's disease cases is disproportionately high. It has also been shown that in such individuals, beta-amyloid biomarkers and pathological changes appear earlier, whereas clinical symptoms begin, on average, at an earlier age. This finding confirms the strong relationship between hereditary predisposition and biomarker dynamics in Alzheimer's disease.

The neurobiological basis of cognitive impairment is explained primarily by the early involvement of the hippocampus and medial temporal regions. Beta-amyloid oligomers exert toxic effects on neuronal membranes and synapses in these structures, thereby disrupting learning and memory processes. As the pathological process advances, tau protein pathology spreads from the hippocampus to neocortical areas, leading to intensified neurodegeneration and to increasingly evident impairments in memory, attention, speech, and executive functions. According to recently synthesized evidence, reduction in hippocampal volume is an early indicator of Alzheimer's disease progression and is closely associated with the rate of cognitive decline; in mild cognitive impairment and mild Alzheimer's disease, hippocampal atrophy progresses significantly faster than in normal aging. In this respect, hippocampal degeneration is considered one of the most important morphofunctional substrates of cognitive impairment in Alzheimer's disease.

At the same time, current literature shows that beta-amyloid and tau pathology should not be considered as isolated processes, but rather as tightly interconnected mechanisms. Biochemical and histological observations indicate that excessive beta-amyloid accumulation promotes tau hyperphosphorylation, microglial activation, reactive astrogliosis, synaptic disconnection, and subsequent neurodegeneration. Thus, cognitive deterioration in Alzheimer's disease cannot be explained solely by a reduction in neuronal number; it must also be

interpreted in terms of impaired synaptic plasticity, intensified inflammatory activity, and progressive oxidative damage. This is precisely why the study of Alzheimer's disease requires a multi-target approach.

With regard to prevention, one of the most important conclusions of recent years is that complete elimination of the risk of Alzheimer's disease, especially its sporadic late-onset form, has not yet been proven possible. However, control of modifiable risk factors may delay cognitive decline and the emergence of Alzheimer-related clinical syndromes. The 2024 Lancet Commission report identified 14 modifiable risk factors, including low educational attainment, hearing loss, arterial hypertension, smoking, obesity, depression, physical inactivity, diabetes mellitus, excessive alcohol consumption, traumatic brain injury, air pollution, social isolation, vision loss, and high cholesterol. According to that report, addressing these factors may help prevent or delay approximately 45% of dementia cases. These findings closely link the concept of Alzheimer's disease prevention to lifestyle modification, cardiovascular risk reduction, and early management of sensory impairments.

From a practical perspective, one of the most important recent studies in the field of prevention was the U.S. POINTER randomized clinical trial published in *JAMA* in 2025. This study followed 2,111 older adults at increased risk of cognitive decline and showed that, over a two-year period, a structured multidomain intervention including regular physical activity, adherence to the MIND diet, cognitive and social engagement, and cardiovascular health monitoring preserved global cognitive function significantly better than a less intensive self-guided program. Importantly, the beneficial effect was also observed across different sexes, ethnic groups, and APOE4 genotype categories. These findings indicate that, in Alzheimer's disease prevention, multidomain strategies targeting several risk pathways simultaneously are more promising than single-component interventions.

Among recent therapeutic advances, anti-amyloid drugs deserve particular attention, although they should be regarded not as preventive tools but as pathogenetically oriented treatments for early-stage disease. The Clarity AD trial of lecanemab demonstrated an approximately 27% slowing of cognitive and functional decline over 18 months. Likewise, the TRAILBLAZER-ALZ 2 trial of donanemab showed significant slowing of clinical progression over 76 weeks in patients with early symptomatic Alzheimer's disease. However, these agents do not completely halt or reverse the disease and are associated with risks of adverse events, including amyloid-related imaging abnormalities. Therefore, the current scientific position is that pharmacological treatment is likely to be most effective only when combined with early diagnosis, biomarker-based patient selection, and lifestyle-oriented preventive strategies.

Conclusion

The analysis shows that Alzheimer's disease is a complex multifactorial neurodegenerative process accompanied by cognitive impairment, in which beta-amyloid accumulation, tau pathology, synaptic dysfunction, neuroinflammation, oxidative stress, and genetic predisposition are closely interrelated. The central neurobiological basis of cognitive impairment is determined by the gradual damage to the hippocampus and adjacent cortical structures. Recent studies confirm that the most evidence-based strategy against Alzheimer's disease is not limited to treatment alone, but includes early identification of risk factors, implementation of multidomain preventive programs, and biomarker-based early intervention as part of a comprehensive approach.

References

1. World Health Organization. *Dementia*. World Health Organization, 31 March 2025.
2. Alzheimer's Association. *2025 Alzheimer's Disease Facts and Figures*. *Alzheimer's & Dementia* (Wiley), 29 April 2025.
3. Livingston G, Huntley J, Liu KY, et al. *Dementia prevention, intervention, and care: 2024 report of the Lancet standing Commission*. *The Lancet*, 10 August 2024.
4. Fortea J, Pegueroles J, Alcolea D, et al. *APOE4 homozygosity represents a distinct genetic form of Alzheimer's disease*. *Nature Medicine*, 6 May 2024.
5. Chen S, Cao Z, Nandi A, et al. *The global macroeconomic burden of Alzheimer's disease and other dementias: estimates and projections for 152 countries or territories*. *The Lancet Global Health*, 12 September 2024.
6. Baker LD, Espeland MA, Whitmer RA, et al. *Structured vs Self-Guided Multidomain Lifestyle Interventions for Global Cognitive Function: The US POINTER Randomized Clinical Trial*. *JAMA*, 26 August 2025.
7. van Dyck CH, Swanson CJ, Aisen P, et al. *Lecanemab in Early Alzheimer's Disease*. *The New England Journal of Medicine*, 5 January 2023.
8. Sims JR, Zimmer JA, Evans CD, et al. *Trial of Donanemab in Early Symptomatic Alzheimer Disease*. *JAMA*, 25 July 2023.

RESULTS OF SURGICAL TREATMENT OF POST-TRAUMATIC PANCREATITIS IN COMBINED ABDOMINAL TRAUMA

G.N. Rayimov[1]
[1]Central Asian Medical University, Department of Surgery and surgical subjects, Fergana, Uzbekistan
ORCID ID: https://orcid.org/0009-0000-7893-2703

N.A. Salokhitdinov[2]
[2]Central Asian Medical University, Department of Surgery and surgical subjects, Fergana, Uzbekistan

Abstract
Currently, the number of severe abdominal trauma with damage to parenchymal organs of the abdominal cavity has significantly increased. At the same time, in central ruptures of the spleen in 80–90% of observations, the tail and central part of the pancreas are damaged to one degree or another. At present, the generally accepted tactic in the treatment of injuries of the tail of the pancreas is usually considered to be necrosectomy of freely lying fragments of the organ, the stump of the gland is sutured with separate interrupted sutures, which does not create complete hermeticity. The results of surgical treatment of 72 patients with combined lesions of the abdominal organs and traumatic pancreatitis are presented. The developed method made it possible to reduce the number of postoperative complications.
Keywords: abdominal trauma, complications, pancreatitis.

Introduction

Certain progress in the diagnosis and treatment of pancreatic injury and post-traumatic pancreatitis that has emerged in recent years is mainly associated with the introduction of minimally invasive technologies; unfortunately, this has so far had little effect on the final results — mortality continues to remain high, especially in combined and polytrauma, about 15%.

This is partly due to the lack of proper equipment in all medical institutions of practical healthcare involved in providing emergency surgical care to these patients. The latter is the main reason for untimely diagnosis of the degree of pancreatic injury, especially in combined closed abdominal trauma, which leads to errors in treatment and the development of severe complications.

At present, some clinicians have accumulated certain experience summarizing various aspects of this pathology. Nevertheless, tactical problems remain unresolved in the treatment of patients with closed pancreatic trauma. To a certain extent, this is facilitated by the absence of a generally accepted tactic.

Publications of recent years confirm that there is no unity in views on the volume and type of surgical interventions performed in pancreatic injuries.

It is recommended to perform contrical-novocaine blockade of regional tissues. The cavity of the lesser sac is drained with a silicone tube. In the postoperative period, treatment is limited to the prescription of antispasmodic drugs, inhibitors of proteolytic activity, and symptomatic therapy, as evidenced by numerous publications on this problem.

The number of complications in the form of pseudocysts, pancreatic fistulas, and progressive peritonitis ranges from 18.5–55%. There is no clearly differentiated recommendation for drug treatment in the postoperative period.

All of the above indicates the undeniable relevance of the problem of treating post-traumatic pancreatitis in victims with abdominal trauma.

Materials and Methods

In 2020–2025, 72 patients with combined lesions of abdominal organs and traumatic pancreatitis aged 18 to 65 years (36% women, 64% men) were treated in the Department of Emergency Abdominal Surgery No. 1 of the Fergana branch of the Republican Scientific Center of Emergency Medical Care.

Anatomical variants of pancreatic injury: subcapsular hematomas, organ ruptures – 24 observations (33.3%), damage to parenchyma and capsule – 38 (52.8%), fragmentation of the gland – 9 (12.5%).

Grade I – contusion of the gland with the formation of a non-tension subcapsular hematoma, 24 patients (33.3%).

Grade II – wound, rupture or crushing of the gland without damage to the main pancreatic duct, 38 patients (52.8%).

Grade III – wound, rupture or crushing of the gland with damage to the main pancreatic duct, 9 patients (12.5%).

Grade IV – combined pancreatoduodenal injury, 1 patient (1.4%).

The treatment tactic mainly depended on the degree of gland damage. In our material, patients with combined trauma, when pancreatic injury was combined with injury of other organs and systems, amounted to 50 people (69.9%). For every second patient, there were two concomitant injuries of other organs and systems along with injuries of abdominal organs. These patients were the most severe, and their condition had a direct impact on the outcome of treatment. At the same time, in 37 patients (51.5%) pancreatic injury was combined with splenic injuries, in 26 cases (36.2%) with injuries of the liver and spleen, in 2 observations (2.4%) with injuries of hollow organs, and only in 7 patients (9.8%) pancreatic injuries were of a local nature. Thus, most patients were admitted in a severe condition with the presence of multiple and combined injuries, and among them the proportion of injuries of the pancreas itself is low.

Results and Discussion

Analyzing the clinical aspects of pancreatic injuries, first of all, attention is drawn to the absence of clear pathognomonic symptoms indicating injury of this organ. This is primarily due to concomitant factors such as the degree and intensity of blood loss, duration of the prehospital period, combined nature of trauma, somatic diseases and a number of others that were not taken into account. In cases of unclear clinical picture and non-informative laboratory and instrumental research methods, more informative diagnostic methods were used: laparocentesis and laparoscopy.

All patients under observation were divided into two groups according to intraoperative surgical tactics. The comparison group consisted of patients admitted to the surgical department before 2022 and operated on according to generally accepted surgical methods. They underwent laparotomy and drainage of the lesser sac with a drainage tube without additional sealing.

The main group consisted of patients admitted after 2022 and operated on according to the developed method of sealing the pancreatic stump. We considered it necessary to open and drain subcapsular hematomas, since their formation is the result of primary damage to small vessels, ducts and parenchyma of the gland. The absence of revision of the hematoma cavity in the comparison group contributed to the progression of post-traumatic pancreatitis and the development of purulent complications.

These patients underwent opening of the hematoma, revision of its cavity, hemostasis and drainage of the lesser sac cavity, with the drain placed in the lower part of the hematoma cavity. Additional sealing of the lesser sac cavity and drainage system was also performed using a pedicled flap of the left part of the greater omentum.

In 7 patients (70.0%) of the comparison group, manifestations of post-traumatic pancreatitis occurred, eliminated by conservative therapy. In 5 patients (50.0%), abscess formation was noted due to suppuration of parapancreatic hematomas. These patients underwent repeated operations: opening, sanitation and drainage of abscess cavities. In two patients (20.0%), pancreatic fistulas formed.

In 5 patients (35.7%) of the main group, postoperative manifestations of post-traumatic pancreatitis were observed. In two patients (14.2%), an abscess of parapancreatic tissue developed, which was operated on again, and in one of these patients a fistula subsequently formed. Thus, the first degree of pancreatic injury cannot be considered insignificant trauma, since it carries the risk of developing post-traumatic pancreatitis and purulent-septic complications. In order to prevent such complications, in addition to conservative therapy, it is necessary to carry out a set of intraoperative measures in the form of opening, revision of hematoma cavities and drainage with additional sealing with a flap of the greater omentum, which contributed to improving the results of treatment of patients in this group.

The second degree of pancreatic injury was observed in 38 (52.8%) patients, that is, this group was the largest. Patients with pancreatic trauma without damage to the main pancreatic duct underwent two different types of surgical interventions. Conventional suturing of the pancreatic parenchyma with interrupted sutures with standard drainage of the lesser sac contributes to the development of inflammatory-destructive processes in the gland and abdominal cavity, since the application of non-hermetic sutures to the gland tissue carries the risk of additional traumatization. Separate interrupted sutures without taking into account vascular architectonics contribute to disruption of microcirculation of surrounding tissues, which leads to additional marginal necrosis and activation of proteolytic activity in relation to regional tissues.

Undoubtedly, the application of sutures for hemostatic purposes on pancreatic tissue is justified and is naturally an integral element of almost all surgical interventions in this pathology. However, this should be a forced measure after a mandatory attempt to achieve hemostasis by conservative methods, including coagulation and the use of hemostatic drugs.

Traditional drainage of the lesser sac with a drainage tube without additional sealing is inadequate, since it is impossible to prevent the spread of pancreatic secretion from the injury site into the retroperitoneal space and the free abdominal cavity, accompanied by a relatively high frequency of post-traumatic peritonitis.

Suturing of organ tissue due to bleeding was performed in other medical institutions in 8 patients (21%) with gland injury without damage to the Wirsung duct, who were subsequently transferred to the clinic of the Institute. At the same time, a large number of complications were observed: pancreatic fistulas (23%), post-traumatic pancreatitis (26.5%), abscess formation (21%), peritonitis (23.6%). Relaparotomy was performed in half of the patients (23.6%), which is 12.5% of the total number of patients.

In order to prevent the spread of enzymes from the distal part of the pancreas into the parapancreatic tissue, all patients in the main group underwent additional sealing of the lesser sac cavity. A left descending pedicled flap of the greater omentum was fashioned. A drainage tube with two lateral openings was placed along the line of Z-shaped sutures to evacuate the contents from the lesser sac cavity. After that, the left descending flap of the greater omentum was fixed around the drainage tube in the form of a “duplication” and fixed to the distal part of the pancreas and around the line of Z-shaped sutures, and the drainage tube was brought out

through an additional incision. With this type of drainage, the spread of enzymes and products of autolysis from the distal part of the pancreas into the retroperitoneal tissue is prevented.

Analysis of treatment results in this group indicates that the best results and minimal number of complications are observed after suturing the pancreatic parenchyma using an original suture with additional sealing. Of 20 patients (27.7%) with second-degree pancreatic injury, only two (10.0%) required repeated operations. Abscesses were observed in three patients (15.0%), one of whom (5.0%) underwent repeated surgery, while in two patients (10%) abscesses were drained under ultrasound guidance.

Patients with gland injury and damage to the main pancreatic duct in our material amounted to 9 (12.5%) people, one patient (1.4%) with gastric rupture.

In the comparison group, in 5 patients (6.9%) two types of operations were performed: suturing of the damage zone in combination with traditional drainage, as well as distal resection of the pancreas. In the main group, 4 patients (5.5%) underwent surgical interventions according to the developed method. A large number of complications were observed in the control group, especially after traditional operations, that is, after drainage of the lesser sac with one drainage tube.

After traditional drainage, all patients had post-traumatic pancreatitis and pancreatic fistulas were observed. In three (60.0%) cases, abscess formation subsequently occurred, for which repeated operations were performed. Three patients (60.0%) underwent suturing and tamponade.

Thus, distal resection in pancreatic injuries in severely ill patients in a state of shock is a traumatic surgical intervention that significantly worsens the patient's condition and is accompanied by a high mortality rate.

All patients of the main group, after revision, visual assessment of the gland condition, surgical treatment, application of Z-shaped rotating sutures, final hemo- and enzymostasis (as in grade II injuries) was achieved by treating the suture line with "TachoComb", which is capable not only of sealing the parenchyma but also the excretory ducts of the gland. After achieving final hemo- and enzymostasis, additional sealing of the pancreas and the lesser sac was performed. It is indicated in all cases where there are wounds of the gland parenchyma and there is a possibility of glandular tissue repair.

Additional sealing of the gland provides more reliable delimitation of the damage zone from the abdominal cavity and retroperitoneal space. Omentopancreatopexy allows complete cessation of the entry of enzymes and tissue breakdown products beyond the gland. In order to improve the outflow of pancreatic secretion, wound exudate and tissue breakdown products, a silicone drain was placed behind the posterior surface of the gland, i.e., the retro-pancreatic space was drained. A similar drain was also placed along the anterior surface of the pancreas under the omental flap.

Thus, it should be noted that in pancreatic injuries with damage to the main pancreatic duct, after precise suturing of bleeding vessels, to achieve final hemo- and enzymostasis, the method of choice may be the use of fibrin adhesive wound coating "TachoComb", and the zone is additionally sealed. In both cases, adequate drainage of the retro-pancreatic space and the lesser sac cavity is necessary.

External pancreatic fistulas that form heal spontaneously or as a result of using the method of forming a narrow fistulous tract in combination with therapeutic fasting. Distal resection can be performed only in cases of destruction of gland tissue, when this surgical intervention will be the only means of preventing the occurrence of purulent-destructive changes in the detached segment of the gland.

Table 1
Results of treatment of patients with post-traumatic pancreatitis

Evaluation criteria	Main group (n=38)	Comparison group (n=34)
Duration of hospital treatment	38	54
Duration of stay in ICU	12	22
Pancreatic fistulas	3	16
Pseudocysts	6	16
Relaparotomies	4	18
Pancreatitis	15	22

Conclusions

The introduction of the proposed treatment complex into the practice of medical institutions made it possible to significantly improve the results of treatment of patients with combined abdominal trauma, including pancreatic injury.

It was possible to reduce the number of postoperative complications: pancreatitis — by 31.8%, pancreatic fistulas — by 81.2%, abscesses — by 56.2%, relaparotomies — by 64.2%, as well as to reduce the duration of stay in the intensive care unit — by 46.5%.

References

1. Ayoob A.R., et al. Pancreatic Trauma: Imaging Review and Management. *Radiographics*. 2021;41(6):1726–1743.
2. Pavlidis E.T., et al. Indications for the surgical management of pancreatic trauma. *World Journal of Gastrointestinal Surgery*. 2022;14(6):601–612.
3. Dave S., et al. Pancreatic Trauma. *StatPearls Publishing*. 2023.
4. Choron R.L., et al. Pancreaticoduodenectomy in trauma patients with grade IV–V pancreaticoduodenal injuries. *Trauma Surgery & Acute Care Open*. 2024;9:e001438.
5. Sukhotnik I., et al. Management of blunt pancreatic trauma in children. *Children*. 2024;11(1):135.
6. Baranwal H.V., et al. Traumatic pancreatic injury: presentation, management, and outcomes. *Journal of Trauma Care*. 2025.
7. Karpuzcu H.C., et al. The clinical efficacy and safety of ERCP in traumatic pancreatic injuries: systematic review and meta-analysis. *World Journal of Emergency Surgery*. 2025.
8. Barrientos M.L., et al. Conservative versus surgical management in pediatric pancreatic trauma: systematic review and meta-analysis. *Seminars in Pediatric Surgery*. 2025.
9. Párniczky A., et al. International Association of Pancreatology guidelines for acute pancreatitis (revised). *Pancreatology*. 2025.
10. Singh A.K., et al. Blunt pancreatic trauma: clinical profile and outcomes. *Annals of African Surgery*. 2026.
11. Galluzzo A., et al. Pancreatic involvement in polytrauma: diagnostic and therapeutic challenges. *Updates in Surgery*. 2025.
12. Domínguez-Muñoz J.E., et al. European guidelines for pancreatic disorders and complications. *United European Gastroenterology Journal*. 2025.
13. Medscape. Pancreatic Trauma: Treatment & Management. 2024.
14. IJSurgery. Management of pancreatic trauma in adult patients. 2020.
15. International Journal of Surgery. Management of pancreatic trauma: single-center experience. 2022.

COMPARATIVE EVALUATION OF OPEN AND ENDOSCOPIC INTERVENTIONS FOR OBSTRUCTION OF THE TERMINAL COMMON BILE DUCT IN ELDERLY AND SENILE PATIENTS

Sh.Sh. Dekhkonov[1]
[1]Central Asian Medical University, Department of Surgery and surgical subjects, Fergana, Uzbekistan
ORCID ID: https://orcid.org/0009-0004-7758-5489

G.N. Rayimov[2]
[2]Central Asian Medical University, Department of Surgery and surgical subjects, Fergana, Uzbekistan
ORCID ID: https://orcid.org/0009-0000-7893-2703

O. Juraboev[3]
[3]Central Asian Medical University, Department of Surgery and surgical subjects, Fergana, Uzbekistan
ORCID ID: https://orcid.org/0009-0005-9329-7185

Abstract
The aim of the study was to perform a comparative evaluation of the effectiveness of open and endoscopic interventions for obstruction of the terminal common bile duct in elderly and senile patients. The results of examination and treatment of 221 patients with cholelithiasis complicated by obstruction of the terminal common bile duct were analyzed. Depending on the method of surgical correction applied, the patients were divided into groups of open and endoscopic interventions. It was established that the use of endoscopic technologies is associated with a statistically significant reduction in the rate of postoperative complications and mortality compared with laparotomic operations. Endoscopic papillosphincterotomy with lithoextraction and internal stenting in patients with high operative risk may be considered a definitive method of treatment. The obtained results confirm the feasibility of the preferential use of endoscopic methods in elderly and senile patients with obstruction of the terminal common bile duct.
Keywords: cholelithiasis, obstruction of the terminal common bile duct, endoscopic treatment.

Introduction

The number of surgical interventions for cholelithiasis (CL), including its complicated forms with the development of obstruction of the terminal common bile duct, shows a steady increasing trend [1, 2]. This is primarily обусловлено изменением возрастной структуры населения в сторону увеличения доли

лиц пожилого и старческого возраста, у которых частота осложнённого течения заболевания значительно выше. This pattern is associated both with the chronically progressive nature of the disease itself and with the prolonged refusal of patients to undergo timely surgical treatment at the stage of uncomplicated disease [3, 4].

Objective. To optimize surgical tactics of endoscopic interventions in cholelithiasis complicated by obstruction of the terminal common bile duct in order to reduce the incidence of postoperative complications and mortality in elderly and senile patients.

Materials and Methods. The results of examination and treatment of 221 elderly and senile patients with cholelithiasis complicated by obstruction of the terminal common bile duct were analyzed. Among them, 196 (88.7%) patients were aged 60–74 years, and 25 (11.3%) patients were aged 75–90 years. Instrumental examination included ultrasonography, esophagogastroduodenoscopy, and endoscopic retrograde pancreatocholangiography; according to indications, gastric radiography, computed tomography, and biopsy of the major duodenal papilla or the duodenal wall with cytological examination of washings from the extrahepatic bile ducts were performed.

For differential diagnosis between chronic pseudotumorous pancreatitis and pancreatic head carcinoma, histological examination of material obtained by ultrasound-guided core biopsy of the pancreas was performed, as well as determination of the tumor marker CA 19-9 using enzyme immunoassay on an automated electrochemiluminescence analyzer “Elecsys 2010” (Roche Diagnostics, Germany). Bacterioscopic and bacteriological examination of the contents of the ampulla of the major duodenal papilla, obtained during endoscopic examination or external nasobiliary drainage, was carried out with determination of pathogen sensitivity to antibacterial agents.

Statistical analysis was performed using the software package “STATGRAPHICS Plus for Windows 7.0”.

Results and Discussion

The most common cause of obstruction of the terminal common bile duct in cholelithiasis was stenosis of the major duodenal papilla combined with choledocholithiasis, identified in 80 (36.2%) patients. Other causes included isolated stenosis of the major duodenal papilla in 64 (28.9%) patients, impacted stone of the major duodenal papilla in 33 (14.9%), choledocholithiasis in 26 (11.8%), chronic pseudotumorous pancreatitis in 16 (7.3%), and compression of the common bile duct by a cyst of the pancreatic head in 2 (0.9%) patients.

Obstructive jaundice syndrome was observed in 202 (91.4%) patients, acute cholangitis in 73 (33.1%), and acute biliary pancreatitis in 18 (8.1%). Previous

cholecystectomy had been performed in 35 (15.8%) patients. Concomitant somatic diseases were identified in 205 (92.8%) of 221 patients.

Of the 221 patients, open laparotomic operations were performed in 116 (52.5%), while endoscopic and/or laparoscopic interventions were performed in 105 (47.5%) patients. Open surgical procedures for obstruction of the terminal common bile duct included transduodenal or transcholedochal papillosphincterotomy with external drainage of the common bile duct or the formation of choledochoduodenostomy. According to indications, choledocholithotomy and cholecystectomy were also performed.

Endoscopic methods of correction included endoscopic papillosphincterotomy, endoscopic choledocholithotomy and/or stone extraction, less frequently endoscopic suprapapillary choledochoduodenostomy, as well as various options for drainage of the extrahepatic bile ducts (external transnasal, internal, or combined external-internal drainage). In 3 patients with obstruction of the terminal common bile duct associated with chronic pseudotumorous pancreatitis, endoscopic internal stenting was performed using an original technique.

Following endoscopic suprapapillary choledochoduodenostomy, tunneling of the narrowed intrapancreatic segment of the common bile duct was carried out by inserting an internal polyvinyl chloride stent up to 12 cm in length with side openings arranged in a staggered pattern into the lumen of the hepatic choledochus through the choledochoduodenostomy. This technique made it possible to avoid laparotomy and choledochotomy, general anesthesia, reduce surgical trauma and operative time, prevent suture failure of the common bile duct, and decrease postoperative mortality.

In 13 patients, including 9 with unresolved choledocholithiasis due to ineffective endoscopic stone extraction and high risk of open surgery, combined external-internal drainage of the extrahepatic bile ducts was performed to prevent recurrence of cholangitis and obstructive jaundice. After endoscopic papillosphincterotomy and, when possible, stone extraction, internal drainage of the common bile duct was performed with placement of the drain into the duodenal lumen, followed by external drainage through the nasal route. After resolution of cholangitis and obstructive jaundice, the external drain was removed, while the internal drain was left in place to ensure adequate bile outflow.

After normalization of clinical and laboratory parameters, laparoscopic cholecystectomy was performed as a second stage in 11 (12.9%) of 85 patients with gallbladder stones. In an additional 26 (30.6%) patients, cholecystectomy was performed by an open approach combined with choledocholithotomy.

In 48 (58.5%) of 82 patients, cholecystectomy was not performed due to high operative risk, and endoscopic intervention served as the definitive treatment. In 6 of these cases, endoscopic papillosphincterotomy with internal stenting constituted the final stage of treatment. Among 20 patients with prior cholecystectomy, endoscopic interventions were effective in 17 (85.0%). In 3

cases, treatment was completed with internal stenting due to невозможности эндоскопического удаления конкрементов и высокого риска открытой операции. In 3 patients, open choledocholithotomy was performed due to parapapillary diverticulum, large calculi, or невозможности катетеризации большого дуоденального сосочка.

Thus, in 9 (8.6%) of 105 patients treated with endoscopic methods, stones of the hepatic choledochus were not removed, and treatment was completed with endoscopic papillosphincterotomy and internal stenting.

The incidence of postoperative complications was significantly higher after open surgery compared to endoscopic interventions (χ^2=4.6; $p<0.05$) and amounted to 19.8% and 9.5%, respectively. Postoperative mortality was also significantly higher in the open surgery group (χ^2=4.9; $p<0.05$), amounting to 10.3% versus 2.9%.

Thus, the use of existing and newly developed endoscopic methods for the treatment of cholelithiasis complicated by obstruction of the terminal common bile duct made it possible to significantly reduce, in elderly and senile patients, the rate of postoperative complications from 19.8% to 9.5% and mortality from 10.3% to 2.9%.

Conclusions

1. In elderly and senile patients with obstruction of the terminal common bile duct and choledocholithiasis at high operative risk, endoscopic papillosphincterotomy with stone extraction is an adequate method of treatment.
2. In selected cases of unresolved choledocholithiasis, endoscopic papillosphincterotomy and internal stenting make it possible to avoid revision of the common bile duct and, in high-risk patients, to отказаться от открытого хирургического вмешательства.

A promising direction is the study of the combined use of endoscopic interventions and interventional sonography in obstruction of the terminal common bile duct complicated by acute purulent cholangitis and cholangiogenic liver abscesses.

References

1. Ivashkin VT, Maev IV. *Cholelithiasis*. Moscow: GEOTAR-Media; 2016.
2. Saveliev VS, Petukhov VA, Filimonov MI. Surgical treatment of complicated forms of cholelithiasis. *Khirurgiia (Mosk)*. 2013;(5):4–10.
3. Shalimov AA, Grubnik VV. *Surgery of the biliary tract*. Kyiv: Zdorovya; 2009.
4. Cotton PB, Leung JW. *Advanced Digestive Endoscopy: ERCP*. Oxford: Blackwell Publishing; 2006.

5. Freeman ML, Nelson DB, Sherman S, Haber GB, Herman ME, Dorsher PJ, et al. Complications of endoscopic biliary sphincterotomy. *N Engl J Med*. 1996;335(13):909–918.

6. Boerma D, Rauws EAJ, Keulemans YC, Janssen IMC, Bolwerk CJ, Timmer R, et al. Wait-and-see policy or laparoscopic cholecystectomy after endoscopic sphincterotomy for bile-duct stones. *Lancet*. 2002;360(9335):761–765.

7. Leung JW, Liu YL, Lau GC, Chan RC, Lai CW, Ling TK. Endoscopic biliary drainage for acute cholangitis in elderly patients. *Gastrointest Endosc*. 2000;51(3):301–306.

8. Costamagna G, Tringali A, Shah SK, Mutignani M, Perri V, Spada C. Long-term results of endoscopic management of bile duct stones in high-risk patients. *Endoscopy*. 2004;36(9):831–836.

ONE HEALTH APPROACH IN THE SURVEILLANCE OF EMERGING ZOONOTIC DISEASES IN SOUTHEAST ASIA

Dr. Nattapong Wattanakul Chiang
Mai University, Thailand

Kanya Srisuk
Chiang Mai University, Thailand

Abstract
Emerging zoonotic diseases represent a growing threat to global health security, particularly in Southeast Asia where dense populations, biodiversity, agricultural activities, and extensive human-animal interactions create favorable conditions for disease emergence. Outbreaks of diseases such as avian influenza, Nipah virus, and other zoonotic infections have demonstrated the importance of integrated surveillance systems capable of detecting health threats at an early stage. The One Health approach, which promotes collaboration among human health, animal health, and environmental sectors, has become a critical framework for addressing these challenges. Application of the One Health approach in the surveillance of emerging zoonotic diseases in Southeast Asia. The research evaluates collaborative surveillance mechanisms, datasharing systems, laboratory capacities, and multidisciplinary partnerships among public health agencies, veterinary services, and environmental organizations. The study also investigates regional efforts to strengthen preparedness and response capabilities for emerging infectious diseases. İntegrated surveillance systems significantly improve the early detection of zoonotic disease outbreaks and facilitate coordinated responses among relevant stakeholders. Crosssectoral collaboration enhances information exchange, risk assessment, and outbreak management. Furthermore, regional partnerships support knowledge sharing and strengthen public health capacities across national borders. However, challenges including limited financial resources, inconsistent surveillance standards, and insufficient interagency coordination continue to affect implementation effectiveness. One Health approach provides a valuable framework for strengthening disease surveillance and improving regional health security. Increased investment in surveillance infrastructure, professional training, and international cooperation is essential for preventing future zoonotic disease outbreaks and protecting public health. **Keywords**: One Health, Zoonotic Diseases, Disease Surveillance, Public Health, Veterinary Health, Emerging Infectious Diseases, Southeast Asia